

Nome: Gustavo Murilo Corvalante Carvalho
Turma: CT11 348

01. $A = (a_{ij})_{3 \times 2}$

$$a_{ij} = 2i + 3j$$

$$a_{11} = 2 \cdot 1 + 3 \cdot 1 = 5$$

$$a_{12} = 2 \cdot 1 + 3 \cdot 2 = 8$$

$$a_{21} = 2 \cdot 2 + 3 \cdot 1 = 7$$

$$a_{22} = 2 \cdot 2 + 3 \cdot 2 = 10$$

$$a_{31} = 2 \cdot 3 + 3 \cdot 1 = 9$$

$$a_{32} = 2 \cdot 3 + 3 \cdot 2 = 12$$

$$A = \begin{bmatrix} 5 & 8 \\ 7 & 10 \\ 9 & 12 \end{bmatrix}$$

02. $A = (a_{ij})_{2 \times 2}$

$$a_{ij} = i^2 + 4j^2$$

$$a_{11} = 1^2 + 4 \cdot 1^2 = 5$$

$$a_{12} = 1^2 + 4 \cdot 2^2 = 17$$

$$a_{21} = 2^2 + 4 \cdot 1^2 = 8$$

$$a_{22} = 2^2 + 4 \cdot 2^2 = 20$$

$$(A) \rightarrow A = \begin{bmatrix} 5 & 17 \\ 8 & 20 \end{bmatrix}$$

03. $X + 2 = -X \rightarrow 2X = -2 \rightarrow X = -1$

$$X = -1$$

$$2y = y - 1 \rightarrow y = -1$$

$$y = -1$$

$$Z + 1 = -2Z \rightarrow 3Z = -1 \rightarrow Z = -1/3$$

$$Z = -\frac{1}{3}$$

04. $3X = 2X + 1 \rightarrow X = 1$

$$X = 1$$


$$y = -X \rightarrow y = -1$$

$$y = -1$$

$$x = -2Z \rightarrow 2Z = -x \rightarrow 2Z = -1 \rightarrow Z = -\frac{1}{2}$$

$$Z = -\frac{1}{2}$$

05.



Para chegarmos a resposta exata a matriz deve ser definida pela seguinte lei:

$$a_{ij} = 0 \text{ se } i = j \quad \left\{ \begin{array}{l} a_{ij} = 1 \text{ se } i - j = 1 \text{ ou } -1 \\ a_{ij} = \sqrt{2} \text{ se } i - j = 2 \text{ ou } -2 \end{array} \right.$$

$$a_{ij} = \sqrt{2} \text{ se } i - j = 2 \text{ ou } -2$$

Assim chegamos na alternativa

$$(B) \begin{bmatrix} 0 & 1 & \sqrt{2} & 1 \\ 1 & 0 & 1 & \sqrt{2} \\ \sqrt{2} & 1 & 0 & 1 \\ 1 & \sqrt{2} & 1 & 0 \end{bmatrix}$$

06.

$$2A = \begin{bmatrix} -2 \\ 4 \\ 6 \end{bmatrix} - B = \begin{bmatrix} 0 \\ -2 \\ 1 \end{bmatrix} \rightarrow 2A - B = \begin{bmatrix} -2 \\ 6 \\ 5 \end{bmatrix} (D)$$

07.

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} - B^T = \begin{bmatrix} -1 & 2 \\ 3 & 0 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 4 \\ 3 & 5 \end{bmatrix} (B)$$

08.

$$\begin{bmatrix} 2 & -1 & 2y \\ x & 0 & -z \\ 4 & 3 & 2 \end{bmatrix} = \begin{bmatrix} 2 & x & 4 \\ -1 & 0 & 3 \\ 2y & -z & 2 \end{bmatrix}$$

$$x = -1$$

$$2y = 4 \rightarrow y = 2$$

$$-z = 3 \rightarrow z = -3$$

$$2 - 1 - 3 \rightarrow 2 - 4 = -2$$

$$(A) -2$$

$$09. A = (a_{ij})_{3 \times 2} \quad B = (b_{ij})_{3 \times 2}$$

$\text{se } i=j, a_{ij}=1$ $\text{se } i=j, b_{ij}=2i-j$
 $\text{se } i \neq j, a_{ij}=i+j$ $\text{se } i \neq j, b_{ij}=0$

$$A+B \rightarrow \begin{bmatrix} 1 & 3 \\ 3 & 1 \\ 4 & 5 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 2 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 3 & 3 \\ 4 & 5 \end{bmatrix} \quad (C)$$

$$10. \begin{bmatrix} \frac{3x}{2} & 12 \\ 15 & \frac{3y}{2} \end{bmatrix} + \begin{bmatrix} \frac{2y}{3} & 4 \\ 8 & \frac{2(x+4)}{3} \end{bmatrix} = \begin{bmatrix} 7 & 16 \\ 23 & 13 \end{bmatrix}$$

$$\frac{3x}{2} + \frac{2y}{3} = 7 \rightarrow 9x + 4y = 42$$

$$\frac{3y}{2} + \frac{2(x+4)}{3} = 13 \rightarrow 9y + 4x + 16 = 78 \rightarrow 9y + 4x = 62$$

$$\begin{array}{r} 9y + 4x = 62 \\ - 4y + 9x = 42 \\ \hline 5y - 5x = 20 \end{array}$$

$$\frac{5y - 5x = 20}{5} \rightarrow y - x = 4 \quad (B)$$