

# CHAPTER 7


# QUANTUM THEORY OF

# THE ATOM

Dr. Yuan Dan



# 7. Quantum Theory of the Atom




**FIGURE 7.1** ▲

## **Flame tests of Groups IA and IIA elements**

A wire loop containing a sample of a metal compound is placed in a flame. *Left to right:* flames of lithium (red), sodium (yellow), strontium (red), and calcium (orange).

# 7. Quantum Theory of the Atom

- Emission (line) spectra of some elements resolved (separated) by means of a prism





# 7. Quantum Theory of the Atom



SOLVAY CONFERENCE 1927

colourized by pastincolour.com

A. PICARD	E. HENRIOT	P. EHRENFEST	Ed. HERSEN	Th. DE DONDER	E. SCHRÖDINGER	E. VERSCHAFFELT	W. PAULI	W. HEISENBERG	R.H FOWLER	L. BRILLOUIN
P. DEBYE	M. KNUDSEN	W.L. BRAGG	H.A. KRAMERS	P.A.M. DIRAC	A.H. COMPTON	L. de BROGLIE	M. BORN	N. BOHR		
I. LANGMUIR	M. PLANCK	Mme CURIE	H.A. LORENTZ	A. EINSTEIN	P. LANGEVIN	Ch.E. GUYE	C.T.R. WILSON	O.W. RICHARDSON		


# 7.1 The Wave Nature of Light

---

- A **wave** is a continuously repeating change or oscillation in matter or in a physical field.
- **Light** is also a wave. It consists of oscillations in **electric** and **magnetic** fields that can travel through space.
- Visible lights, X-rays, and radio waves are all forms of **electromagnetic radiation**.


# 7.1 The Wave Nature of Light

- **Wavelength ( $\lambda$ )**: the distance between any two adjacent identical points of a wave.



# 7.1 The Wave Nature of Light

- **Frequency ( $\nu$ )**: the number of wavelengths of that wave that pass a fixed point in one unit of time (usually one second).
- Unit:  $\text{s}^{-1}$ , also *hertz* (Hz)



wavelength and frequency are **inversely related**: the greater the wavelength, the lower the frequency, and *vice versa*

# 7.1 The Wave Nature of Light

---

$$c = v\lambda$$

- The speed of light waves in vacuum is a constant:  
 $3.00 \times 10^8 \text{ m/s}$



## P266 Example 7.1

What is the wavelength of the yellow sodium emission, which has a frequency of  $5.09 \times 10^{14}/\text{s}$ ?

$$\lambda = \frac{3.00 \times 10^8 \text{ m/s}}{5.09 \times 10^{14}/\text{s}} = 5.89 \times 10^{-7} \text{ m, or } \mathbf{589 \text{ nm}}$$


## P266 Example 7.2

What is the frequency of violet light with a wavelength of 408 nm?

$$\nu = \frac{3.00 \times 10^8 \text{ m/s}}{408 \times 10^{-9} \text{ m}} = 7.35 \times 10^{14} / \text{s}$$

# 7.1 The Wave Nature of Light

- Electronic spectrum: the range of frequencies or wavelengths of electromagnetic radiation



# 7.2 Quantum Effects and Photons

---

- Light is...

Newton: a beam of particles


Young: like waves – diffraction

Einstein: light has both wave and particle properties (photoelectric effect)

- Diffraction is a property of waves in which the waves spread out when they encounter an obstruction or small hole about the size of the wavelength.

# 7.2 Quantum Effects and Photons

- Diffraction





# 7.2 Quantum Effects and Photons

- Planck's Quantization of Energy
  - An atom could have only certain energies of vibration,  $E$

$$E = nh\nu, \quad n = 1, 2, 3, \dots$$

- $h$ : Planck's constant, a physical constant relating energy and frequency, having the value  $6.63 \times 10^{-34} \text{ J}\cdot\text{s}$
- $n$ : quantum numbers
- The vibrational energies of the atoms are said to be **quantized**; that is, the possible energies are limited to certain values.

## 7.2 Quantum Effects and Photons


- Photoelectric Effect (by Einstein)
  - Light consists of photons, or particles of electromagnetic energy, with energy  $E$  proportional to the observed frequency of the light.

$$E = h\nu$$

- The wave-particle duality of light:  $E$  is the energy of a light particle or photon;  $\nu$  is the frequency of the associated wave.

# 7.2 Quantum Effects and Photons

- Photoelectric Effect (by Einstein)
  - Photoelectric effect: the ejection of electrons from the surface of a metal or from another material when light shines on it.
  - Electrons are ejected only when the frequency of light exceeds a certain threshold value characteristic of the particular metal.



**FIGURE 7.6** ▲  
**The photoelectric effect**


## P269 Example 7.3

The red spectral line of lithium occurs at 671 nm ( $6.71 \times 10^{-7}$  m). Calculate the energy of one photon of this light.

$$\nu = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{6.71 \times 10^{-7} \text{ m}} = 4.47 \times 10^{14} / \text{s}$$

## 7.3 The Bohr Theory of the Hydrogen Atom


- The light emitted by a heated solid. E.g., A heated tungsten filament in a lightbulb
- **Continuous spectrum**: a spectrum containing light of all wavelengths





# 7.3 The Bohr Theory of the Hydrogen Atom


- The light emitted by a heated gas
- **Line spectrum:** a spectrum showing only certain colors or specific wavelengths of light.



# 7.3 The Bohr Theory of the Hydrogen Atom

- The visible spectrum of hydrogen:

$$\frac{1}{\lambda} = 1.097 \times 10^7 / \text{m} \left( \frac{1}{2^2} - \frac{1}{n^2} \right)$$



# 7.3 The Bohr Theory of the Hydrogen Atom

---

- Bohr's Postulates
  - To solve problems of 1) the stability of the hydrogen atom (the atom exists and its electron does not continuously radiate energy and spiral into the nucleus); 2) the line spectrum of the atom.

# 7.3 The Bohr Theory of the Hydrogen Atom


- Bohr's Postulates
  - **Energy-level Postulate**: an electron can have only specific energy values in an atom.

$$E = -\frac{R_H}{n^2} \quad n = 1, 2, 3, \dots \infty \quad (\text{for H atom})$$

- $R_H = 2.179 \times 10^{-18} \text{ J}$
- $n$ : principle quantum number, integral values from 1

# 7.3 The Bohr Theory of the Hydrogen Atom

- Bohr's Postulates




**FIGURE 7.10** ▲  
Energy-level diagram for the  
electron in the hydrogen atom



# 7.3 The Bohr Theory of the Hydrogen Atom

- Bohr's Postulates
  - Transitions Between Energy Levels in an atom can change the energy of the electron from one energy level to another. By so doing, the electron can emit or absorb light.



**FIGURE 7.10** ▲  
Energy-level diagram for the  
electron in the hydrogen atom

# 7.3 The Bohr Theory of the Hydrogen Atom

- Bohr's Postulates
  - Transitions Between Energy Levels: An electron in an atom can change energy only by going from one energy level to another energy level. By so doing, the electron undergoes a transition.

$$\text{Energy of emitted photon} = h\nu = -\Delta E = -(E_f - E_i)$$

$$h\nu = R_H \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

## P273 Example 7.4

What is the wavelength of light emitted when the electron in a hydrogen atom undergoes a transition from energy level  $n = 4$  to level  $n = 2$ ?

$$E_i = \frac{-R_H}{4^2} = \frac{-R_H}{16} \quad \text{and} \quad E_f = \frac{-R_H}{2^2} = \frac{-R_H}{4}$$

$$\left( \frac{-R_H}{16} \right) - \left( \frac{-R_H}{4} \right) = \frac{-4R_H + 16R_H}{64} = \frac{-R_H + 4R_H}{16} = \frac{3R_H}{16} = h\nu$$

$$\nu = \frac{3R_H}{16h} = \frac{3}{16} \times \frac{2.179 \times 10^{-18} \text{ J}}{6.626 \times 10^{-34} \text{ J}\cdot\text{s}} = 6.17 \times 10^{14} / \text{s}$$

$$\lambda = \frac{3.00 \times 10^8 \text{ m/s}}{6.17 \times 10^{14} / \text{s}} = 4.86 \times 10^{-7} \text{ m, or } \mathbf{486 \text{ nm}}$$

## 7.4 Quantum Mechanics

- A theory that applies to submicroscopic (that is, extremely small) particles of matter, such as electrons.
- Stimulated by the discovery of the de Broglie relation.
- Particle of matter of mass  $m$  and speed  $v$  has an associated wavelength, by analogy with light:

$$\lambda = \frac{h}{mv}$$

## 7.4 Quantum Mechanics

- Particle of matter of mass  $m$  and speed  $v$  has an associated wavelength, by analogy with light:

$$\lambda = \frac{h}{mv}$$

- Baseball, 0.145 kg, 27 m/s,  $10^{-34}$  m
- Electron,  $9.11 \times 10^{-31}$  kg,  $10^{-12}$  m



## 7.4 Quantum Mechanics

- Electron microscope: To resolve detail the size of several hundred picometers, we need a wavelength on that order.



## P278 Example 7.5

- Calculate the wavelength (in meters) of the wave associated with a 1.00-kg mass moving at 1.00 km/hr.
- What is the wavelength (in picometers) associated with an electron, whose mass is  $9.11 \times 10^{-31}$  kg, traveling at a speed of  $4.19 \times 10^6$  m/s? (This speed can be attained by an electron accelerated between two charged plates differing by 50.0 volts; voltages in the kilovolt range are used in electron microscopes.)

$$1.00 \frac{\text{km}}{\text{hr}} \times \frac{1 \text{ hr}}{3600 \text{ s}} \times \frac{10^3 \text{ m}}{1 \text{ km}} = 0.278 \text{ m/s}$$

$$\lambda = \frac{h}{mv} = \frac{6.63 \times 10^{-34} \text{ J}\cdot\text{s}}{1.00 \text{ kg} \times 0.278 \text{ m/s}} = \mathbf{2.38 \times 10^{-33} \text{ m}}$$

## P278 Example 7.5

- Calculate the wavelength (in meters) of the wave associated with a 1.00-kg mass moving at 1.00 km/hr.
- What is the wavelength (in picometers) associated with an electron, whose mass is  $9.11 \times 10^{-31}$  kg, traveling at a speed of  $4.19 \times 10^6$  m/s? (This speed can be attained by an electron accelerated between two charged plates differing by 50.0 volts; voltages in the kilovolt range are used in electron microscopes.)

$$\lambda = \frac{h}{mv} = \frac{6.63 \times 10^{-34} \text{ J}\cdot\text{s}}{9.11 \times 10^{-31} \text{ kg} \times 4.19 \times 10^6 \text{ m/s}} = 1.74 \times 10^{-10} \text{ m} = \mathbf{174 \text{ pm}}$$

## 7.4 Quantum Mechanics

---

- de Broglie relation: applies quantitatively only to particles in a force-free environment
- Erwin Schrödinger's theory: could be used to find the wave properties of electrons in atoms and molecules.
- Quantum mechanics or wave mechanics:  
The branch of physics that mathematically describes the wave properties of submicroscopic particles.



## 7.4 Quantum Mechanics

- Heisenberg pointed out: it is impossible to know simultaneously, with absolute precision, both the position and the momentum of a particle such as an electron.
- Heisenberg's uncertainty principle: a relation that states that the product of the uncertainty in position and the uncertainty in momentum of a particle can be no smaller than Planck's constant divided by  $4\pi$ .


$$(\Delta x)(\Delta p_x) \geq \frac{h}{4\pi} \longrightarrow (\Delta x)(\Delta v_x) \geq \frac{h}{4\pi m}$$

# 7.4 Quantum Mechanics

- Make statistical statements on the probability of finding an electron at a certain point in a hydrogen atom.



A



B

## 7.5 Quantum Numbers and Atomic Orbitals

- According to quantum mechanics, each electron in an atom is described by four different quantum numbers, three of which ( $n$ ,  $l$ , and  $m_l$ ) specify the wave function that gives the probability of finding the electron at various points in space.
- A wave function for an electron in an atom is called an atomic orbital.
- An atomic orbital is pictured qualitatively by describing the region of space where there is high probability of finding the electrons.

## 7.5 Quantum Numbers and Atomic Orbitals

---

- **Principal Quantum Number ( $n$ ):** The energy of an electron in an atom depends principally on  $n$ .
- $n = 1, 2, 3$ , and so on.
- The size of an orbital also depends on  $n$ . The larger the value of  $n$  is, the larger the orbital.




## 7.5 Quantum Numbers and Atomic Orbitals

- **Angular Momentum Quantum Number ( $l$ ):**  
This quantum number distinguishes orbitals of given  $n$  having different shapes; it can have any integer value from 0 to  $n-1$ .
- For a given  $n$ , the energy of an orbital increases with  $l$ .
- Orbitals of the same  $n$  but different  $l$  are said to belong to different subshells of a given shell.

<i>Letter</i>	<i>s</i>	<i>p</i>	<i>d</i>	<i>f</i>	<i>g</i> . . .
<i>l</i>	0	1	2	3	4 . . .

# 7.5 Quantum Numbers and Atomic Orbitals



## 7.5 Quantum Numbers and Atomic Orbitals

- **Magnetic Quantum Number ( $m_l$ ):** This quantum number distinguishes orbitals of given  $n$  and  $l$  - that is, of given energy and shape but having a different orientation in space; the allowed values are the integers from  $-l$  to  $+l$ .
- The orbitals have the same shape but different orientations in space.
- All orbitals of a given subshell have the same energy.
- There are  $2l+1$  orbitals in each subshell of quantum number  $l$ .

## 7.5 Quantum Numbers and Atomic Orbitals

---

- A fourth quantum number ( $m_s$ ) refers to a magnetic property of electrons called **spin**.
- This quantum number refers to the two possible orientations of the spin axis of an electron; possible values are  $+1/2$  and  $-1/2$ .

# 7.5 Quantum Numbers and Atomic Orbitals

**TABLE 7.1**

**Permissible Values of Quantum Numbers for Atomic Orbitals**


$n$	$l$	$m_l^*$	Subshell Notation	Number of Orbitals in the Subshell
1	0	0	1s	1
2	0	0	2s	1
2	1	-1, 0, +1	2p	3
3	0	0	3s	1
3	1	-1, 0, +1	3p	3
3	2	-2, -1, 0, +1, +2	3d	5
4	0	0	4s	1
4	1	-1, 0, +1	4p	3
4	2	-2, -1, 0, +1, +2	4d	5
4	3	-3, -2, -1, 0, +1, +2, +3	4f	7

\*Any one of the  $m_l$  quantum numbers may be associated with the  $n$  and  $l$  quantum numbers on the same line.

# 7.5 Quantum Numbers and Atomic Orbitals


- Atomic Orbital Shapes

99% contour



**1s orbital**


99% contour




**2s orbital**

# 7.5 Quantum Numbers and Atomic Orbitals


- Atomic Orbital Shapes



$2p_x$  orbital



$2p_y$  orbital



$2p_z$  orbital

# 7.5 Quantum Numbers and Atomic Orbitals

- Atomic Orbital Shapes

