

Discrete and Algorithmic Geometry 2013(Part 2)

Julian Pfeifle

Version of January 8, 2014

This is the preliminary version of the lecture notes for the second part of *Discrete and Algorithmic Geometry* (Universitat Politècnica de Catalunya), held in the fall semester of 2013 by Ferran Hurtado and Julian Pfeifle.

These notes are fruit of the collaborative effort of all participating students, who have taken turns in assembling this text. The name of each scribe figures in each corresponding section.

Suggestions for improvements will always be gladly received by `julian.pfeifle@upc.edu`.

Contents

Lecture 1. Convex Polytopes	5
1. Faces	5
Lecture 2. Asymptotic f-vectors of families of polytopes	9
1. The Unimodality Conjecture	9
2. Operations on polytopes	11
Lecture 3. Cyclic Polytopes	13
Lecture 4. Kalai's simple way of telling a simple polytope from its graph	15
Lecture 5. Lattice points in multiples of polytopes	17
1. Generating functions for rational cones	17
Lecture 6. Ehrhart-Macdonald Reciprocity	19
1. Statement of the theorem	19
2. Auxiliary results	19
3. Proof of Ehrhart-Macdonald Reciprocity	20
4. Degree of a lattice polytope	20
5. Reflexive polytopes	21
Scribes 2013	23
1. Alex Alvarez	23
2. Cecilia Girón Albert	23
3. Anna Somoza	23
4. Daniel Torres	23
5. Borja Elizalde	24
6. Xavier Tapia	24
Bibliography	25
Papers to referee	27
Bibliography	27

LECTURE 1

Convex Polytopes

Scribe: Cecilia Girón

A convex polytope can be defined in two different ways:

- *V-polytope* (discrete geometry): is the convex hull of the finite non-empty point set in \mathbf{R}^d .
- *H-polytope* (linear/integer optimization): An *H-polyhedron* is an intersection of a finite number of linear half spaces in some \mathbf{R}^d , if non-empty. And an *H-polytope* is a bounded *H-polyhedron*.

1. Faces

One of the properties studied about polytopes is their faces. A **face** F of a polytope P is a set of the form:

$$F = \{x \in \mathbf{R}^d : \langle a, x \rangle = n\} \cap P$$

where $a \in (\mathbf{R}^d)^*$ (dual space), $b \in \mathbb{R}$ and $P \subseteq \{x \in \mathbf{R}^d : \langle a, x \rangle \leq b\} \iff$ The inequality $\langle a, x \rangle \leq b$ is valid for P . Notice that P is actually a face of itself.

We can also study the dimension $\dim F$ of a face. Let P be a d dimensional polytope, then if a face F of P has dimension:

- $d - 1$, it is called a **facet**.
- $d - 2$, it is called a **ridge**.
- 1, it is called an **edge**.
- 0, it is called a **vertex**.
- -1 then $F = \emptyset$.


The partially ordered set of all faces $\mathcal{F}(P)$ of a convex polytope P forms a Eulerian lattice called **face lattice**. The face lattice can be used, for instance, to count the number of faces of same dimension:

$$f_i = \#\{\text{faces } F \text{ of } P \text{ with } \dim F = i\}$$

EXAMPLE. Let P be a cube in two dimensions, so a square. Notice that $\dim P = 2$, for every edge $ij \in P$ $\dim(ij) = 1$ and for every vertex i $\dim i = 0$, for $i, j = 1, 2, 3, 4$ and $i \neq j$. With this information we can construct the face lattice and study some properties of P .



EXAMPLE. Let's study now the dimension of the faces of a hypercube of dimension d $\square^d = \{x \in \mathbf{R}^d : -1 \leq x_i \leq 1, i = 1, \dots, d\}$: $f_{-1}(\square^d) = 1$, $f_0(\square^d) = 2^d$, $f_{d-1}(\square^d) = 2d$ and $f_d(\square^d) = 1$. Notice that the radius r from the center of the cube to one of its vertices is $r = \sqrt{d}/2$, thus the exterior circle of the polytope has radius r and the interior radius 1.



d	2	3	4	5	...	100	...	10^{100}
$r = \sqrt{d} - 1$	$\sqrt{2} - 1$	$\sqrt{3} - 1$	1	$\sqrt{5} - 1$...	9	...	$10^{50} - 1$
f_0	4	8	16	32	...	2^{100}	...	$2^{10^{100}}$
f_1	4	6	8	10	...	200	...	$2 \cdot 10^{100}$



An other property that can be studied about the faces of convex polytopes is whether they are a simplex or not. Let P be a polytope such that $\dim P = d$ and $\mathcal{F}(p) = k + 1$. It is said to be **simplicial** if it is k -simplex, i.e. if each of its faces is a simplex; and it is called **simple** if each of its vertices is contained in exactly d faces where $\dim P = d$.

1.1. Exercises done during the lecture 8/11/2013. Each one includes one.

Alex Alvarez.: The set of vertices is the following:

$(1 : -1 : -1 : -1 : 0 : 0)$
 $(1 : 1 : -1 : -1 : 0 : 0)$
 $(1 : -1 : 1 : -1 : 0 : 0)$
 $(1 : 1 : 1 : -1 : 0 : 0)$
 $(1 : 0 : 0 : 1 : 0 : 0)$
 $(1 : 0 : 0 : 1 : 1 : 0)$
 $(1 : 0 : 0 : 1 : 0 : 1)$

And this can be obtained as the join of a 2-cube with a 2-simplex. If we construct that in Polymake:

```
polytope > $p = join_polytopes(cube(2), simplex(2));
```

```
polytope > print($p->VERTICES);
```

```
1 -1 -1 -1 0 0
1 1 -1 -1 0 0
1 -1 1 -1 0 0
1 1 1 -1 0 0
1 0 0 1 0 0
1 0 0 1 1 0
1 0 0 1 0 1
```

Thus, we can use the program to see the number of facets and the vertices in each facet:

```
polytope > print $p->N_FACETS;
```

```
polymake: used package cddlib
```

Implementation of the double description method of Motzkin et al.

Copyright by Komei Fukuda.

http://www.ifor.math.ethz.ch/~fukuda/cdd_home/cdd.html

```
polymake: used package lrslib
```

Implementation of the reverse search algorithm of Avis and Fukuda.

Copyright by David Avis.

<http://cgm.cs.mcgill.ca/~avis/lrs.html>

```
7
polytope > print $p->POINTS_IN_FACETS;
{0 2 4 5 6}
{0 1 4 5 6}
{1 3 4 5 6}
{2 3 4 5 6}
{0 1 2 3 5 6}
{0 1 2 3 4 6}
{0 1 2 3 4 5}
```

If we focus now in the graph of the polytope, we can check the number of edges and we can also see it:

```
polytope > print $p->GRAPH->N_NODES;
```

```
7
```

```
polytope > print $p->GRAPH->N_EDGES;
```

```
19
```

```
polytope > print $p->GRAPH->VISUAL;
```

Therefore, we can see that the last two vertices are not connected, but they are connected to all the other vertices.

Exercise ?? (team members):

LECTURE 2

Asymptotic f-vectors of families of polytopes

Scribe: Cecilia Girón

1. The Unimodality Conjecture

In this section we are going to study the *unimodality conjecture* which says that there exists an $l = P(L) \in \mathbb{N}$ such that $f_0 \leq f_1 \leq \dots \leq f_l \leq f_{l+1} \leq \dots \leq f_{d-1} \leq f_d$. We would like to know if it is true.

First, we define the **f-vector** as the vector of the form $(f_0, f_1, \dots, f_{d-1})$ where f_i is as defined before in (1). We will say that it is a **flag f-vector** $(f_s)_s = [d]$ such that f_s count the number of flags $F_{i_1} \subset F_{i_2} \subset \dots \subset F_{i_k}$ where $s = \{i_1, i_2, \dots, i_k\}$ and $\dim F_{i_k} = i_k$ ¹.

The unimodal conjecture described before is known to be false for simplicial polytopes of dimension $d \geq 19$ and for non-simplicial polytopes of dimension $d \geq 8$. The following conjecture is not known to be false.

Restricted unimodal conjecture (Anders Björner):

$$f_0 \leq f_1 \leq \dots \leq f_{\lfloor \frac{d-1}{4} \rfloor} \\ f_{\lfloor \frac{3(d-1)}{4} \rfloor} \geq \dots \geq f_{d-1}$$

Intuitively we are sure that there is no way this conjecture could be false, but there is not proof of this. We don't even know if $f_k \geq \frac{1}{10000} \min\{f_0, f_{d-1}\}$ is true.

1.1. Exercises worked on during the lecture 11/11/2013. Each team includes one.

Exercise 1b (Alex Alvarez and Ivan Geffner): Using the simple form $n! \approx \left(\frac{n}{e}\right)^n$ of Stirling's formula, show that $\phi_d(x) := \log\left(\frac{d}{xd}\right)$ is asymptotically proportional to $-x \log x - (1-x) \log(1-x)$, for $x \in (0, 1)$ and $d \rightarrow \infty$. Discuss the real function ϕ_d on $[0, 1]$.

Using the simple form of Stirling's formula, we obtain the following form:

$$\binom{d}{xd} \approx \frac{d^d}{xd^{xd}(d-xd)^{d-xd}}$$

So applying the logarithm, we get

$$\begin{aligned} \phi_d(x) &= d \log d - xd \log xd - (d - xd) \log(d - xd) \\ &= d \log d - xd \log x - xd \log d - (d - xd)(\log d + \log(1 - x)) \\ &= d(-x \log x - (1 - x) \log(1 - x)) \end{aligned}$$

¹You can also read about *cd-index*

Thus, the first part of the exercise is proven. Let us study now the shape of the function.

The function clearly vanishes when x tends to 0 and 1 and in this interval is non-negative. The first derivative of the function is

$$\frac{d}{dx}(-(1-x)\log(1-x) - x\log(x)) = \log(1-x) - \log(x)$$

and therefore, there is only one point in which derivative vanishes, that is $x = \frac{1}{2}$. Now, the second derivative is

$$\frac{d^2}{dx^2} \log(1-x) - \log(x) = \frac{1}{x(x-1)}$$

We can conclude then that the point is a maximum, so we have characterized the shape of the function.

Exercise 2b (Borja and Cecilia): Using the simple form $n! \approx \left(\frac{n}{e}\right)^n$ of Stirling's formula, show that $\psi_d(x) := d(1-x) + \log \binom{d}{xd}$ is asymptotically proportional to $1 - x - x \log x - (1-x) \log(1-x)$, where $\log = \log_2$ denotes the binary logarithm. Find an approximation to the maximum of this function on $(0, 1)$.

For the first part of the exercise, by applying the Stirling's formula, in the binomial for of the given function:

$$\binom{d}{xd} = \left(\frac{d}{xd^x(d(1-x))^{1-x}} \right)^d$$

Then, substituting in $\psi_d(x)$:

$$\begin{aligned} \psi_d(x) &= d(1-x) + \log \left(\frac{d}{xd^x(d(1-x))^{1-x}} \right)^d \\ &= d(1-x) + d(\log d - x \log x - x \log d - (1-x) \log d - (1-x) \log(1-x)) \\ &= d(1-x - x \log x - (1-x) \log(1-x)) \end{aligned}$$

Hence, $\psi_d(x)$ is asymptotically proportional to $1 - x - x \log x - (1-x) \log(1-x)$

For the second part of the exercise, in order to find the maximum of the function

$$f(x) = 1 - x - x \log(x) - (1-x) \log(1-x)$$

in $(0,1)$ we will the points that have first derivative equal to 0, that is x such that $f'(x) = 0$, and computing $f'(x)$ we get:

$$f'(x) = -1 - (\log x + 1) - (-\log(1-x) - 1) = \log\left(\frac{1-x}{x}\right) - 1$$

Now the points that make $f'(x) = 0$ are the ones that make $\log\left(\frac{1-x}{x}\right) = 1$, which is the same as x such that $\frac{1-x}{x} = e$, which translates into:

$$x_{\max} = \frac{1}{e+1}$$

The shape of this function is a growing function from $x = 0$ starting at $f(0) = 1$ to $x = x_{\max}$, where it gets it's maximum, that is approximated by $f(x_{\max}) \approx 1.0414$ and then decreases to 0 at $x = 1$.

2. Operations on polytopes

- **Cartesian (direct) product** $P \times Q$.
- **Direct sum** $P^d \oplus Q^e \subset \mathbb{R}^{d+e}$.
- $P * Q \subset \mathbb{R}^{d+e+1}$. It is like \oplus but the subspaces are skew (i.e. affine and they have no point on common). For example $\square^1 * \square^1 = Pyr(P)$.

EXAMPLE: Given $f_k(P)$, calculate the k -th entry of $Pyr(P)$:

$$\begin{aligned} f(P) &= (f_0, f_1, \dots, f_{d-1}) \\ f_k(Pyr(P)) &= (f_0 + 1, f_1 + f_0, f_2 + f_1, \dots, f_{d-2} + f_{d-3}, f_{d-1} + f_{d-2}, 1 + f_{d-1}) \end{aligned}$$



- **Connected sum** $P^d \# Q^d$ where P has as simplicial face f and Q has a simplicial face G .

This last operation is used to join the asymptotic function $f(\square^d)$ and its dual $f(\diamond^d)$. To make it work, since \square^d has no triangulations in its faces, it is enough to cut away one vertex and, this way, get a simplex. Merging both functions using the connected sum gives place to a new function which is a non-unimodal function.

LECTURE 3

Cyclic Polytopes

Scribe: Daniel Torres

Let $x: \mathbb{R} \rightarrow \mathbb{R}^d$ an algebraic curve. Then we can choose $n > d$ points of the curve, for example, for $t_1 < \dots < t_n$ choosing $x(t_1), \dots, x(t_n)$, in order to construct a polytope resulting convex hull of these points. An example of this could be the degree d *moment curve* $\mu_d(t) = (t, t^2, \dots, t^d)$.

At first, one can think that this way to construct a polytope it's just silly example, since choosing random points in a algebraic curve seems to be the same that choosing random points in \mathbb{R}^d . However, an algebraic curve of degree d relating vertices let us obtain an interesting result.

Theorem 3.1. *Let μ be a degree d algebraic curve in \mathbb{R}^d . Then, the combinatorial type of the polytope $\text{conv}\{\mu(t_1), \dots, \mu(t_n)\}$ is independent of the choice of t_1, \dots, t_n .*

LECTURE 4

Kalai's simple way of telling a simple polytope from its graph

Scribe: Alex Alvarez

Conjecture 4.1 (Micha Perles, ~ 1970). *Any d -polytope is determined by its graph, i.e., $sk^1(P)$ determines $\mathcal{F}(P)$.*

The conjecture is false, e.g. for simplicial polytopes. On the other hand, K_n is dimensionally ambiguous, e.g. $sk^1(C_d(n)) = sk^1(C_e(n))$ for $d, e < n$. Besides there are neighborly graphs that are not combinatorially isomorphic to cyclic ones. However, the result is true for simple polytopes:

Theorem 4.2 (Roswitha Blind, Peter Mani, 1987). *Any simple d -polytope is determined by its graph.*

Their proof is not constructive, but in 1988 Kalai found an algorithmic approach. Let us see the details of that approach.

Definition 4.3 (Acyclic orientation). An acyclic orientation of a graph G is an orientation of $E(G)$ without directed cycles.

Definition 4.4 (Abstract Objective Function, AOF). An AOF of $G = sk^1(P)$ is an acyclic orientation that has a unique sink on each face of P .

Definition 4.5 (Unique Sink Orientation). A Unique Sink Orientation is an orientation such that every face has a unique sink. The orientation is not necessarily acyclic.

Notice that AOFs exist because of the existence of generic linear objective functions. Consider the set of all AOFs and define the following function:

$$(4.1) \quad f(O) = h_0(O) + 2h_1(O) + \dots + 2^k h_k(O) + \dots + 2^d h_d(O)$$

where O is an AOF and $h_k(O) \in \mathbb{N}$ counts the vertices of in-degree k in O .

Let f denote the number of non-empty faces of P . Given an acyclic orientation O , then

- (1) $f(O) \geq f$, because as O is acyclic each face has at least one sink.
- (2) $f(O) = f$ if O is an AOF.

To determine all AOFs of P just from its graph $G = sk^1(P)$ do the following steps:

- (1) Enumerate all acyclic orientations O of G .
- (2) Calculate $f(O)$ for each such orientation O .
- (3) Keep the orientations O with minimal $f(O)$.

The only remaining step to complete Kalai's method is showing a way to identify the faces of P from the knowledge of all the AOFs. Notice that the subgraphs of G that are graphs of faces are connected, k -regular and induced. With that, the following proposition is enough to characterize the faces:




FIGURE 1. An example of AOF. For this orientation, $h_0 = 1$, $h_1 = 3$, $h_2 = 3$ and $h_3 = 1$. Therefore, $f(O) = 1 + 6 + 12 + 8 = 27 = 3^3$.

Proposition 4.6. *An induced connected k -regular subgraph H is the graph of a face of $P \Leftrightarrow H$ is an initial set for some AOF O .*

PROOF. \Rightarrow : Perturb a face-defining inequality to obtain a linear function with respect to which the vertices of F lie below all other vertices.

\Leftarrow : Let H be a connected k -regular subgraph of $sk^1(P)$ and let O be an AOF such that H is an initial set of O . As H has a unique sink v and is k -regular, the sink v has k incoming edges and as the polytope is simple, those k edges determine a k -face F and v is a sink in that face.

As v is the unique sink of O in F , all vertices of F are $\leq v$ with respect to O . But the initial set of v in G is H , so $\text{vert}(F) \subseteq \text{vert}(H)$. As both H and F are connected and k -regular, we have that $\text{vert}(F) = \text{vert}(H)$. \square

Notice that this algorithm is exponential, but in 1995 Friedman gave a polynomial algorithm to reconstruct the 2-faces, which had been proven before to be enough to reconstruct the whole polytope.

LECTURE 5

Lattice points in multiples of politopes

Scribe: Xavier Tapia

Let's define $L_P(t) = \# \{tP^d \cap \mathbb{Z}^d\}$, this number is equal to $\# \{P^d \cap \frac{1}{t}\mathbb{Z}^d\}$ where P is a polytope. Take now $P^o = \text{int}P$ as a topological space, and consider $L_{P^o}(t) = \# \{t(P^o)^d \cap \mathbb{Z}^d\}$.

Let's consider first some examples of this:

If we take $d = 2$ and $t = 2$ we will have 9 points

Consider some tables.

	0	1	2	3	4	...	t
$L_{\square^2}(t)$	1	4	9	16	25	...	$(t+1)^2$
$L_{(\square^2)^o}(t)$	0	0	1	4	9	...	$(t-1)^2$
$L_{\square^1}(t)$	1	2	3	4	5	...	$t+1$
$L_{(\square^1)^o}(t)$	0	0	1	2	3	...	$t-1$

It's possible to see that, in general:

Theorem 5.1. $L_{\square^d}(t) = (t+1)^d$, $L_{(\square^d)^o}(t) = (t-1)^d = (-1)^d L_{\square^d}(-t)$

In fact, this result is true for every polytope P not only for \square^d , this theorem is known as the Ehrhart-Macdonald Reciprocity and it's the principal result we want to prove.

Theorem 5.2 (Ehrhart-Macdonald Reciprocity).

$$L_P(-t) = (-1)^d L_{P^o}(t)$$

To proof this theorem we need some previous results,

1. Generating functions for rational cones

Take S a rational cone or polytope (i.e. the generators of 1-d rays with rational vertices coordinates). We define:

$$G_S(z) = G_S(z_1, z_2, \dots, z_d) = \sum_{m \in S \cap \mathbb{Z}^d} z^m = \sum_{m \in S \cap \mathbb{Z}^d} z_1^{m_1} \dots z_d^{m_d}$$

$$G(z) = 1 + x + y + xy = (1+x)(1+y)$$

Goal: Find $G_C(z)$ where $C = \{v_1, \dots, v_d\} \subset$

\mathbb{R}^d

Assume that C is simplicial (that means C is a cone over a simplex), so the parallelepiped $\Pi_C = \{\lambda_1 v_1 + \dots + \lambda_d v_d : 0 \leq \lambda_i \leq 1\}$ till C disjointly.

In particular, $G_C(z) = \sum_{s_i \in \mathbb{N}} \text{tr}_{\sum s_i v_i} G_{\Pi}(z) = \sum_{s_i \in \mathbb{N}} z^{\sum s_i v_i} G_{\Pi}(z) = \frac{G_{\Pi}(z)}{(1-z^{v_1}) \dots (1-z^{v_d})} \in \mathbb{Q}[[z_1, \dots, z_d]]$

Theorem 5.3 (Beck-Robins theorem 3.5). *Suppose $C = \text{cone}\{v_1, \dots, v_d\}$ cone over a simplex, then for all $v \in \mathbb{R}^d$ the generating function satisfies:*

$$G_{v+C} = \frac{G_{v+\Pi_C}(z)}{(1-z^{v_1})\dots(1-z^{v_d})}$$

PROOF. In $G_{v+C}(z) = \sum_{m \in (v+C) \cap \mathbb{Z}^d} z^m$ write $m = w + \lambda_1 v_1 + \dots + \lambda_d v_d$ for $\lambda_i \geq 0$, and take $\lambda_k = \lfloor \lambda_k \rfloor + \{\lambda_k\}$ so for that, we can write:

$$m = w + \lfloor \lambda_1 \rfloor v_1 + \dots + \lfloor \lambda_d \rfloor v_d + \{\lambda_1\} v_1 + \dots + \{\lambda_d\} v_d$$

The vector $P = w + \{\lambda_1\} v_1 + \dots + \{\lambda_d\} v_d \in \mathbb{Z}^d$ is unique and any $u \in v + C \cap \mathbb{Z}^d$ can be written uniquely as $P + \sum n_i v_i$ with $P \in \Pi_C$ and $n_i \in \mathbb{Z}$.

We have:

$$\frac{\sigma_{w+\Pi_C}(z)}{(1-z^{v_1})\dots(1-z^{v_d})} = \left(\sum_{m \in (v+C) \cap \mathbb{Z}^d + w} z^m \right) \left(\sum_{k_i \in \mathbb{N}} z^{k_1 v_1} \right) \dots \left(\sum_{k_d \in \mathbb{N}} z^{k_d v_d} \right)$$

Note that if $\Delta C \cap \mathbb{Z}^d = \emptyset$ it suffices to take $\Pi_C^0 = \{\sum_{i=1}^d \lambda_i v_i, 0 < \lambda_i < 1\}$ the open fundamental parallelepiped. \square

Theorem 5.4. *For any pointed cone (it's maximum linear subspace contained in C is $\{0\}$) $C = \{w + \sum \lambda_i v_i, \lambda_i \geq 0\}$ with $w \in \mathbb{R}^d$ and $v_i \in \mathbb{Z}^d$, the generating function $\sigma_C(z)$ is a rational function in the z_i .*

LECTURE 6

Ehrhart-Macdonald Reciprocity

Scribe: Albert Santiago

1. Statement of the theorem

Let P be a lattice polytope. Let $L_P : \mathbb{N} \longrightarrow \mathbb{N}$ be the lattice point counting function,

$$\begin{aligned} L_P : \mathbb{N} &\longrightarrow \mathbb{N} \\ t &\longmapsto L_P(t) = \# \{tP \cap \mathbb{Z}^d\} \end{aligned}$$

and let $L_{P^0} : \mathbb{N} \longrightarrow \mathbb{N}$ be the lattice point counting function in the interior of P .

By Ehrhart's Theorem seen in the previous lecture, it is known that $L_P, L_{P^0} \in \mathbb{Q}[t]$.

Theorem 6.1 (Ehrhart-Macdonald Reciprocity).

$$L_P(-t) = (-1)^d L_{P^0}(t)$$

Note that in previous lecture we have already seen an example for this EM reciprocity for the cube. Indeed,

$$\begin{aligned} L_{\square^d}(t) &= (1+t)^d \\ L_{(\square^d)^0}(t) &= (-1)^d (1-t)^d \end{aligned}$$

2. Auxiliary results

Proposition 6.2. *Let $w_1, \dots, w_d \in \mathbb{Z}^d$ be linearly independent. Let $C = \text{cone } w_1, \dots, w_d$ be cone over a simplex. Let $v \in \mathbb{R}^d$ such that $\partial(v+C) \cap \mathbb{Z}^d = \emptyset$. Then,*

$$\sigma_{v+C} \left(\frac{1}{z_1}, \dots, \frac{1}{z_d} \right) = (-1)^d \sigma_{-v+C}(z_1, \dots, z_d)$$

Notes:

- For the existence of such $v \in \mathbb{R}^d$, see exercise sheet #6.
- $\sigma_{v+C} \left(\frac{1}{z_1}, \dots, \frac{1}{z_d} \right)$ is not a power series, so the theorem is meaningless at this level. At most, we can say it is an element from $\mathbb{Q}[z_1, \dots, z_d]$, where the theorem holds.
- Try to see $\frac{1}{z_i}$ as the variables corresponding to $-P$.

PROOF. To be done. □

Proposition 6.3 (Stanley Reciprocity). *Let C be a rational pointed cone (i. e. $C = \text{cone } w_1, \dots, w_d, w_i \in \mathbb{Q}^d$) with apex at the origin. Then,*

$$\sigma_C \left(\frac{1}{z} \right) = (-1)^d \sigma_{C^0}(z)$$

PROOF. To be done. □

Definition 6.4. We define the Ehrhart function for the interior lattice points of a polytope naturally as

$$\text{Ehr}_{P^0}(z) := \sum_{t \geq 1} L_{P^0}(t) z^t$$

Note: We can start the sum at $t = 1$ rather than at $t = 0$ because the origin is not an interior point of P .

Observation 6.5. As it happened with the original Ehrhart function, the following equality holds:

$$\text{Ehr}_{P^0}(z) = \sigma_{\text{cone}(P^0)}(1, \dots, 1, z)$$

Proposition 6.6. Let P be a lattice polytope. Then,

$$\text{Ehr}_{(P^0)}(z) = (-1)^{\dim P+1} \text{Ehr}_P\left(\frac{1}{z}\right)$$

Notes:

- The proposition is also valid for P any *rational* polytope, not only for lattice polytopes.
- Not every polytope can be perturbed into a rational one. There exist irrational polytopes.

PROOF. To be done. □

3. Proof of Ehrhart-Macdonald Reciprocity

To be done.

4. Degree of a lattice polytope

Observation 6.7. Let $p \in \mathbb{Q}[t]$ be a polynomial of degree d . If the following equality holds:

$$\sum_{t \geq 0} p(t) z^t = \frac{h_d z^d + \dots h_1 z + h_0}{(1 - z)^d}$$

then

$$\left. \begin{array}{l} h_d = \dots = h_{k+1} = 0 \\ h_k \neq 0 \end{array} \right\} \iff \left\{ \begin{array}{l} p(-1) = p(-2) = \dots = p(-(d-k)) = 0 \\ p(-(d-k+1)) \neq 0 \end{array} \right.$$

PROOF. See Theorem 3.18 at Beck-Robins' *Computing the continuous discretely*. □

Proposition 6.8. Let $P \subset \mathbb{R}^d$ be a lattice polytope with Ehrhart function:

$$\text{Ehr}_P(z) = \frac{h_d z^d + \dots h_1 z + h_0}{(1 - z)^d}$$

Then,

$$\left. \begin{array}{l} h_d = \dots = h_{k+1} = 0 \\ h_k \neq 0 \end{array} \right\} \iff \left\{ \begin{array}{l} L_P(-1) = L_P(-2) = \dots = L_P(-(d-k)) = 0 \\ L_P(-(d-k+1)) \neq 0 \end{array} \right.$$

Definition 6.9. We call k the *degree of the lattice polytope* P .

Observation 6.10. If P is a d -polytope of degree k , then $L_P(-(d-k)) = (-1)^d L_{P^0}(-(d-k)) = 0$, so the $(d-k)$ -th dilate of P does not contain any interior lattice point, and $(d-k+1)P$ is the smallest integer dilate that contains an interior lattice point.

5. Reflexive polytopes

Definition 6.11. A lattice polytope P with $0 \in P$ is *reflexive* if P can be expressed as:

$$P = \{x \in \mathbb{R}^d : Ax \leq \mathbb{1}\}$$

where $A \in \mathbb{Z}^{n \times d}$ is an integer matrix.

Observation 6.12. To be done.

To be ended.

Scribes 2013

1. Alex Alvarez

I have done a double degree in Informatics Engineering and Mathematics at the UPC and during these years I have participated in programming contests both individually and representing the UPC. I am mainly interested in Algorithms and Data Structures and I would like to start a PhD in those topics next year, but I also enjoy learning about Discrete Mathematics, in particular Combinatorics and Graph Theory.

2. Cecilia Girón Albert

I've got my Degree in Mathematics at the Universidad Autónoma de Madrid, which I completed my fifth and sixth semesters at the University of Jyväskylä (Finland) as an Erasmus student. My degree has been mainly focused on subjects like analysis, statistics and numerical methods, although I am more interested in algebra and graph theory.

I decided to study the Master in Advanced Mathematics and Mathematical Engineering to keep developing my mathematics skills in some theoretical subjects that can be applied in real life problems. Therefore, I believe that this course is a great opportunity to learn more about algorithmic and computing science and even more importantly, it may help me to find out what field I would like to focus on in the future.

3. Anna Somoza

I have recently finished a degree in Mathematics at the Universitat Politècnica de Catalunya. During this degree I developed a great interest in Algebra fields. In particular, I took the optional subjects *Algebraic Geometry*, *Algebraic Topology* and *Galois Theory* and I wrote my Final Degree Thesis on a topic of Number Theory.

Now I'm taking the Master in Advanced Mathematics and Mathematical Engineering to develop my knowledge in these and other fields, and I my aim is to start a PhD in Number Theory next year. I enrolled this subject because I have always liked both computer science and geometry, and it seemed to be interesting. Therefore, I would be interested in the topic related to algebraic geometry.

4. Daniel Torres

My name is Daniel Torres and I am graduate in mathematics in UPC. Along the degree I have developed much interest in fields of topology, algebra and geometry, and some loathing to study (not to programming) numerical methods and modelling. I decided study this master, and particularly this subject, for expand my knowledge about my interests.

More concretely, I am doing this course with the hope it shows me about geometry.

5. Borja Elizalde

I have finished the degree in Mathematics at the UPC and I have also finished Industrial Engineering also at the UP (not CFIS).

I am interested in Number Theory and Algebraic Geometry in general, because the problems I like solving and thinking on usually belong to these fields. I am not sure at all about how I want to develop my professional career.

6. Xavier Tapia

I have finished a degree in Mathematics in UB, I spent 5 years to finish it because I have been living in Valencia for a year studying in UV. Along these years I have developed interest on Algebra and geometry, specially on finite groups and commutative algebra. Now I'm working and studying at the same time this Master but I don't have any idea what is my professional future.

Bibliography

Papers to referee

Bibliography

1. David Avis, *A revised implementation of the reverse search vertex enumeration algorithm.*, Polytopes - combinatorics and computation. DMV-seminar Oberwolfach, Germany, November 1997, Basel: Birkhäuser, 2000, pp. 177–198 (English).
2. Matthias Beck, Steven V. Sam, and Kevin M. Woods, *Maximal periods of (Ehrhart) quasi-polynomials.*, J. Comb. Theory, Ser. A **115** (2008), no. 3, 517–525 (English).
3. Matthias Beck and Thomas Zaslavsky, *Inside-out polytopes.*, Adv. Math. **205** (2006), no. 1, 134–162 (English).
4. Jürgen Bokowski and Jürgen Richter, *On the finding of final polynomials.*, Eur. J. Comb. **11** (1990), no. 1, 21–34 (English).
5. Gheorghe Craciun, Luis Garcia, and Frank Sottile, *Some geometrical aspects of control points for toric patches*, Mathematical Methods for Curves and Surfaces, Lecture Notes in Computer Science, vol. 5862, 2010, pp. 111–135.
6. Mike Develin and Josephine Yu, *Tropical polytopes and cellular resolutions.*, Exp. Math. **16** (2007), no. 3, 277–291 (English).
7. Dave Donoho, *Neighborly polytopes and sparse solutions of underdetermined linear equations*, Tech. report, Dep. Statist., Stanford Univ., Stanford, CA, 2005.
8. David Eppstein, Greg Kuperberg, and Günter M. Ziegler, *Fat 4-polytopes and fatter 3-spheres.*, Discrete geometry. In honor of W. Kuperberg’s 60th birthday, New York, NY: Marcel Dekker, 2003, pp. 239–265 (English).
9. Eric J. Friedman, *Finding a simple polytope from its graph in polynomial time.*, Discrete Comput. Geom. **41** (2009), no. 2, 249–256 (English).
10. Christian Haase and Josef Schicho, *Lattice polygons and the number $2i + 7$.*, Am. Math. Mon. **116** (2009), no. 2, 151–165 (English).
11. Birkett Huber, Jörg Rambau, and Francisco Santos, *The Cayley trick, lifting subdivisions and the Bohne-Dress theorem on zonotopal tilings.*, J. Eur. Math. Soc. (JEMS) **2** (2000), no. 2, 179–198 (English).
12. Michael Joswig, *Beneath-and-beyond revisited.*, Algebra, geometry, and software systems, Berlin: Springer, 2003, pp. 1–21 (English).
13. Felix Klein, *On the order-seven transformation of elliptic functions.*, The eightfold way. The beauty of Klein’s quartic curve, Cambridge: Cambridge University Press, 1999, pp. 287–331 (English).
14. Christophe Oguey, Michel Duneau, and André Katz, *A geometrical approach of quasiperiodic tilings.*, Commun. Math. Phys. **118** (1988), no. 1, 99–118 (English).
15. Alexander Postnikov, *Permutohedra, associahedra, and beyond.*, Int. Math. Res. Not. **2009** (2009), no. 6, 1026–1106 (English).
16. Jürgen Richter-Gebert, Bernd Sturmfels, and Thorsten Theobald, *First steps in tropical geometry.*, Idempotent mathematics and mathematical physics. Proceedings of the international workshop,

- Vienna, Austria, February 3–10, 2003, Providence, RI: American Mathematical Society (AMS), 2005, pp. 289–317 (English).
17. Raman Sanyal, *Topological obstructions for vertex numbers of Minkowski sums.*, J. Comb. Theory, Ser. A **116** (2009), no. 1, 168–179 (English).
 18. Alexander Schwartz and Günter M. Ziegler, *Construction techniques for cubical complexes, odd cubical 4-polytopes, and prescribed dual manifolds.*, Exp. Math. **13** (2004), no. 4, 385–413.
 19. Richard P. Stanley, *A zonotope associated with graphical degree sequences.*, A dual forest algorithm for the assignment problem, DIMACS, 1991, pp. 555–570 (English).
 20. Tibor Szabó and Emo Welzl, *Unique sink orientations of cubes*, Proc. 42nd Ann. IEEE Symp. on Foundations of Computer Science (FOCS), 2001, pp. 547–555.
 21. Emo Welzl, *Entering and leaving j -facets.*, Discrete Comput. Geom. **25** (2001), no. 3, 351–364.