

Further Mathematical Methods: II


In this lecture ...

- Double Integration
 - Review and examples
 - Applications to joint probability distributions
 - The gamma function
- Fourier Transforms
 - Definition and standard results

- Applications to differential equations
- Power series solutions of Ordinary Differential Equations

1 Double Integration

1.1 Evaluation of $\iint_A f(x, y) dx dy$




$$\iint_A f(x, y) \, dx \, dy$$

$$= \int_{\alpha}^{\beta} \underbrace{\left\{ \int_{x_P(y)}^{x_Q(y)} f(x, y) \, dx \right\}}_{\text{}} dy$$

So limits are given by:

5




x_p } $f_{us} \cdot f$ } x_q

6

Example: Evaluate

$$\iint_A (x+y) \, dx \, dy$$

where A is the Δ in the following diagram:



$$\alpha = 0$$

$$\beta = 1$$

$$x_P = y \quad P(y, y)$$

$$x_Q = 1 \quad Q(1, y)$$

$$I = \int_{y=0}^{y=1} \{x_P=yx + y \, dx\} dy$$

$$\int_y^1 (x+y) \, dx = \left[\frac{x^2}{2} + xy \right]_y^1 = \left(\frac{1}{2} + y \right) - \left(\frac{y^2}{2} + y^2 \right) \quad \text{inner integral}$$


$$I = \int_0^1 \left(\frac{1}{2} + y - \frac{3y^2}{2} \right) dy = \left(\frac{y}{2} + \frac{y^2}{2} - \frac{y^3}{2} \right) \Big|_0^1 \quad \text{outer integral}$$

$$= \frac{1}{2}$$

So generally

$$\iint_A f(x, y) dx dy$$


where A is defined as



x_P , x_Q function of y

$$= \int_{\alpha}^{\beta} \underbrace{\left\{ \int_{x_P}^{x_Q} f(x, y) \right\}}_{\text{repeated integral}} dy$$

We note in passing that



$$\iint_A f \, dx \, dy = \iint_{A_1} f \, dx \, dy + \iint_{A_2} f \, dx \, dy$$

$$A: A_1 + A_2$$

The main problem lies in the limits. We consider the following examples –
(Main work)

Examples:

1. A Rectangle

$$a \leq x \leq b$$


$$\alpha \leq y \leq \beta$$

Here $x_P = a$, $x_Q = b$

$$\alpha \leq y \leq \beta$$


\therefore

$$\iint_A f \, dx \, dy = \int_{\alpha}^{\beta} \left\{ \int_a^b f \, dx \right\} dy$$



2. A Triangle

with sides



$$\begin{cases} x+y = 0 \\ x-y = 0 \\ y = 2 \end{cases}$$

A is Region bounded by these 3 lines

In this case

$$\begin{aligned} x_P &= -y; \quad x_Q = y \\ \alpha &= 0; \quad \beta = 2 \end{aligned}$$


We go from $x=-y$ to $x=y$
Solved $y=0$ to 2 in the y -axis

$$\iint_A f \, dx \, dy = \int_0^2 \left\{ \int_{-y}^y f \, dx \right\} dy$$

3 A is the region defined by

$$\text{unit circle } x^2 + y^2 \leq 1, \quad \boxed{x, y \geq 0} \quad \text{first quadrant} \\ y \in [0, 1]$$

$$\iint_A f \, dx \, dy = \int_0^1 \left\{ \int_0^{\sqrt{1-y^2}} f \, dx \right\} dy$$



Difficulty: A parallelogram


For this A we do not have a simple value for x_P (or x_Q) 2 stages

For A_1 $x_P = 0, x_Q = y$

For A_2 $x_P = y - 1, x_Q = 1$

So

We don't have a unique set x_P, x_Q (simple)



$$\iint_A f \, dx \, dy = \iint_{A_1} f \, dx \, dy + \iint_{A_2} f \, dx \, dy$$


$$\iint_{A_1} = \int_0^1 \left\{ \int_0^y f \, dx \right\} dy \quad (0 \leq y \leq 1 \text{ in } A_1)$$

$$\iint_{A_2} = \int_1^2 \left\{ \int_{y-1}^1 f \, dx \right\} dy \quad (1 \leq y \leq 2 \text{ in } A_2)$$

Sometimes, then, we want to do the y -integration first:

$$\iint_A f \, dx \, dy$$

$$= \int_a^b \left\{ \int_{y_R}^{y_S} f \, dy \right\} dx$$




Here y_R , y_S depend on x

Example:

A is the parallelogram discussed earlier

$$y_R = x \quad a = 0$$

$$y_S = x + 1 \quad b = 1$$



$$\iint_A f \, dx \, dy = \int_0^1 \left\{ \int_x^{x+1} f \, dy \right\} dx$$

1.2 Uses of Double Integration

AREAS

Theorem

$$\iint_A 1 \, dx \, dy = \text{area of } A$$


↖ integrate over |

Here we have $f(x, y) = 1 \, \forall (x, y) \text{ in } A$

Example

A rectangle $a \leq x \leq b, \alpha \leq y \leq \beta$

$$\text{area} = \iint_A 1 \cdot dx dy = \int_{\alpha}^{\beta} \left\{ \int_a^b 1 dy \right\} dx$$



$$\begin{aligned}
&= \int_a^b [y]_{\alpha}^{\beta} dx = \int_a^b (\beta - \alpha) dx = (\beta - \alpha) [x]_a^b \\
&= (\beta - \alpha)(b - a)
\end{aligned}$$

1.3 Changing to Plane Polars

If

$$x = r \cos \theta$$

$$y = r \sin \theta$$

then

$$\iint_A f(x, y) \, dx \, dy = \iint_{A'} F(r, \theta) \, r \, dr \, d\theta$$

where

1. $F(r, \theta) = f(r \cos \theta, r \sin \theta)$
2. A' is the region A described in (r, θ) coordinates.

Ex: 1

1.4 Joint PDF for Continuous Random Variables

Recall that the cumulative distribution function $F(x)$ of a RV X is

$$F(x) = P(X \leq x) = \int_{-\infty}^x p(s) ds$$

$F(x)$ is related to the PDF $p(x)$ by

$$p(x) = \frac{dF}{dx}.$$

Consider the pair (X, Y) with joint pdf $p_{XY}(x, y)$ and cdf $F_{XY}(x, y)$. They are related through a similar fashion

$$p_{XY}(x, y) = \frac{\partial^2 F_{XY}(x, y)}{\partial x \partial y}.$$

Integrating this (as before) gives the cdf as

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y p_{XY}(s, t) dt ds$$

which allows to calculate the probability

$$\mathbb{P}(X \leq x, Y \leq y).$$

We can extend the simple properties of $p_{XY}(x, y)$ to two dimensions:

- $p_{XY}(x, y) \geq 0$
- $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p_{XY}(x, y) dx dy = 1$
- $\iint_R p_{X, Y}(x, y) dx dy = \mathbb{P}((X, Y) \in R).$
- $\mathbb{P}(a < X \leq b, c < Y \leq d) = \int_c^d \int_a^b f_{XY}(x, y) dx dy$

If X and Y are independent random variables the cdf can be expressed in separable form

$$F_{XY}(x, y) = F_X(x) F_Y(y).$$

Then differentiating gives

$$\frac{\partial^2 F_{XY}(x, y)}{\partial x \partial y} = \frac{\partial F_X}{\partial x} \frac{\partial F_Y}{\partial y} = p_X(x) p_Y(y).$$

$$\left. \begin{aligned} P_X(x) &= \int_{\mathbb{R}} f_{XY}(x, y) dy \\ P_Y(y) &= \int_{\mathbb{R}} p_{XY}(x, y) dx \end{aligned} \right\} \text{Marginal pdfs}$$

Ex: 2

1.5 The Gamma Function Revisited

The Gamma Function $\Gamma(x)$ is defined as

$$\Gamma(x) = \int_0^{\infty} e^{-t} t^{x-1} dt \quad (x > 0)$$

The condition on x is a convergent criterion.

If $a > 0$

$$\begin{array}{ll} \int_0^a x^p dx \text{ exists for } p > -1 & \text{to avoid } 1/0 \text{ or } \ln 0 \\ \int_a^{\infty} x^p dx \text{ exists for } p < -1 & \text{to avoid } \infty^{p+1} \end{array}$$

Integration by parts gives us $\Gamma(x)$

$$\begin{aligned} \Gamma(x+1) &= \int_0^{\infty} e^{-t} t^x dt = x \underbrace{\int_0^{\infty} e^{-t} t^{x-1} dt}_{\Gamma(x)} = x(x-1) \int_0^{\infty} e^{-t} t^{x-2} dt = \\ &\dots\dots\dots = x! \quad \text{already seen} \end{aligned} \quad (\dagger)$$

Important results:

$$\begin{aligned}\Gamma(n+1) &= n! \quad (n \geq 0) \\ \Gamma(1) &= 1\end{aligned}$$

and also from (‡)

$$\Gamma(x+1) = x\Gamma(x).$$

Theorem

$$\int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta \, d\theta = \frac{\Gamma(m)\Gamma(n)}{2\Gamma(m+n)}$$

Proof Start with the definition of the gamma function

$$\Gamma(m) = \int_0^\infty t^{m-1} e^{-t} dt$$

and make the substitution $t = x^2$ which gives $dt = 2x \, dx$

$$\begin{aligned}\Gamma(m) &= \int_0^\infty (x^2)^{m-1} \exp(-x^2) \cdot \underbrace{2x dx}_{dt} \\ &= 2 \int_0^\infty x^{2m-1} \exp(-x^2) dx\end{aligned}$$

Similarly

$$\Gamma(n) = 2 \int_0^\infty y^{2n-1} \exp(-y^2) dy$$

therefore


$$\begin{aligned}\Gamma(m)\Gamma(n) &= 4 \left(\int_0^\infty x^{2m-1} \exp(-x^2) dx \right) \left(\int_0^\infty y^{2n-1} \exp(-y^2) dy \right) \\ &= 4 \iint_A x^{2m-1} y^{2n-1} e^{-(x^2+y^2)} dx dy\end{aligned}$$

where A is the region of integration defined by the first (positive) quadrant.

Introduce polar coordinates

$$x = r \cos \theta$$

$$y = r \sin \theta$$



$$\lim_{R \rightarrow \infty}$$

$$4 \iint_A r^{2m-1} \cos^{2m-1} \theta \, r^{2n-1} \sin^{2n-1} \theta \cdot e^{-r^2} \underbrace{(r dr d\theta)}_{dx dy}$$

to transform the integrand to

$$r^{2m+2n-2} \cos^{2m-1} \theta \sin^{2n-1} \theta \exp(-r^2)$$

and $\boxed{dx dy \rightarrow r dr d\theta}$ group r & θ terms separately

$$\Gamma(m) \Gamma(n) = 4 \underbrace{\int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta \int_0^\infty r^{2(m+n)-1} e^{-(r^2)} dr}_{\text{integral we want}} \underbrace{\int_0^\infty (r^2)^{m+n-1} e^{-r^2} r dr}_{\frac{1}{2} \Gamma(m+n)}$$

so rearranging gives the result

$$\boxed{\int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta = \frac{\Gamma(m) \Gamma(n)}{2 \Gamma(m+n)}}$$

Example Calculate $\int_0^{\pi/2} \cos^4 \theta \sin^3 \theta d\theta = \int_0^{\pi/2} \cos^{2m-1} \theta \sin^{2n-1} \theta d\theta$

Hence

$$2m-1 = 4 \rightarrow m = 5/2$$

$$2n-1 = 3 \rightarrow n = 2$$

$$= \frac{1}{2} \Gamma(m+n)$$

$$\begin{aligned} & \int_0^\infty (r^2)^{m+n-1} e^{-r^2} r dr \\ & \text{Put } t=r^2 \\ & dt=2r dr \\ & = \int_0^\infty (t)^{m+n-1} e^{-t} \frac{1}{2} dt \\ & = \frac{1}{2} \int_0^\infty t^{m+n-1} e^{-t} dt \end{aligned}$$

so integral equals

$$\frac{\Gamma\left(\frac{5}{2}\right) \Gamma(2)}{2\Gamma\left(\frac{9}{2}\right)} \overset{\text{Verify}}{=} \frac{\frac{3\sqrt{\pi}}{2} \cdot 1}{2\left(\frac{7}{2} \cdot \frac{5}{2} \cdot \frac{3}{2} \cdot \frac{\sqrt{\pi}}{2}\right)} = \frac{2}{35}$$

Example $I = \int_0^{\pi/2} \cos^6 \theta \, d\theta$ No \sin^{even} term

$$2m - 1 = 6 \longrightarrow m = 7/2$$

$$2n - 1 = 0 \longrightarrow n = 1/2$$

Hence $I =$

$$\frac{\Gamma\left(\frac{7}{2}\right) \Gamma\left(\frac{1}{2}\right)}{2\Gamma(4)} = \frac{\frac{5}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} \cdot \frac{\sqrt{\pi}}{2} \cdot \sqrt{\pi}}{2(3 \cdot 2)} = \frac{5\pi}{32}$$

2 The Fourier Transform

If $f = f(x)$ then consider

$$\hat{f}(\omega) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{ix\omega} dx. \quad \text{or} \quad \frac{1}{2\pi}$$

If this special integral converges, it is called the *Fourier Transform* of $f(x)$.
Similar to the case of Laplace Transforms, it is denoted as $\mathcal{F}(f)$, i.e.

$$\int_0^{\infty} f(t) e^{-st} dt \quad \mathcal{F}(f) = \int_{-\infty}^{\infty} f(x) e^{ix\omega} dx = \hat{f}(\omega).$$

The *Inverse Fourier Transform* is then i.e. \mathbb{C} -plane to \mathbb{R} -plane

$$\mathcal{F}^{-1}(\hat{f}(\omega)) = \int_{-\infty}^{\infty} \hat{f}(\omega) e^{-ix\omega} d\omega = f(x).$$

The convergent property means that $\hat{f}(\omega)$ is bounded and we have

$$\int_{-\infty}^{\infty} |f(x)| dx < \infty.$$

Functions of this type $f(x) \in L_1(-\infty, \infty)$ and are called square integrable.

We know from integration that

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx. \quad \text{Standard result}$$

Hence

$$\begin{aligned} |\hat{f}(\omega)| &= \left| \int_{\mathbb{R}} f(x) e^{ix\omega} dx \right| \\ &\leq \int_{\mathbb{R}} |f(x) e^{ix\omega}| dx \quad \text{from above result.} \end{aligned}$$

and Euler's identity $e^{i\theta} = \cos \theta + i \sin \theta$ implies that $|e^{i\theta}| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1$, therefore

$$|\hat{f}(\omega)| \leq \int_{\mathbb{R}} |f(x)| dx < \infty.$$

In addition to the boundedness of $\hat{f}(\omega)$, it is also continuous (requires a $\delta - \epsilon$ proof).


Ex: Fourier transform of $e^x = f(x)$

$$\begin{aligned}\hat{f}(\omega) &= f(\omega) = \int_{\mathbb{R}} f(x) e^{i\omega x} dx = \int_{-\infty}^{\infty} e^x e^{i\omega x} dx \\ &= \int_{-\infty}^{\infty} e^{(1+i\omega)x} dx = \left. \frac{1}{(1+i\omega)} e^{(1+i\omega)x} \right|_{-\infty}^{\infty} \rightarrow \infty \text{ i.e. integral diverges.}\end{aligned}$$

\therefore Fourier transform of e^x does not exist.

Example: Obtain the Fourier transform of $f(x) = e^{-|x|}$

$$\begin{aligned}
 \hat{f}(\omega) &= \mathcal{F}(f) = \int_{-\infty}^{\infty} f(x) e^{ix\omega} dx \\
 &= \int_{-\infty}^{\infty} e^{-|x|} e^{ix\omega} dx \\
 &= \int_{-\infty}^0 e^{-|x|} e^{ix\omega} dx + \int_0^{\infty} e^{-|x|} e^{ix\omega} dx \\
 &= \int_{-\infty}^0 e^x e^{ix\omega} dx + \int_0^{\infty} e^{-x} e^{ix\omega} dx = \\
 &\quad \int_{-\infty}^0 \exp[(1+i\omega)x] dx + \int_0^{\infty} \exp[-(1-i\omega)x] dx \\
 &= \frac{1}{(1+i\omega)} \exp[(1+i\omega)x] \Big|_{-\infty}^0 \oplus \frac{1}{(1-i\omega)} \exp[-(1-i\omega)x] \Big|_0^{\infty} \\
 &= \frac{1}{(1+i\omega)} + \frac{1}{(1-i\omega)} = \frac{2}{(1+\omega^2)}
 \end{aligned}$$



Our interest in differential equations continues, hence the reason for introducing this transform. We now look at obtaining Fourier transforms of derivative terms. We assume that $f(x)$ is continuous and $f(x) \rightarrow 0$ as $x \rightarrow \pm\infty$. Consider

$$\mathcal{F}\{f'(x)\} = \int_{\mathbb{R}} f'(x) e^{ix\omega} dx$$

$$\begin{aligned} v &= e^{ix\omega} & v' &= f' \\ v' &= i\omega e^{ix\omega} & v &= f \end{aligned}$$

which is simplified using integration by parts

$$\underbrace{f(x) e^{ix\omega}}_u \underbrace{\Big|_{-\infty}^{\infty}}_v - i\omega \underbrace{\int_{\mathbb{R}} f(x) e^{ix\omega} dx}_{\int u v'}$$

so

$$\mathcal{F}\{f'(x)\} = -i\omega \underbrace{\int_{\mathbb{R}} f(x) e^{ix\omega} dx}_{\hat{f}(\omega)} = -i\omega \hat{f}(\omega).$$

We can obtain the Fourier transform for the second derivative by performing integration by parts (twice) to give

$$\mathcal{F}\{f''(x)\} = (-i\omega)^2 \mathcal{F}\{f(x)\} = -\omega^2 \hat{f}(\omega).$$

Check

$$\begin{aligned} v &= e^{ix\omega} & v' &= f' \\ v' &= i\omega e^{ix\omega} & v &= f \end{aligned}$$

Example: Solve the diffusion equation problem

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad -\infty < x < \infty, \quad t > 0$$

$$u(x, 0) = e^{-|x|}, \quad -\infty < x < \infty$$

Here $u = u(x, t)$, so we begin by defining

$$\mathcal{F}\{u(x, t)\} = \int_{-\infty}^{\infty} u(x, t) e^{ix\omega} dx = \hat{u}(\omega, t). \quad \text{So } \mathcal{F}\{u(x, t)\} = \hat{u}(\omega, t)$$

Now take Fourier transforms of our PDE, i.e.

$$\mathcal{F}\left\{\frac{\partial u}{\partial t}\right\} = \mathcal{F}\left\{\frac{\partial^2 u}{\partial x^2}\right\}$$

to obtain

$$\frac{d\hat{u}}{dt} = -\omega^2 \hat{u}(\omega, t). \quad \mathcal{F}\left(\frac{\partial u}{\partial t}\right) = \frac{\partial}{\partial t} \underbrace{\mathcal{F}(u)}_{\hat{u}}$$

We note that the second order PDE has been reduced to a first order equation of type variable separable. This has solution

$$\hat{u}(\omega, t) = C e^{-\omega^2 t}.$$

$$\begin{aligned} \int \frac{d\hat{u}}{\hat{u}} &= -\omega^2 \int dt \\ \ln \hat{u} &= -\omega^2 t + k \\ \hat{u} &= C e^{-\omega^2 t} \end{aligned}$$

We can find the constant of integration transforming the initial condition

$$\mathcal{F}\{u(x, 0)\} = \mathcal{F}\{e^{-|x|}\}$$

$$\hat{u}(\omega, 0) = \int_{-\infty}^{\infty} e^{-|x|} e^{i\omega x} dx = \frac{2}{(1+\omega^2)} \quad \text{from earlier example}$$

Applying this to the solution $\hat{u}(\omega, t)$ gives

$$\hat{u}(\omega, 0) = C = \frac{2}{(1+\omega^2)},$$

hence

$$\hat{u}(\omega, t) = \frac{2}{(1+\omega^2)} e^{-\omega^2 t}.$$

this is in ω - t plane

How do we get $u(x, t)$?

$$\hat{u} = C e^{-\omega^2 t}$$

at $t=0$

$$\frac{2}{1+\omega^2} = C e^0$$

We now use the inverse transform to get $u(x, t) = \mathcal{F}^{-1}(\hat{u}(\omega, t))$

$$\begin{aligned}
 &= \int_{-\infty}^{\infty} \hat{u}(\omega, t) e^{-i\omega x} d\omega \\
 &= 2 \int_{-\infty}^{\infty} \frac{1}{(1 + \omega^2)} e^{-\omega^2 t} e^{-i\omega x} d\omega \\
 &= 2 \int_{-\infty}^{\infty} \frac{1}{(1 + \omega^2)} e^{-\omega^2 t} (\cos \omega x - i \sin \omega x) d\omega \\
 &= 2 \int_{-\infty}^{\infty} \frac{1}{(1 + \omega^2)} e^{-\omega^2 t} \cos \omega x d\omega - 2i \int_{-\infty}^{\infty} \frac{1}{(1 + \omega^2)} e^{-\omega^2 t} \underbrace{\sin \omega x}_{\text{odd}} d\omega.
 \end{aligned}$$

This now simplifies nicely because $\frac{1}{(1 + \omega^2)} e^{-\omega^2 t} \sin \omega x$ is an odd function,

hence

$$\int_{-\infty}^{\infty} \frac{1}{(1 + \omega^2)} e^{-\omega^2 t} \sin \omega x d\omega = 0.$$

Therefore

$$u(x, t) = 2 \int_{-\infty}^{\infty} \frac{1}{(1 + \omega^2)} e^{-\omega^2 t} \cos \omega x \, d\omega.$$

Now needs treatment of Residue Theory.

3 Power Series Solutions

3.1 Introduction

The Euler equation has a nice structure, i.e.

$$ax^2 \frac{d^2 y}{dx^2} + bx \frac{dy}{dx} + cy = 0$$

where the order of each derivative term and power of its coefficient in x is the same. The next step is to move away from this "nice pattern" and consider a more general equation of the form

$$\frac{d^2 y}{dx^2} + p(x) \frac{dy}{dx} + q(x) y = 0 \quad (1)$$

We look for solutions in the neighbourhood of $x = 0$.

We say that $x = 0$ is an ordinary point of the differential equation (1) if both $p(x)$ and $q(x)$ have Taylor expansions about $x = 0$.

i.e.

$$\begin{aligned} p(x) &= p_0 + p_1x + p_2x^2 + O(x^3) \\ q(x) &= q_0 + q_1x + q_2x^2 + O(x^3) \end{aligned}$$

i.e. p & q are $\mathcal{O}(x^3)$ diff'ble at $x=0$

with both $p_i, q_i \sim O(1)$ where $i = 0, 1, \dots, n$.

If either or both $p(x), q(x)$ do not have Taylor expansions about $x = 0$, then $x = 0$ is a singular point for the D.E.

Regular Singular Point: $x^2p(x)$ and $x^2q(x)$ have Taylor expansions about $x = 0$.

Irregular Singular Point: all other points.

Examples:

$$1. \quad x \frac{d^2 y}{dx^2} + x^3 \frac{dy}{dx} + xy = 0$$

This can be written in standard form as $\frac{d^2 y}{dx^2} + x^2 \frac{dy}{dx} + y = 0 \Rightarrow p(x) = x^2$ & $q(x) = 1$ which both have Taylor expansions about $x = 0$.

Therefore $x = 0$ is an ordinary point of the differential equation.

2. $x^3 \frac{d^2 y}{dx^2} + 2x^2 \frac{dy}{dx} + 5x^2 y = 0$

which becomes $\frac{d^2 y}{dx^2} + \frac{2dy}{x dx} + \frac{5y}{x^2} = 0$ and $p(x) = \frac{2}{x}$ & $q(x) = \frac{5}{x^2}$ do not have a Taylor expansion about $x = 0$ - however $xp(x) = 2$ & $x^2 q(x) = 5x$ do.

Therefore $x = 0$ is a regular singular point of the differential equation.

3. $\frac{d^2 y}{dx^2} - \frac{1}{x^2} \frac{dy}{dx} + \frac{4}{x^3} y = 0$

$p(x) = O\left(\frac{1}{x^2}\right)$ & $xp(x) = O\left(\frac{1}{x}\right); \quad q(x) = O\left(\frac{1}{x^3}\right)$ & $x^2 q(x) = O\left(\frac{1}{x}\right)$

None of these expressions have a Taylor expansion about $x = 0$.

Therefore $x = 0$ is an irregular singular point of the given differential equation.

Ex: Consider $(1-x^2)y'' - 2xy' + 6y = 0$ Legendre's eqⁿ of order 2

$x=0$ is an ordinary point. $y'' - \frac{2x}{1-x^2}y' + \frac{6}{1-x^2}y = 0$

$p = \frac{2x}{(1-x)(1+x)}$ regular everywhere except $x = x_0 = \pm 1$

$q = \frac{6}{(1-x)(1+x)}$ Look at $(x-x_0)p(x) = \frac{2x(x-1)}{-(x-1)(1+x)} = -\frac{2x}{1+x}$ both have TSF at $x=1$ $\therefore x=1$ is a regular singular point

$x_0 = -1$

$(x+1)p(x) = \frac{2x}{1-x}$ both diff. at $x=1$
 $(x+1)^2q(x) = \frac{6(x+1)}{(1-x)}$ \therefore Reg. sing. pt.

3.2 Ordinary Point

Assume a solution of (1) of the form

$$y(x) = \sum_{n=0}^{\infty} A_n x^n \quad (A_0 \neq 0) \quad (2)$$

$$A_{-1} = A_{-2} = \dots = 0$$

with A_n constant.

Since no boundary conditions are imposed, the general solution involves two arbitrary constants - else the constants can be determined.

Substitute (2) into the equation given by (1) and equate to zero the coefficients of various powers of x .

$$y(x) = \sum_{n=0}^{\infty} A_n x^n \rightarrow q(x)y \sim (q_0 + q_1x + q_2x^2)(A_0 + A_1x + A_2x^2)$$

$$y'(x) = \sum_{n=0}^{\infty} n A_n x^{n-1} \rightarrow p(x)y' \sim (p_0 + p_1x + p_2x^2)(A_1 + 2A_2x + 3A_3x^2)$$

$$y''(x) = \sum_{n=0}^{\infty} n(n-1) A_n x^{n-2} \rightarrow y'' \sim 2A_2 + 6A_3x + 12A_4x^2$$

$$\underbrace{y''}_{p} + \underbrace{6A_3x}_{y'} + (p_0 + p_1x)(A_1 + 2A_2x) + (q_0 + q_1x)(A_0 + A_1x) = 0$$

$$O(1): A_0q_0 + A_1p_0 + 2A_2 = 0$$

$$O(x): q_0A_1 + 2p_0A_2 + p_1A_1 + q_1A_0 + 6A_3 = 0$$

All coefficients can be expressed in terms of A_0 and A_1 which can be arbitrary.

Go to first
order want
 A_0, A_1
 \therefore 2nd order
eqn

Example

Obtain the general solution of

$$y'' - 2xy' + y = 0$$

$$y' = \sum_n A_n x^{n-1}$$
$$y'' = \sum_n n(n-1) A_n x^{n-2}$$

about the ordinary point $x = 0$.

We assume a solution of the form $y(x) = \sum_{n=0}^{\infty} A_n x^n$ and substitute the expression and its derivatives into the ODE to yield

$$\sum_{n=0}^{\infty} n(n-1) A_n x^{n-2} - 2x \sum_{n=0}^{\infty} n A_n x^{n-1} + \sum_{n=0}^{\infty} A_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) A_n x^{n-2} + \underbrace{\sum_{n=0}^{\infty} (1-2n) A_n x^n}_{-2 \times y' + y} = 0$$

Want A_{n+2} in terms of A_n i.e. $\sum (\quad) x^{n-2} = 0$

(45)

We require a recurrence relation for which a "trick" is used in the summation.

In the second summation above, the n term is changed to $(n-2)$ to give

$$\sum_{n-2=0}^{\infty} (1-2(n-2)) A_{n-2} x^{n-2} \text{ which is equivalent to having } \sum_{n=2}^{\infty} \dots$$

$$\sum_{n=0}^{\infty} n(n-1) A_n x^{n-2} + \underbrace{\sum_{n=2}^{\infty} (1-2(n-2)) A_{n-2} x^{n-2}}_{\substack{\text{trick} \\ n=2 \equiv n=0 \\ n=3 \equiv n=1 \text{ etc.}}} = 0$$

We are still unable to write the lhs of the expression above as one term of $O(x^{n-2})$, because the lower limit of the first summation starts at $n=0$, whilst the other begins at $n=2$. This minor problem can be easily overcome by writing

$$\sum_{n=0}^{\infty} n(n-1) A_n x^{n-2} + \sum_{n=0}^{\infty} (5-2n) A_{n-2} x^{n-2} = 0 \quad (\dagger)$$

with $A_{-2} = A_{-1} = 0$

because $A_{-2} = A_{-1} = 0$ and $A_0 \neq 0$, and (*) can now be expressed as

$$\sum_{n=0}^{\infty} \{n(n-1)A_n + (5-2n)A_{n-2}\} x^{n-2} = 0.$$

Collecting coefficients of x^{n-2} : ↗ equals zero rearrange

$$A_n = \frac{(2n-5)}{n(n-1)} A_{n-2} \quad (n \geq 2)$$

or

$$A_{n+2} = \frac{(2n-1)}{(n+2)(n+1)} A_n$$

which gives us the recurrence relationship which we sought.

$$n=0: \quad A_2 = -\frac{1}{2}A_0; \quad n=1: \quad A_3 = \frac{1}{6}A_1 = \frac{1}{3!}A_1$$

So we see that all terms A_{2k} will be in terms of A_0 and odd ones in terms of A_1 .

↖ correction

A_{2k+1}

(7)

$$n = 2: \quad A_4 = \frac{3}{4.3} A_2 = -\frac{3}{4.32} A_0 = -\frac{3}{4!} A_0$$

$$n = 3: \quad A_5 = \frac{5}{5.4} A_3 = \frac{5}{5.43!} A_1 = \frac{5}{5!} A_1$$

$$n = 4: \quad A_6 = \frac{7}{6.5} A_4 = -\frac{7}{6.54!} A_0 = -\frac{21}{6!} A_0$$

$$n = 5: \quad A_7 = \frac{9}{7.6} A_5 = \frac{9}{7.65!} A_1 = \frac{45}{7!} A_1$$

The solution is

$$y(x) = \sum_{n=0}^{\infty} A_n x^n = \sum_{k=0}^{\infty} \left(A_{2k} x^{2k} + A_{2k+1} x^{2k+1} \right)$$

$$= A_0 \underbrace{\left[1 - \frac{1}{2!} x^2 - \frac{3}{4!} x^4 - \frac{21}{6!} x^6 + O(x^8) \right]}_{=y_1} + \underbrace{\left[\frac{5}{5!} x^5 + \frac{451}{7!} x^7 + O(x^9) \right]}_{=y_2} + \text{even powers in terms of } A_0$$

$$= A_0 y_1(x) + A_1 y_2(x) + \text{odd in terms of } A_1$$

The linear combination $A_0 y_1(x) + A_1 y_2$ becomes the general solution of the equation. The terms A_0 , A_1 are arbitrary.

$$I \cdot c / \beta \cdot c \rightarrow A_0, A_1$$

(1a)

Ex: 1


$$\int_{-\infty}^{\infty} e^{-x^2} dx \quad \text{put } I = \int_{-\infty}^{\infty} e^{-x^2} dx$$

$$\text{then also } I = \int_{-\infty}^{\infty} e^{-y^2} dy \rightarrow$$

$$I^2 = \left(\int_{\mathbb{R}} e^{-x^2} dx \right) \left(\int_{\mathbb{R}} e^{-y^2} dy \right)$$

$$= \int_{\mathbb{R}} \int_{\mathbb{R}} e^{-(x^2+y^2)} dx dy$$

Use plane polars $x = r \cos \theta$
 $y = r \sin \theta$



$$dx dy = r dr d\theta$$

$$e^{-(x^2+y^2)} = e^{-r^2}$$

$$0 < r < \infty$$

$$0 < \theta < 2\pi$$

$$\int_0^{2\pi} \int_0^{\infty} e^{-r^2} r dr d\theta$$

①

②

$$\textcircled{1} \int_0^{\infty} e^{-r^2} r \, dr \quad \text{use substit. } u=r^2$$

$$= \frac{1}{2}$$

$$\frac{1}{2} \int_0^{2\pi} d\theta = \frac{\theta}{2} \Big|_0^{2\pi} = \pi$$

$$\therefore I^2 = \pi \rightarrow I = \sqrt{\pi}$$

$$\therefore \int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$$


$$\Rightarrow \int_0^{\infty} e^{-x^2} dx = \frac{\sqrt{\pi}}{2}$$

(1c)

$$\int_{-a}^a \int_0^{\sqrt{a^2-x^2}} (x^2+y^2)^{3/2} dy dx$$

$$y = \sqrt{a^2-x^2} \Rightarrow x^2+y^2 = a^2 = r^2$$

$$-a \leq x \leq a$$



$$0 \leq r \leq a$$

$$dy dx = r dr d\theta$$

$$0 \leq \theta \leq \pi$$

$$\int_0^\pi \int_0^a (r^2)^{3/2} r dr d\theta$$

$$\int_0^\pi \int_0^a r^4 dr d\theta = \int_0^\pi \left. \frac{r^5}{5} \right|_0^a d\theta$$

$$= \frac{1}{5} \int_0^\pi a^5 d\theta = \left. \frac{a^5}{5} \theta \right|_0^\pi = \frac{\pi a^5}{5}$$

(2a)

Ex: 2 Joint pdf of (X, Y) is given by

$$f_{XY} = \begin{cases} k e^{-(x+2y)} & x > 0, y > 0 \\ 0 & \text{otherwise} \end{cases}$$

a) find k . $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(x, y) dx dy = 1$

$$k \int_0^{\infty} \int_0^{\infty} e^{-(x+2y)} dx dy = k \int_0^{\infty} -e^{-(x+2y)} \Big|_0^{\infty} dy$$


$$k \int_0^{\infty} + e^{-2y} dy = -\frac{k}{2} e^{-2y} \Big|_0^{\infty} = 1 \rightarrow \boxed{k=2}$$

b) $P[1 < X < \infty, 0 < Y < 1] = 2 \int_0^1 \int_1^{\infty} e^{-(x+2y)} dx dy$

$$= 2 \int_0^1 e^{-(1+2y)} dy = 2e^{-1} \int_0^1 e^{-2y} dy$$

$$= -e^{-(1+2y)} \Big|_0^1 = -[e^{-3} - e^{-1}] = e^{-1} - e^{-3}$$

$$P[X < Y]$$



$$P[X < Y] = 2 \int_0^{\infty} \int_{x=0}^{x=y} e^{-(x+2y)} dx dy$$

$$= -2 \int_0^{\infty} (e^{-3y} - e^{-2y}) dy = \frac{1}{3}$$

To check indep. Must show

$$p_{xy}(x,y) = p_x(x) p_y(y)$$

$$p_x(x) = \int_0^{\infty} p_{xy}(x,y) dy = k e^{-x} \int_0^{\infty} e^{-2y} dy = e^{-x}$$

$$p_y(y) = \int_0^{\infty} p_{xy}(x,y) dx = k e^{-2y} \int_0^{\infty} e^{-x} dx = 2 e^{-2y}$$

$$\text{i.e. } p_x(x) p_y(y) = 2 e^{-x} e^{-2y} = 2 e^{-(x+2y)} = p_{xy}(x,y)$$

\therefore indep.