Explanations

Joanne Dumont

November 2021

1 AllDifferent($\{X_1, \ldots, X_n\}$)

$$X_{i} = t \iff b_{it} \qquad \forall i \in [1, n] \forall t \in [1, m]$$
$$\sum_{i \in [1, n]} b_{it} \leq 1 \qquad \forall t \in [1, m]$$

$$\frac{X_{i'} = t, \ \forall i', \ i' \neq i, \ i' \in [[1, n]], \ i \in [[1, n]]}{X_i \neq t}$$

2 AllEqual($\{X_1,\ldots,X_n\}$)

$$X_{i} \geq t \iff b_{it} \qquad \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket$$

$$\bigwedge_{i \in \llbracket 1, n \rrbracket} b_{it} \iff b1_{t} \qquad \forall t \in \llbracket 1, m \rrbracket$$

$$\bigwedge_{i \in \llbracket 1, n \rrbracket} \neg b_{it} \iff b2_{t} \qquad \forall t \in \llbracket 1, m \rrbracket$$

$$(b1_{t} \lor b2_{t}) \qquad \forall t \in \llbracket 1, m \rrbracket$$

$$\frac{X_{i} \geq t, \ \exists i, \ i \in \llbracket 1, n \rrbracket}{X_{i} \geq t}$$

$$\frac{X_{i'} < t, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket}{X_{i} \geq t}$$

$$\frac{X_{i'} \geq t, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket}{X_{i} < t}$$

$$\frac{X_{i} < t, \ \exists i, \ i \in \llbracket 1, n \rrbracket}{X_{i} < t}$$

3 NValue $(N, \{X_1, \ldots, X_n\})$

$$X_{i} = t \iff b_{it} \qquad \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket$$

$$\bigvee_{i \in \llbracket 1, n \rrbracket} b_{it} \iff b2_{t} \qquad \forall t \in \llbracket 1, m \rrbracket$$

$$\sum_{t \in \llbracket 1, m \rrbracket} b2_{t} = p \iff b3_{p} \qquad \forall t \in \llbracket 1, m \rrbracket \forall p \in \llbracket 1, n \rrbracket$$

$$N = p \iff b3_{p} \qquad \forall p \in \llbracket 1, n \rrbracket$$

$$\underbrace{X_{i} \neq t, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, m \rrbracket \quad N = p, \ \forall p, \ p \in \llbracket 1, n \rrbracket}_{X_{i} = t}$$

$$\underbrace{X_{i'} \neq t, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket}_{X_{i} = t}$$

$$\underbrace{X_{i} = t, \ \exists i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, m \rrbracket \quad N = p, \ \forall p, \ p \in \llbracket 1, n \rrbracket}_{X_{i} \neq t}$$

$$\underbrace{X_{i} = t, \ \exists i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, m \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket}_{N = p}$$

$$\underbrace{X_{i} \neq t, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, n \rrbracket}_{N = p} \qquad X_{i} = t, \ \exists i, \ i \in \llbracket 1, n \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket}_{N = p}$$

4 AtLeastNValue($N, \{X_1, \dots, X_n\}$)

$$X_{i} = t \iff b_{it} \qquad \forall i \in [1, n] \forall t \in [1, m]$$

$$\bigvee_{i \in [1, n]} b_{it} \iff b2_{t} \qquad \forall t \in [1, m]$$

$$\sum_{t \in [1, m]} b2_{t} \ge p \iff b3_{p} \qquad \forall t \in [1, m]$$

$$N \ge p \iff b3_{p} \qquad \forall t \in [1, m]$$

 $\frac{X_i=t, \ \exists i, \ i \in \llbracket 1,n \rrbracket, \ \forall t, \ t \in \llbracket 1,m \rrbracket, \ \forall i, \ i \in \llbracket 1,n \rrbracket}{N \neq p}$

 $\frac{X_i \neq t, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, m \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket}{N \neq p}$

 $\frac{X_i \neq t', \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t', \ t' \neq t, \ t' \in \llbracket 1, m \rrbracket, \ t \in \llbracket 1, m \rrbracket \quad N \geq p, \ \forall p, \ p \in \llbracket 1, n \rrbracket}{X_i = t}$

$$\frac{X_{i'} \neq t, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket}{X_i = t}$$

 $\underbrace{X_i = t', \ \exists i, \ i \in [\![1,n]\!], \ \forall i, \ i \in [\![1,n]\!], \ \forall t', \ t' \neq t, \ t' \in [\![1,m]\!], \ t \in [\![1,m]\!] \quad N < p, \ \forall p, \ p \in [\![1,n]\!]}_{X_i \neq t}$

$$\frac{X_i=t, \ \exists i, \ i \in \llbracket 1,n \rrbracket, \ \forall i, \ i \in \llbracket 1,n \rrbracket, \ \forall t, \ t \in \llbracket 1,m \rrbracket, \ \forall i, \ i \in \llbracket 1,n \rrbracket}{N \geq p}$$

$$\frac{X_i \neq t, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, m \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket}{N < p}$$

5 AtMostNValue($N, \{X_1, \dots, X_n\}$)

$$X_{i} = t \iff b_{it} \qquad \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket$$

$$\bigvee_{i \in \llbracket 1, n \rrbracket} b_{it} \iff b2_{t} \qquad \forall t \in \llbracket 1, m \rrbracket$$

$$\sum_{t \in \llbracket 1, m \rrbracket} b2_{t}
$$N \geq p \iff b3_{p} \qquad \forall t \in \llbracket 1, m \rrbracket$$

$$\underbrace{X_{i} \neq t', \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t', \ t' \neq t, \ t' \in \llbracket 1, m \rrbracket, \ t \in \llbracket 1, m \rrbracket \qquad N \geq p, \ \forall p, \ p \in \llbracket 1, n \rrbracket}_{X_{i} = t}$$

$$\underbrace{X_{i'} \neq t, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket}_{X_{i} = t}$$

$$\underbrace{X_{i} = t', \ \exists i, \ i \in \llbracket 1, n \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t', \ t' \neq t, \ t' \in \llbracket 1, m \rrbracket, \ t \in \llbracket 1, m \rrbracket \qquad N < p, \ \forall p, \ p \in \llbracket 1, n \rrbracket}_{X_{i} \neq t}$$$$

 $X_{i} = t, \exists i, i \in \llbracket 1, n \rrbracket, \forall i, i \in \llbracket 1, n \rrbracket, \forall t, t \in \llbracket 1, m \rrbracket, \forall i, i \in \llbracket 1, n \rrbracket}$ $N \geq p$ $X_{i} \neq t, \forall i, i \in \llbracket 1, n \rrbracket, \forall i, i \in \llbracket 1, n \rrbracket, \forall t, t \in \llbracket 1, m \rrbracket, \forall i, i \in \llbracket 1, n \rrbracket}$

 $\frac{X_i \neq t, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket, \ \forall t, \ t \in \llbracket 1, m \rrbracket, \ \forall i, \ i \in \llbracket 1, n \rrbracket}{N < p}$

6 Cumulative $(\{X_1, \ldots, X_n\}, \{d_1, \ldots, d_n\}, c)$

$$X_{i} \geq t \iff b_{it} \qquad \forall i \in [1, n] \forall t \in [1, m]$$

$$(b_{i(t-d_{i})} \wedge \neg b_{it}) \iff b2_{it} \qquad \forall i \in [1, n] \forall t \in [1, m]$$

$$\sum_{i \in [1, n]} b2_{it} \leq c \qquad \forall t \in [1, m]$$

 $\frac{X_i \geq t', \ t' = t - d_i \quad \ X_{i'} < t, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket, \ \ X_{i'} \geq t', \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket, \ t' = t - d_{i'} \leq t', \ \forall i', \ i' \neq i, \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i' = t - d_{i'} \leq t', \ \forall i' \in \llbracket 1, n \rrbracket, \ i' \in \llbracket 1, n \rrbracket, \ i' = t - d_{i'} \leq t', \ \forall i' \in \llbracket 1, n \rrbracket, \ i' \in \llbracket$

 $\frac{X_{i} < t', \ t' = t + d_{i} \quad X_{i'} < t', \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket, \ t' = t + d_{i} \quad X_{i'} \geq t'', \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ t'' = t' - d_{i'}, \ t' = t + d_{i'}, \ X_{i} < t' = t + d_{i'}, \ X_{i'} \geq t'', \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket, \ t'' = t' - d_{i'}, \ t' = t + d_{i'}, \ X_{i'} < t' = t + d_{i'}, \ X_{i'} \leq t'', \ X_{i'} \leq t'$

7 Element $(I, \{X_1, \ldots, X_n\}, V)$

$$\begin{split} X_i &= t \iff b^X_{it} & \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket \\ I &= i \iff b^I_i & \forall i \in \llbracket 1, n \rrbracket \\ V &= t \iff b^V_t & \forall t \in \llbracket 1, m \rrbracket \\ \neg b^V_t \wedge \neg b^I_i \wedge b^X_{it} & \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket \\ b^V_t \wedge \neg b^I_i \wedge \neg b^X_{it} & \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket \end{split}$$

$$\begin{split} \frac{I=i \quad V=t}{X_i=t} \\ \frac{I=i \quad V\neq t}{X_i\neq t} \\ \frac{X_i\neq t, \ \forall t, \ t\in \llbracket 1,m\rrbracket \quad V=t, \ \forall t, \ t\in \llbracket 1,m\rrbracket}{I\neq i} \\ \frac{X_i=t, \ \forall t, \ t\in \llbracket 1,m\rrbracket \quad V\neq t, \ \forall t, \ t\in \llbracket 1,m\rrbracket}{I\neq i} \\ \frac{X_i=t, \ \forall i, \ i\in \llbracket 1,n\rrbracket \quad I=i, \ \forall i, \ i\in \llbracket 1,n\rrbracket}{V=t} \\ \frac{X_i\neq t, \ \forall i, \ i\in \llbracket 1,n\rrbracket \quad I=i, \ \forall i, \ i\in \llbracket 1,n\rrbracket}{V\neq t} \end{split}$$

$\mathbf{Gcc}(\{X_1,\ldots,X_n\},V,O)$

$$\begin{aligned} X_i &= t \iff b_{it} & \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket \\ O_t &= p \iff b 2_{tp} & \forall t \in \llbracket 1, m \rrbracket \forall p \in \llbracket 1, n \rrbracket \\ \sum_{i \in \llbracket 1, n \rrbracket} b_{it} \geq p \iff b 2_{tp} & \forall t \in \llbracket 1, m \rrbracket \forall p \in \llbracket 1, n \rrbracket \end{aligned}$$

$$\begin{split} \frac{X_{i'} \neq t, \ \forall i', \ i' \neq i, \ i' \in [\![1,n]\!], \ i \in [\![1,n]\!] \quad O_t \geq p, \ \forall p, \ p \in [\![1,n]\!]}{X_i = t} \\ \frac{X_{i'} = t, \ \forall i', \ i' \neq i, \ i' \in [\![1,n]\!], \ i \in [\![1,n]\!] \quad O_t < p, \ \forall p, \ p \in [\![1,n]\!]}{X_i \neq t} \\ \frac{X_i = t, \ \forall i, \ i \in [\![1,n]\!], \ \forall i, \ i \in [\![1,n]\!]}{O_t \geq p} \\ \frac{X_i \neq t, \ \forall i, \ i \in [\![1,n]\!], \ \forall i, \ i \in [\![1,n]\!]}{O_t < p} \end{split}$$

9 Increasing($\{X_1, \ldots, X_n\}$)

$$X_{i} \geq t \iff b_{it} \qquad \forall i \in [1, n] \forall t \in [1, m]$$

$$(\neg b_{(i-1)t} \lor b_{it}) \qquad \forall i \in [2, n] \forall t \in [1, m]$$

$$\frac{X_{i'} \geq t, \ i' = i - 1}{X_{i} \geq t}$$

$$\frac{X_{i'} < t, \ i' = i + 1}{X_{i} < t}$$

10 Decreasing($\{X_1, \dots, X_n\}$)

$$X_{i} \geq t \iff b_{it} \qquad \forall i \in [1, n] \forall t \in [1, m]$$

$$(b_{(i-1)t} \vee \neg b_{it}) \qquad \forall i \in [2, n] \forall t \in [1, m]$$

$$\frac{X_{i'} \geq t, \ i' = i + 1}{X_{i} \geq t}$$

$$\frac{X_{i'} < t, \ i' = i - 1}{X_{i} < t}$$

11 **Among** $(c, \{X_1, \ldots, X_n\}, D_4)$

$$\begin{split} X_i &= t \iff b_{it} & \forall i \in \llbracket 1, n \rrbracket \forall t \in \llbracket 1, m \rrbracket \\ \bigvee_{t \in D_4} b_{it} \iff b2_i & \forall i \in \llbracket 1, n \rrbracket \\ \sum_{i \in \llbracket 1, n \rrbracket} b2_i &= c \\ & \underbrace{X_i \neq t, \ \forall t, \ t \in D_4, \ \forall i, \ i \in \llbracket 1, n \rrbracket}_{X_i = t} \\ & \underbrace{X_i \neq t', \ \forall t', \ t' \neq t, \ t' \in D_4, \ t \in D_4}_{X_i = t} \\ & \underbrace{X_i = t, \ \exists t, \ t \in D_4, \ \forall i, \ i \in \llbracket 1, n \rrbracket}_{X_i \neq t} \end{split}$$

12 Roots $({X_1, ..., X_n}, I, V)$

$$X_{i} = t \iff b_{it} \qquad \forall it$$

$$\sum_{t \in V} b_{it} = 1 \qquad \forall i \in I$$

$$\sum_{t \in D \setminus V} b_{it} = 1 \qquad \forall i \in [1, n] \setminus I$$

$$V=D_5$$
 et $D\setminus V=D_6$
$$\frac{X_i\neq t,\ \forall t,\ t\in D_5}{X_i=t}$$

$$\frac{X_i\neq t,\ \forall t,\ t\in D_6}{X_i=t}$$

$$\begin{aligned} & X_i = t, \ \forall t, \ t \in D_5 \\ & X_i \neq t \\ & X_i = t, \ \forall t, \ t \in D_6 \\ & X_i \neq t \end{aligned}$$

13 Range($\{X_1, ..., X_n\}, I, V$)

$$X_i = t \iff b_{it}$$
 $\forall it$

$$\sum_{i \in I} b_{it} \ge 1$$
 $\forall t \in V$

$$\sum_{t \in V} b_{it} = 1$$
 $\forall i \in I$

 $I = D_5$ et $V = D_6$

$$\begin{aligned} X_i \neq t', \ \forall t', \ t' \neq t, \ t' \in D_5, \ t \in D_5 \\ X_i = t \\ X_i \neq t, \ \forall t, \ t \in D_6 \\ \overline{X_i = t} \\ X_i = t \\ X_i \neq t \end{aligned}$$

14 Xor(b1, b2, b3)

$$b1 = (b2 \neq b3)$$

b1,b2,b3 intervertibles

$$\frac{b2 - b3}{b1} \\
 \frac{b3 - b2}{b1} \\
 \frac{b2 \ b3}{\neg b1} \\
 \frac{-b2 \ \neg b3}{\neg b1}$$

15 **BinPacking**($\{X_1, \ldots, X_n\}, \{w_1, \ldots, w_n\}, c$)

$$X_i = p \iff b_{ip} \qquad \forall i \in [1, n] \forall p \in [1, n]$$
$$\sum_{i \in [1, n]} w_i \times b_{ip} \le c \qquad \forall p \in [1, n]$$

$$\frac{X_{i'} = p, \ \forall i', \ i' \neq i, \ i' \in \llbracket 1, n \rrbracket, \ i \in \llbracket 1, n \rrbracket}{X_i \neq p}$$