

CS70: Discrete Math and Probability

Fan Ye

June 23, 2016

More inductions!

Bunch of examples

More inductions!

Bunch of examples
Good ones

More inductions!

Bunch of examples
Good ones and bad ones

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case:

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp:

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps:

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1:

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp).

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have ...0**1001**0..., it cannot create two 1's in a row.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have ...0**1001**0..., it cannot create two 1's in a row.

Case 2:

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have $\dots 010010 \dots$, it cannot create two 1's in a row.

Case 2: If we choose a 0, and change it to 010,

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have ...0**1001**0..., it cannot create two 1's in a row.

Case 2: If we choose a 0, and change it to 010, then its previous neighbors will still have 0 as their neighbors.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have ...0**1001**0..., it cannot create two 1's in a row.

Case 2: If we choose a 0, and change it to 010, then its previous neighbors will still have 0 as their neighbors. Thus after the change we still do not have two 1's.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have ...0**1001**0..., it cannot create two 1's in a row.

Case 2: If we choose a 0, and change it to 010, then its previous neighbors will still have 0 as their neighbors. Thus after the change we still do not have two 1's.

Therefore after the $(n+1)_{th}$ step there are still not two 1's in a row.

More inductions

Suppose I start with 0 written on a piece of paper. Each time, I choose a digit written on the paper and erase it. If it was a 0, I replace it with 010. If it was a 1, I replace it with 1001. Prove that it's not possible for me to get two 1's in a row.

Induct on number of operations I make.

Base case: After the first step, we get 010, which does not have two 1's in a row.

Ind hyp: Assume after n steps, we do not have two 1's in a row.

Ind steps: At the $(n+1)_{th}$ step,

Case 1: If we choose a 1, its neighbor must be 0's (based on ind hyp). Therefore after the change we will have ...0**1001**0..., it cannot create two 1's in a row.

Case 2: If we choose a 0, and change it to 010, then its previous neighbors will still have 0 as their neighbors. Thus after the change we still do not have two 1's.

Therefore after the $(n+1)_{th}$ step there are still not two 1's in a row.

By principle of induction, ...



More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction?

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction?

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case:

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps:

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1:

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$,

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2:

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$,

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

Base on induction hypothesis, $n+1 - 2^k$ can be written as a sum of distinct powers of 2.

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

Base on induction hypothesis, $n+1 - 2^k$ can be written as a sum of distinct powers of 2.

Done?

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

Base on induction hypothesis, $n+1 - 2^k$ can be written as a sum of distinct powers of 2.

Done? Need to make sure 2^k is unique!

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

Base on induction hypothesis, $n+1 - 2^k$ can be written as a sum of distinct powers of 2.

Done? Need to make sure 2^k is unique!

Since $n+1 - 2^k < 2^{k+1} - 2^k < 2^k$, all terms must have power less than k ,

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

Base on induction hypothesis, $n+1 - 2^k$ can be written as a sum of distinct powers of 2.

Done? Need to make sure 2^k is unique!

Since $n+1 - 2^k < 2^{k+1} - 2^k < 2^k$, all terms must have power less than k , thus 2^k must be unique in this sum for $n+1$.

More power, more sum!

Theorem: Every positive integer n can be written as a sum of distinct powers of 2.

Discuss with your neighbors

Vanilla induction? Strong induction? Strengthen ind hyp?

Base case: For $n = 1 = 2^0$, $p(1)$ is true.

Induction hypothesis:

Assume all integers between 1 and n can be written as sums of distinct powers of 2.

Induction steps: Need to show that $n+1$ can be written as a sum of distinct powers of 2.

We can find a k such that $2^k \leq (n+1) < 2^{k+1}$

Case 1: $n+1 = 2^k$, we are done

Case 2: $2^k < (n+1) < 2^{k+1}$, then we have $n+1 = 2^k + (n+1 - 2^k)$

Base on induction hypothesis, $n+1 - 2^k$ can be written as a sum of distinct powers of 2.

Done? Need to make sure 2^k is unique!

Since $n+1 - 2^k < 2^{k+1} - 2^k < 2^k$, all terms must have power less than k , thus 2^k must be unique in this sum for $n+1$.

By principle of induction, ...



Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$.

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\sum_{i=1}^{k+1} \frac{1}{i^2}$$

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned} \sum_{i=1}^{k+1} \frac{1}{i^2} \\ = \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2}. \end{aligned}$$

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \cdot \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm...

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less?

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

$$"S_k \leq 2 - \frac{1}{(k+1)^2}"$$

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Induction step works!

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Induction step works! **No!**

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Induction step works! **No! Not the same statement!!!!**

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Induction step works! **No! Not the same statement!!!!**

Need to prove " $S_{k+1} \leq 2 - \frac{1}{(k+2)^2}$ ".

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Induction step works! **No! Not the same statement!!!!**

Need to prove " $S_{k+1} \leq 2 - \frac{1}{(k+2)^2}$ ".

Strengthening: need to...

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Base: $P(1)$. $1 \leq 2$.

Ind Step: $\sum_{i=1}^k \frac{1}{i^2} \leq 2$.

$$\begin{aligned}\sum_{i=1}^{k+1} \frac{1}{i^2} &= \sum_{i=1}^k \frac{1}{i^2} + \frac{1}{(k+1)^2} \\ &\leq 2 + \frac{1}{(k+1)^2}\end{aligned}$$

Uh oh?

Hmmm... It better be that any sum is *strictly less than* 2.

How much less? At least by $\frac{1}{(k+1)^2}$ for S_k .

" $S_k \leq 2 - \frac{1}{(k+1)^2}$ " \implies " $S_{k+1} \leq 2$ "

Induction step works! **No! Not the same statement!!!!**

Need to prove " $S_{k+1} \leq 2 - \frac{1}{(k+2)^2}$ ".

Ooops.....

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - \frac{1}{n}$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - \frac{1}{n}$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - \frac{1}{n}$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k)$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1)$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$S(k+1) = S_k + \frac{1}{(k+1)^2}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \end{aligned}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

Choose $f(k+1) \leq f(k) - \frac{1}{(k+1)^2}$.

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ — “} S_k \leq 2 - f(k) \text{”}$

Prove: $P(k+1) \text{ — “} S_{k+1} \leq 2 - f(k+1) \text{”}$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

$$1 \leq \frac{k+1}{k} - \frac{1}{k+1}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \text{ By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

$$1 \leq \frac{k+1}{k} - \frac{1}{k+1} \quad \text{Multiplied by } k+1.$$

$$1 \leq 1 + \left(\frac{1}{k} - \frac{1}{k+1} \right)$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ --- } "S_k \leq 2 - f(k)"$

Prove: $P(k+1) \text{ --- } "S_{k+1} \leq 2 - f(k+1)"$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \quad \text{By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

$$1 \leq \frac{k+1}{k} - \frac{1}{k+1} \quad \text{Multiplied by } k+1.$$

$$1 \leq 1 + \left(\frac{1}{k} - \frac{1}{k+1} \right) \quad \text{Some math.}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ — “} S_k \leq 2 - f(k) \text{”}$

Prove: $P(k+1) \text{ — “} S_{k+1} \leq 2 - f(k+1) \text{”}$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \quad \text{By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

$$1 \leq \frac{k+1}{k} - \frac{1}{k+1} \quad \text{Multiplied by } k+1.$$

$$1 \leq 1 + \left(\frac{1}{k} - \frac{1}{k+1} \right) \quad \text{Some math. So yes!}$$

Strengthening: how?

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - f(n)$. ($S_n = \sum_{i=1}^n \frac{1}{i^2}$.)

Proof:

Ind hyp: $P(k) \text{ — “} S_k \leq 2 - f(k) \text{”}$

Prove: $P(k+1) \text{ — “} S_{k+1} \leq 2 - f(k+1) \text{”}$

$$\begin{aligned} S(k+1) &= S_k + \frac{1}{(k+1)^2} \\ &\leq 2 - f(k) + \frac{1}{(k+1)^2} \quad \text{By ind. hyp.} \end{aligned}$$

$$\begin{aligned} \text{Choose } f(k+1) &\leq f(k) - \frac{1}{(k+1)^2}. \\ \implies S(k+1) &\leq 2 - f(k+1). \end{aligned}$$

Can you?

Subtracting off a quadratically decreasing function every time.

Maybe a linearly decreasing function to keep positive?

Try $f(k) = \frac{1}{k}$

$$\frac{1}{k+1} \leq \frac{1}{k} - \frac{1}{(k+1)^2} ?$$

$$1 \leq \frac{k+1}{k} - \frac{1}{k+1} \quad \text{Multiplied by } k+1.$$

$$1 \leq 1 + \left(\frac{1}{k} - \frac{1}{k+1} \right) \quad \text{Some math. So yes!}$$

Theorem: For all $n \geq 1$, $\sum_{i=1}^n \frac{1}{i^2} \leq 2 - \frac{1}{n}$.

Careful!



Horses of the same color...

Theorem: All horses have the same color.

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$. $1, 2, 3, \dots, k, k+1$

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$. 1, 2, 3, ..., $k, k+1$

Second k have same color by $P(k)$. 1, 2, 3, ..., $k, k+1$

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$. 1, 2, 3, ..., k , $k+1$

Second k have same color by $P(k)$. 1, 2, 3, ..., k , $k+1$

A horse in the middle in common! 1, 2, 3, ..., k , $k+1$

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$. $1, 2, 3, \dots, k, k+1$

Second k have same color by $P(k)$. $1, 2, 3, \dots, k, k+1$

A horse in the middle in common! $1, 2, 3, \dots, k, k+1$

All k must have the same color. $1, 2, 3, \dots, k, k+1$

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

How about $P(1) \implies P(2)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

How about $P(1) \implies P(2)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

1,2

Second k have same color by $P(k)$.

A horse in the middle in common!

How about $P(1) \implies P(2)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

1,2

Second k have same color by $P(k)$.

1,2

A horse in the middle in common!

How about $P(1) \implies P(2)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

1,2

Second k have same color by $P(k)$.

1,2

A horse in the middle in common!

1,2

No horse in common!

How about $P(1) \Rightarrow P(2)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

1,2

Second k have same color by $P(k)$.

1,2

A horse in the middle in common!

1,2

No horse in common!

How about $P(1) \Rightarrow P(2)$?

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

New Base Case: $P(2)$: there are two horses with same color.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

Fix base case.

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

New Base Case: $P(2)$: there are two horses with same color.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

Fix base case.

...Still doesn't work!!

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

New Base Case: $P(2)$: there are two horses with same color.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

Fix base case.

...Still doesn't work!!

(There are two horses is \neq For all two horses!!!)

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

New Base Case: $P(2)$: there are two horses with same color.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

Fix base case.

...Still doesn't work!!

(There are two horses is \neq For all two horses!!!)

Of course it doesn't work.

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

New Base Case: $P(2)$: there are two horses with same color.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$.

Second k have same color by $P(k)$.

A horse in the middle in common!

Fix base case.

...Still doesn't work!!

(There are two horses is \neq For all two horses!!!)

Of course it doesn't work.

As we will see, it is more subtle to catch errors in proofs of correct theorems!!

Use induction to prove the follow equality:

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Proof by induction:

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Proof by induction:

Base case:

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Proof by induction:

Base case: for $n = 1$, $1 = \sqrt{1+0} = 1$, equality holds.

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Proof by induction:

Base case: for $n = 1$, $1 = \sqrt{1+0} = 1$, equality holds.

Induction hypothesis:

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Proof by induction:

Base case: for $n = 1$, $1 = \sqrt{1+0} = 1$, equality holds.

Induction hypothesis: Assume this equality holds for $n = k$, i.e.

Use induction to prove the follow equality:

$$n = \sqrt{1 + (n-1)\sqrt{1 + n\sqrt{1 + (n+1)\sqrt{1 + (n+2)\dots}}}}$$

for all positive integers n .

Proof by induction:

Base case: for $n = 1$, $1 = \sqrt{1+0} = 1$, equality holds.

Induction hypothesis: Assume this equality holds for $n = k$, i.e.

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step:

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

By square both sides of the induction hypothesis we can get:

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

By square both sides of the induction hypothesis we can get:

$$k^2 = 1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

By square both sides of the induction hypothesis we can get:

$$k^2 = 1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Easy to get

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

By square both sides of the induction hypothesis we can get:

$$k^2 = 1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Easy to get

$$\frac{k^2 - 1}{k - 1} = k + 1 = \sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

By square both sides of the induction hypothesis we can get:

$$k^2 = 1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Easy to get

$$\frac{k^2 - 1}{k - 1} = k + 1 = \sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Therefore it holds for $n = k + 1$, by principle of induction, ...



Ind hyp:

$$k = \sqrt{1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}}$$

Induction step: Need to show it holds for $n = k + 1$.

By square both sides of the induction hypothesis we can get:

$$k^2 = 1 + (k-1)\sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Easy to get

$$\frac{k^2 - 1}{k - 1} = k + 1 = \sqrt{1 + k\sqrt{1 + (k+1)\sqrt{1 + (k+2)\dots}}}$$

Therefore it holds for $n = k + 1$, by principle of induction, ...

□ Good or bad?

Bad proof!

Bad proof! We need $k \neq 1$ to divide both sides by $k - 1$

Bad proof! We need $k \neq 1$ to divide both sides by $k - 1$

Or in other words, $p(1)$ does not imply $p(2)$

Bad proof! We need $k \neq 1$ to divide both sides by $k - 1$

Or in other words, $p(1)$ does not imply $p(2)$

Be careful.

Graphs!

Graphs!

Definitions: model.



Graphs!

Definitions: model.

Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?



Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).



Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?

Figure 1: "Konigsberg bridges" by Bogdan Giuscă - [License](#).





Can you draw a tour in the graph where you visit each edge once?

Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?

Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).





Can you draw a tour in the graph where you visit each edge once? Yes?

Konigsberg bridges problem.

Can you make a tour visiting each bridge exactly once?

Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).





Can you draw a tour in the graph where you visit each edge once? Yes? No?

Konigsberg bridges problem.


Can you make a tour visiting each bridge exactly once?

Figure 1: "Konigsberg bridges" by Bogdan Giușcă - [License](#).




Can you draw a tour in the graph where you visit each edge once? Yes? No?
We will see!

Graphs: formally.




Graph:

Graphs: formally.



Graph: $G = (V, E)$.


Graphs: formally.



Graph: $G = (V, E)$.

V - set of vertices.

Graphs: formally.




Graph: $G = (V, E)$.

V - set of vertices.

$\{A, B, C, D\}$

Graphs: formally.




Graph: $G = (V, E)$.

V - set of vertices.

$\{A, B, C, D\}$

$E \subseteq V \times V$ -

Graphs: formally.




Graph: $G = (V, E)$.

V - set of vertices.

$\{A, B, C, D\}$

$E \subseteq V \times V$ - set of edges.

Graphs: formally.



Graph: $G = (V, E)$.


V - set of vertices.

$\{A, B, C, D\}$

$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}$

Graphs: formally.



Graph: $G = (V, E)$.


V - set of vertices.

$\{A, B, C, D\}$

$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}, \{A, B\}\}$

Graphs: formally.



Graph: $G = (V, E)$.


V - set of vertices.

$\{A, B, C, D\}$

$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}, \{A, B\}, \{A, C\},$

Graphs: formally.



Graph: $G = (V, E)$.


V - set of vertices.

$\{A, B, C, D\}$

$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}, \{A, B\}, \{A, C\}, \{B, C\}, \{B, D\}, \{B, D\}, \{C, D\}\}$.

Graphs: formally.



Graph: $G = (V, E)$.

V - set of vertices.



$\{A, B, C, D\}$

$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}, \{A, B\}, \{A, C\}, \{B, C\}, \{B, D\}, \{B, D\}, \{C, D\}\}$.

For CS 70, usually simple graphs.

Graphs: formally.



Graph: $G = (V, E)$.

V - set of vertices.

$\{A, B, C, D\}$



$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}, \{A, B\}, \{A, C\}, \{B, C\}, \{B, D\}, \{B, D\}, \{C, D\}\}$.

For CS 70, usually simple graphs.

No parallel edges.

Graphs: formally.



Graph: $G = (V, E)$.

V - set of vertices.

$\{A, B, C, D\}$


$E \subseteq V \times V$ - set of edges.

$\{\{A, B\}, \{A, B\}, \{A, C\}, \{B, C\}, \{B, D\}, \{B, D\}, \{C, D\}\}$.

For CS 70, usually simple graphs.


No parallel edges.

Multigraph above.



$G = (V, E)$.


Directed Graphs



$G = (V, E)$.

V - set of vertices.

Directed Graphs




$G = (V, E).$

V - set of vertices.

$\{1, 2, 3, 4\}$

Directed Graphs




$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

Directed Graphs



$G = (V, E)$.


V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2),$

Directed Graphs



$G = (V, E)$.


V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3),$

Directed Graphs



$G = (V, E)$.


V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4),$

Directed Graphs



$G = (V, E)$.


V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

Directed Graphs



$G = (V, E)$.

V - set of vertices.


$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$


E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament:

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$


E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2,

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.


$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2, ...

Precedence:

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.


$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2, ...

Precedence: 1 is before 2,

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.


$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$


One way streets.

Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Social Network:

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$


One way streets.

Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Social Network: Directed?

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$


One way streets.

Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Social Network: Directed? Undirected?

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.


Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Social Network: Directed? Undirected?

Friends.

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.


Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Social Network: Directed? Undirected?

Friends. Undirected.

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2, ...


Precedence: 1 is before 2, ..

Social Network: Directed? Undirected?

Friends. Undirected.

Likes.

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2, ...


Precedence: 1 is before 2, ..

Social Network: Directed? Undirected?

Friends. Undirected.

Likes. Directed.

Directed Graphs



$G = (V, E)$.

V - set of vertices.

$\{1, 2, 3, 4\}$

E ordered pairs of vertices.

$\{(1, 2), (1, 3), (1, 4), (2, 4), (3, 4)\}$

One way streets.

Tournament: 1 beats 2, ...

Precedence: 1 is before 2, ..

Social Network: Directed? Undirected?

Friends. Undirected.

Likes. Directed.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

Graph Concepts and Definitions.



Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree





Neighbors of 10?

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree





Neighbors of 10? 1,

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree





Neighbors of 10? 1,5,

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree





Neighbors of 10? 1,5,7,

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree





Neighbors of 10? 1,5,7, 8.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree





Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.



u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.



Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.



Edge (u, v) is incident to u and v .

Degree of vertex 1?

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.



Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .



Degree of vertex 1? 2

Degree of vertex u is number of incident edges.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2



Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2



Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Degree of vertex u is number of incident edges.



Equals number of neighbors in simple graph.

Directed graph?

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.



Directed graph?

In-degree of 10?

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.



Directed graph?

In-degree of 10? 1

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.



Directed graph?

In-degree of 10? 1 Out-degree of 10?

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.



Directed graph?

In-degree of 10? 1 Out-degree of 10? 3

Graph Concepts and Definitions.

Graph: $G = (V, E)$

neighbors, adjacent, degree, incident, in-degree, out-degree



Neighbors of 10? 1,5,7, 8.

u is neighbor of v if $(u, v) \in E$.

Neighbors: All vertices that are adjacent to a vertex.

Edge (10,5) is incident to vertex 10 and vertex 5.

Edge (u, v) is incident to u and v .

Degree of vertex 1? 2

Degree of vertex u is number of incident edges.

Equals number of neighbors in simple graph.

Directed graph?

In-degree of 10? 1 Out-degree of 10? 3

Quick Proof.

The sum of the vertex degrees is equal to

Quick Proof.

The sum of the vertex degrees is equal to

(A) the total number of vertices, $|V|$.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)!

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?


Not (A)! Triangle.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.




Not (B)!

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.




Not (B)! Triangle.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.




Not (B)! Triangle.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.


What?

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.


What? For triangle number of edges is 3, the sum of degrees is 6.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.


Could it always be...

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.


Could it always be... $2|E|$?

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?


How many incidences does each edge contribute?

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?


How many incidences does each edge contribute? 2.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?

How many incidences does each edge contribute? 2.


$2|E|$ incidences are contributed in total!

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?

How many incidences does each edge contribute? 2.

$2|E|$ incidences are contributed in total!


What is degree v ?

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?

How many incidences does each edge contribute? 2.

$2|E|$ incidences are contributed in total!


What is degree v ? incidences contributed to v !

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?

How many incidences does each edge contribute? 2.

$2|E|$ incidences are contributed in total!

What is degree v ? incidences contributed to v !


sum of degrees is total incidences

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?

How many incidences does each edge contribute? 2.

$2|E|$ incidences are contributed in total!

What is degree v ? incidences contributed to v !


sum of degrees is total incidences ... or $2|E|$.

Quick Proof.

The sum of the vertex degrees is equal to

- (A) the total number of vertices, $|V|$.
- (B) the total number of edges, $|E|$.
- (C) What?

Not (A)! Triangle.



Not (B)! Triangle.

What? For triangle number of edges is 3, the sum of degrees is 6.

Could it always be... $2|E|$?

How many incidences does each edge contribute? 2.


$2|E|$ incidences are contributed in total!

What is degree v ? incidences contributed to v !

sum of degrees is total incidences ... or $2|E|$.


Thm: Sum of vertex degree is $2|E|$.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.


Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path?


Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1,10\}$, $\{8,5\}$, $\{4,5\}$?


Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1,10\}$, $\{8,5\}$, $\{4,5\}$? No!

Paths, walks, cycles, tour.




A path in a graph is a sequence of edges.

Path? $\{1, 10\}$, $\{8, 5\}$, $\{4, 5\}$? No!

Path?

Paths, walks, cycles, tour.




A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$?

Paths, walks, cycles, tour.




A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Paths, walks, cycles, tour.




A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.


Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check!

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.


Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path?

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.


Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.


Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!


Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!


Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle?

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!


Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!


Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!


Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k).$


Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!

Walk is sequence of edges with possible repeated vertex or edge.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.


Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!

Walk is sequence of edges with possible repeated vertex or edge.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}$, $\{8, 5\}$, $\{4, 5\}$? No!

Path? $\{1, 10\}$, $\{10, 5\}$, $\{5, 4\}$, $\{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.


Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!

Walk is sequence of edges with possible repeated vertex or edge.

Tour is walk that starts and ends at the same node.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.


Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!

Walk is sequence of edges with possible repeated vertex or edge.

Tour is walk that starts and ends at the same node.

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!


Path is usually simple. No repeated vertex!

Walk is sequence of edges with possible repeated vertex or edge.

Tour is walk that starts and ends at the same node.

Quick Check!

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!


Walk is sequence of edges with possible repeated vertex or edge.

Tour is walk that starts and ends at the same node.

Quick Check!

Path is to Walk as Cycle is to ??

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!


Walk is sequence of edges with possible repeated vertex or edge.

Tour is walk that starts and ends at the same node.

Quick Check!

Path is to Walk as Cycle is to ?? Tour!

Paths, walks, cycles, tour.



A path in a graph is a sequence of edges.

Path? $\{1, 10\}, \{8, 5\}, \{4, 5\}$? No!

Path? $\{1, 10\}, \{10, 5\}, \{5, 4\}, \{4, 11\}$? Yes!

Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Quick Check! Length of path? k vertices or $k - 1$ edges.

Cycle: Path with $v_1 = v_k$. Length of cycle? $k - 1$ vertices and edges!

Path is usually simple. No repeated vertex!


Walk is sequence of edges with possible repeated vertex or edge.

Tour is walk that starts and ends at the same node.


Quick Check!

Path is to Walk as Cycle is to ?? Tour!

Directed Paths.




Directed Paths.




Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Directed Paths.




Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Directed Paths.




Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Directed Paths.



Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.


Directed Paths.



Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Paths,


Directed Paths.



Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Paths, walks,


Directed Paths.



Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Paths, walks, cycles,


Directed Paths.



Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.

Paths, walks, cycles, tours


Directed Paths.



Path: $(v_1, v_2), (v_2, v_3), \dots, (v_{k-1}, v_k)$.


Paths, walks, cycles, tours ... are analogous to undirected now.

Connectivity



u and v are **connected** if there is a path between u and v .


Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

Connectivity




u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Connectivity




u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected?

Connectivity




u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes?

Connectivity




u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Connectivity



u and v are **connected** if there is a path between u and v .


A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Proof:

Connectivity



u and v are **connected** if there is a path between u and v .


A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Proof: Use path from u to x and then from x to v .

Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.


If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Proof: Use path from u to x and then from x to v .



Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.


Is graph connected? Yes? No?

Proof: Use path from u to x and then from x to v .



May not be simple!

Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?


Proof: Use path from u to x and then from x to v .



May not be simple!

Either modify definition to walk.

Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Proof: Use path from u to x and then from x to v .




May not be simple!

Either modify definition to walk.

Or cut out cycles.

Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Proof: Use path from u to x and then from x to v .




May not be simple!

Either modify definition to walk.

Or cut out cycles. .

Connectivity



u and v are **connected** if there is a path between u and v .

A connected graph is a graph where all pairs of vertices are connected.

If one vertex x is connected to every other vertex.

Is graph connected? Yes? No?

Proof: Use path from u to x and then from x to v .




May not be simple!

Either modify definition to walk.


Or cut out cycles. .

Connected component




Is graph above connected?

Connected component



Is graph above connected? Yes!


Connected component



Is graph above connected? Yes!

How about now?


Connected component



Is graph above connected? Yes!

How about now? No!

Connected component




Is graph above connected? Yes!

How about now? No!

Connected Components?

Connected component




Is graph above connected? Yes!

How about now? No!

Connected Components? $\{1\}, \{10, 7, 5, 8, 4, 3, 11\}, \{2, 9, 6\}$.

Connected component




Is graph above connected? Yes!

How about now? No!

Connected Components? $\{1\}, \{10, 7, 5, 8, 4, 3, 11\}, \{2, 9, 6\}$.

Connected component - maximal set of connected vertices.

Connected component



Is graph above connected? Yes!


How about now? No!

Connected Components? $\{1\}, \{10, 7, 5, 8, 4, 3, 11\}, \{2, 9, 6\}$.

Connected component - maximal set of connected vertices.

Quick Check: Is $\{10, 7, 5\}$ a connected component?

Connected component



Is graph above connected? Yes!

How about now? No!

Connected Components? $\{1\}, \{10, 7, 5, 8, 4, 3, 11\}, \{2, 9, 6\}$.

Connected component - maximal set of connected vertices.

Quick Check: Is $\{10, 7, 5\}$ a connected component? No.

Thank you!

Congrats on surviving the first week!

Thank you!

Congrats on surviving the first week! Have a good weekend!

Thank you!

Congrats on surviving the first week! Have a good weekend!

Don't forget your homework, homework party tonight.