

CS70: Alex Psomas: Lecture 13.

Modeling Uncertainty: Probability Space

CS70: Alex Psomas: Lecture 13.

Modeling Uncertainty: Probability Space

1. Key Points
2. Random Experiments
3. Probability Space
4. Events

Key Points

Key Points

- ▶ Uncertainty does not mean “nothing is known”

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?
 - ▶ Play games of chance.

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?
 - ▶ Play games of chance.
 - ▶ Design randomized algorithms.

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?
 - ▶ Play games of chance.
 - ▶ Design randomized algorithms.
 - ▶ Catch Pokemon.

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?
 - ▶ Play games of chance.
 - ▶ Design randomized algorithms.
 - ▶ Catch Pokemon.
- ▶ Probability

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?
 - ▶ Play games of chance.
 - ▶ Design randomized algorithms.
 - ▶ Catch Pokemon.
- ▶ Probability
 - ▶ Models knowledge about uncertainty

Key Points

- ▶ Uncertainty does not mean “nothing is known”
- ▶ How to best make decisions under uncertainty?
 - ▶ Buy stocks
 - ▶ Detect signals (transmitted bits, speech, images, radar, diseases, etc.)
 - ▶ Control systems (Internet, airplane, robots, self-driving cars, schedule surgeries in a hospital, etc.)
- ▶ How to best use ‘artificial’ uncertainty?
 - ▶ Play games of chance.
 - ▶ Design randomized algorithms.
 - ▶ Catch Pokemon.
- ▶ Probability
 - ▶ Models knowledge about uncertainty
 - ▶ Discovers best way to use that knowledge in making decisions

The Magic of Probability

The Magic of Probability

Uncertainty:

The Magic of Probability

Uncertainty: vague,

The Magic of Probability

Uncertainty: vague, fuzzy,

The Magic of Probability

Uncertainty: vague, fuzzy, confusing,

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary,

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability:

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise,

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous,

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear



Probability = Serenity

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear



Probability = Serenity

Our mission: help you discover the magic of Probability,

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear



Probability = Serenity

Our mission: help you discover the magic of Probability, i.e., enable you to think clearly about uncertainty.

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear



Probability = Serenity

Our mission: help you discover the magic of Probability, i.e., enable you to think clearly about uncertainty.

Your cost:

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear



Probability = Serenity

Our mission: help you discover the magic of Probability, i.e., enable you to think clearly about uncertainty.

Your cost: focused attention

The Magic of Probability

Uncertainty: vague, fuzzy, confusing, scary, hard to think about.

Probability: A precise, unambiguous, simple way to think about uncertainty.



Uncertainty = Fear



Probability = Serenity

Our mission: help you discover the magic of Probability, i.e., enable you to think clearly about uncertainty.

Your cost: focused attention and practice on examples and problems.

A cool trick

Random Experiment: Flip one Fair Coin

Random Experiment: Flip one Fair Coin

Flip a fair coin:

Random Experiment: Flip one Fair Coin

Flip a fair coin: (*One flips or tosses a coin*)

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



- Possible outcomes:

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



- Possible outcomes: Heads (H)

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



- Possible outcomes: Heads (H) and Tails (T)

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



- Possible outcomes: Heads (H) and Tails (T)
(*One flip yields either 'heads' or 'tails'.*)

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



- ▶ Possible outcomes: Heads (H) and Tails (T)
(*One flip yields either 'heads' or 'tails'.*)
- ▶ Likelihoods:

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: (*One flips or tosses a coin*)



- ▶ Possible outcomes: Heads (H) and Tails (T)
(*One flip yields either 'heads' or 'tails'.*)
- ▶ Likelihoods: H : 50% and T : 50%

Random Experiment: Flip one Fair Coin

Flip a fair coin:



What do we mean by the likelihood of tails is 50%?

Random Experiment: Flip one Fair Coin

Flip a **fair** coin:



What do we mean by **the likelihood of tails is 50%**?

Two interpretations:

Random Experiment: Flip one Fair Coin

Flip a **fair** coin:



What do we mean by **the likelihood of tails is 50%**?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Random Experiment: Flip one Fair Coin

Flip a fair coin:



What do we mean by the likelihood of tails is 50%?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

Random Experiment: Flip one Fair Coin

Flip a **fair** coin:



What do we mean by **the likelihood of tails is 50%**?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

- ▶ Many coin flips: About half yield 'tails'

Random Experiment: Flip one Fair Coin

Flip a **fair** coin:



What do we mean by **the likelihood of tails is 50%**?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

- ▶ Many coin flips: About half yield 'tails'

Makes sense for many flips

Random Experiment: Flip one Fair Coin

Flip a fair coin:



What do we mean by the likelihood of tails is 50%?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

- ▶ Many coin flips: About half yield 'tails'

Makes sense for many flips

- ▶ Question:

Random Experiment: Flip one Fair Coin

Flip a **fair** coin:



What do we mean by **the likelihood of tails is 50%**?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

- ▶ Many coin flips: About half yield 'tails'

Makes sense for many flips

- ▶ Question: Why does the fraction of tails converge to the same value every time?

Random Experiment: Flip one Fair Coin

Flip a fair coin:



What do we mean by the likelihood of tails is 50%?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

- ▶ Many coin flips: About half yield 'tails'

Makes sense for many flips

- ▶ Question: Why does the fraction of tails converge to the same value every time? Statistical Regularity!

Random Experiment: Flip one Fair Coin

Flip a fair coin:



What do we mean by the likelihood of tails is 50%?

Two interpretations:

- ▶ Single coin flip: 50% chance of 'tails'

Willingness to bet on the outcome of a single flip

- ▶ Many coin flips: About half yield 'tails'

Makes sense for many flips

- ▶ Question: Why does the fraction of tails converge to the same value every time? Statistical Regularity! Deep!

Random Experiment: Flip one Fair Coin

Flip a fair coin:

Random Experiment: Flip one Fair Coin


Flip a fair coin: model

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: model



Physical Experiment




Probability Model

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: model



Physical Experiment



Probability Model


- The physical experiment is complex.

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: model



Physical Experiment



Probability Model


- The physical experiment is complex. (Shape, density, initial momentum and position, ...)

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: model



Physical Experiment



Probability Model


- ▶ The physical experiment is complex. (Shape, density, initial momentum and position, ...)
- ▶ The Probability model is simple:

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: model



Physical Experiment



Probability Model


- ▶ The physical experiment is complex. (Shape, density, initial momentum and position, ...)
- ▶ The Probability model is simple:
 - ▶ A set Ω of **outcomes**: $\Omega = \{H, T\}$.

Random Experiment: Flip one Fair Coin

Flip a **fair** coin: model



Physical Experiment



Probability Model

- ▶ The physical experiment is complex. (Shape, density, initial momentum and position, ...)
- ▶ The Probability model is simple:
 - ▶ A set Ω of **outcomes**: $\Omega = \{H, T\}$.
 - ▶ A **probability** assigned to each outcome:
 $Pr[H] = 0.5, Pr[T] = 0.5$.

Random Experiment: Flip one Unfair Coin

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- Possible outcomes:

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- Possible outcomes: Heads (H) and Tails (T)

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods:

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$
- ▶ Frequentist Interpretation:

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$
- ▶ Frequentist Interpretation:

Flip many times \Rightarrow Fraction $1 - p$ of tails

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$
- ▶ Frequentist Interpretation:
Flip many times \Rightarrow Fraction $1 - p$ of tails
- ▶ Question:

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$
- ▶ Frequentist Interpretation:
Flip many times \Rightarrow Fraction $1 - p$ of tails
- ▶ Question: How can one figure out p ?

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$
- ▶ Frequentist Interpretation:
Flip many times \Rightarrow Fraction $1 - p$ of tails
- ▶ Question: How can one figure out p ? Flip many times

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin:



H: 45%

T: 55%

- ▶ Possible outcomes: Heads (H) and Tails (T)
- ▶ Likelihoods: $H : p \in (0, 1)$ and $T : 1 - p$
- ▶ Frequentist Interpretation:
 - Flip many times \Rightarrow Fraction $1 - p$ of tails
- ▶ Question: How can one figure out p ? Flip many times
- ▶ Tautology?

Random Experiment: Flip one Unfair Coin

Random Experiment: Flip one Unfair Coin


Flip an **unfair** (biased, loaded) coin: model

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin: model



Physical Experiment




Probability Model

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin: model



Physical Experiment



Probability Model


- Same set of outcomes as before!

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin: model



Physical Experiment



Probability Model


- ▶ Same set of outcomes as before!
- ▶ Different probabilities!

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin: model



Physical Experiment



Probability Model


- ▶ Same set of outcomes as before!
- ▶ Different probabilities!
- ▶ The most common mistake in Probability:

Random Experiment: Flip one Unfair Coin

Flip an **unfair** (biased, loaded) coin: model



Physical Experiment



Probability Model

- ▶ Same set of outcomes as before!
- ▶ Different probabilities!
- ▶ The most common mistake in Probability: **assuming that outcomes are equally likely.**

Flip Two Fair Coins

Flip Two Fair Coins

- ▶ Possible outcomes:

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\}$

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\} \equiv \{H, T\}^2$.

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\} \equiv \{H, T\}^2$.
- ▶ Note: $A \times B := \{(a, b) \mid a \in A, b \in B\}$

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\} \equiv \{H, T\}^2$.
- ▶ Note: $A \times B := \{(a, b) \mid a \in A, b \in B\}$ and $A^2 := A \times A$.

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\} \equiv \{H, T\}^2$.
- ▶ Note: $A \times B := \{(a, b) \mid a \in A, b \in B\}$ and $A^2 := A \times A$.
- ▶ Likelihoods:

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\} \equiv \{H, T\}^2$.
- ▶ Note: $A \times B := \{(a, b) \mid a \in A, b \in B\}$ and $A^2 := A \times A$.
- ▶ Likelihoods: 1/4 each.

Flip Two Fair Coins

- ▶ Possible outcomes: $\{HH, HT, TH, TT\} \equiv \{H, T\}^2$.
- ▶ Note: $A \times B := \{(a, b) \mid a \in A, b \in B\}$ and $A^2 := A \times A$.
- ▶ Likelihoods: 1/4 each.



Flip Glued Coins

Flip Glued Coins

Flips two coins glued together side by side:

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- Possible outcomes:

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- Possible outcomes: $\{HH, TT\}$.

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- ▶ Possible outcomes: $\{HH, TT\}$.
- ▶ Likelihoods:

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- ▶ Possible outcomes: $\{HH, TT\}$.
- ▶ Likelihoods: $HH : 0.5, TT : 0.5$.

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- ▶ Possible outcomes: $\{HH, TT\}$.
- ▶ Likelihoods: $HH : 0.5, TT : 0.5$.
- ▶ Note: Coins are glued so that they show the same face.

Flip Glued Coins

Flip Glued Coins

Flips two coins glued together side by side:

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- Possible outcomes:

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- Possible outcomes: $\{HT, TH\}$.

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- ▶ Possible outcomes: $\{HT, TH\}$.
- ▶ Likelihoods:

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- ▶ Possible outcomes: $\{HT, TH\}$.
- ▶ Likelihoods: $HT : 0.5, TH : 0.5$.

Flip Glued Coins

Flips two coins glued together side by side:



Glued coins



50%



50%

- ▶ Possible outcomes: $\{HT, TH\}$.
- ▶ Likelihoods: $HT : 0.5, TH : 0.5$.
- ▶ Note: Coins are glued so that they show different faces.


Flip two Attached Coins

Flip two Attached Coins

Flips two coins attached by a spring:

Flip two Attached Coins

Flips two coins attached by a spring:



Flip two Attached Coins

Flips two coins attached by a spring:



- Possible outcomes:

Flip two Attached Coins

Flips two coins attached by a spring:



- Possible outcomes: $\{HH, HT, TH, TT\}$.

Flip two Attached Coins


Flips two coins attached by a spring:



- ▶ Possible outcomes: $\{HH, HT, TH, TT\}$.
- ▶ Likelihoods:

Flip two Attached Coins


Flips two coins attached by a spring:



- ▶ Possible outcomes: $\{HH, HT, TH, TT\}$.
- ▶ Likelihoods: $HH : 0.4, HT : 0.1, TH : 0.1, TT : 0.4$.

Flip two Attached Coins

Flips two coins attached by a spring:



- ▶ Possible outcomes: $\{HH, HT, TH, TT\}$.
- ▶ Likelihoods: $HH : 0.4, HT : 0.1, TH : 0.1, TT : 0.4$.
- ▶ Note: Coins are attached so that they tend to show the same face, unless the spring twists enough.


Flipping Two Coins

Flipping Two Coins


Here is a way to summarize the four random experiments:

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]




[3]




[4]

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]



[3]




[4]


- Ω is the set of *possible* outcomes;

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]



[3]







[4]

- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);

Flipping Two Coins





Here is a way to summarize the four random experiments:



- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;

Flipping Two Coins





Here is a way to summarize the four random experiments:



- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;
- ▶ Fair coins:

Flipping Two Coins





Here is a way to summarize the four random experiments:



- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;
- ▶ Fair coins: [1];

Flipping Two Coins





Here is a way to summarize the four random experiments:



- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;
- ▶ Fair coins: [1]; Glued coins:

Flipping Two Coins





Here is a way to summarize the four random experiments:



- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;
- ▶ Fair coins: [1]; Glued coins: [3], [4];

Flipping Two Coins

Here is a way to summarize the four random experiments:







- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;
- ▶ Fair coins: [1]; Glued coins: [3], [4];

Spring-attached coins:

Flipping Two Coins


Here is a way to summarize the four random experiments:




- ▶ Ω is the set of *possible* outcomes;
- ▶ Each outcome has a **probability** (likelihood);
- ▶ The probabilities are ≥ 0 and add up to 1;
- ▶ Fair coins: [1]; Glued coins: [3], [4];
Spring-attached coins: [2];

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]




[3]




[4]

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]



[3]




[4]


Important remarks:

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]



[3]




[4]

Important remarks:

- Each outcome describes the **two** coins.

Flipping Two Coins

Here is a way to summarize the four random experiments:




Important remarks:


- ▶ Each outcome describes the **two** coins.
- ▶ E.g., HT is **one** outcome of the experiment.

Flipping Two Coins


Here is a way to summarize the four random experiments:




[1]



[2]



[3]




[4]

Important remarks:

- ▶ Each outcome describes the **two** coins.
- ▶ E.g., HT is **one** outcome of the experiment.
- ▶ It is **wrong** to think that the outcomes are $\{H, T\}$ and that one picks twice from that set.

Flipping Two Coins

Here is a way to summarize the four random experiments:




Important remarks:

- ▶ Each outcome describes the **two** coins.
- ▶ E.g., HT is **one** outcome of the experiment.
- ▶ It is **wrong** to think that the outcomes are $\{H, T\}$ and that one picks twice from that set.
- ▶ This viewpoint misses the relationship between the two flips.

Flipping Two Coins

Here is a way to summarize the four random experiments:




Important remarks:

- ▶ Each outcome describes the **two** coins.
- ▶ E.g., HT is **one** outcome of the experiment.
- ▶ It is **wrong** to think that the outcomes are $\{H, T\}$ and that one picks twice from that set.
- ▶ This viewpoint misses the relationship between the two flips.
- ▶ Each $\omega \in \Omega$ describes one outcome of the **complete** experiment.

Flipping Two Coins

Here is a way to summarize the four random experiments:



Important remarks:

- ▶ Each outcome describes the **two** coins.
- ▶ E.g., *HT* is **one** outcome of the experiment.
- ▶ It is **wrong** to think that the outcomes are $\{H, T\}$ and that one picks twice from that set.
- ▶ This viewpoint misses the relationship between the two flips.
- ▶ Each $\omega \in \Omega$ describes one outcome of the **complete** experiment.
- ▶ Ω and the probabilities specify the random experiment.

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- Possible outcomes:

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- ▶ Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

- ▶ Note: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\} = \{H, T\}^n$.

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- ▶ Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

- ▶ Note: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\} = \{H, T\}^n$.

$$A^n := \{(a_1, \dots, a_n) \mid a_1 \in A, \dots, a_n \in A\}.$$

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- ▶ Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

- ▶ Note: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\} = \{H, T\}^n$.

$$A^n := \{(a_1, \dots, a_n) \mid a_1 \in A, \dots, a_n \in A\}. \quad |A^n| = |A|^n.$$

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- ▶ Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

- ▶ Note: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\} = \{H, T\}^n$.

$$A^n := \{(a_1, \dots, a_n) \mid a_1 \in A, \dots, a_n \in A\}. \quad |A^n| = |A|^n.$$

- ▶ Likelihoods:

Flipping n times

Flip a fair coin n times (some $n \geq 1$):

- ▶ Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

- ▶ Note: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\} = \{H, T\}^n$.

$$A^n := \{(a_1, \dots, a_n) \mid a_1 \in A, \dots, a_n \in A\}. \quad |A^n| = |A|^n.$$

- ▶ Likelihoods: $1/2^n$ each.

Flipping n times

Flip a **fair** coin n times (some $n \geq 1$):


- Possible outcomes: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\}$.

Thus, 2^n possible outcomes.

- Note: $\{TT \cdots T, TT \cdots H, \dots, HH \cdots H\} = \{H, T\}^n$.

$A^n := \{(a_1, \dots, a_n) \mid a_1 \in A, \dots, a_n \in A\}$. $|A^n| = |A|^n$.

- Likelihoods: $1/2^n$ each.



Roll two Dice

Roll a **balanced** 6-sided die twice:

Roll two Dice

Roll a **balanced** 6-sided die twice:

- ▶ Possible outcomes:

Roll two Dice

Roll a **balanced** 6-sided die twice:

- Possible outcomes:

$$\{1, 2, 3, 4, 5, 6\}^2 = \{(a, b) \mid 1 \leq a, b \leq 6\}.$$

Roll two Dice

Roll a **balanced** 6-sided die twice:

- ▶ Possible outcomes:
 $\{1, 2, 3, 4, 5, 6\}^2 = \{(a, b) \mid 1 \leq a, b \leq 6\}.$
- ▶ Likelihoods:

Roll two Dice


Roll a **balanced** 6-sided die twice:

- ▶ Possible outcomes:
 $\{1, 2, 3, 4, 5, 6\}^2 = \{(a, b) \mid 1 \leq a, b \leq 6\}.$
- ▶ Likelihoods: $1/36$ for each.

Roll two Dice

Roll a **balanced** 6-sided die twice:

- ▶ Possible outcomes:
 $\{1, 2, 3, 4, 5, 6\}^2 = \{(a, b) \mid 1 \leq a, b \leq 6\}.$
- ▶ Likelihoods: $1/36$ for each.



Probability Space.

1. A “random experiment”:

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;
 - (c) Deal a poker hand.

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;
 - (c) Deal a poker hand.
2. A set of possible outcomes: Ω .

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;
 - (c) Deal a poker hand.
2. A set of possible outcomes: Ω .
 - (a) $\Omega = \{H, T\}$;

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;
 - (c) Deal a poker hand.
2. A set of possible outcomes: Ω .
 - (a) $\Omega = \{H, T\}$;
 - (b) $\Omega = \{HH, HT, TH, TT\}$;

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;
 - (c) Deal a poker hand.
2. A set of possible outcomes: Ω .
 - (a) $\Omega = \{H, T\}$;
 - (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| =$

Probability Space.

1. A “random experiment”:
 - (a) Flip a biased coin;
 - (b) Flip two fair coins;
 - (c) Deal a poker hand.
2. A set of possible outcomes: Ω .
 - (a) $\Omega = \{H, T\}$;
 - (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| = 4$;

Probability Space.

1. A “random experiment”:

- (a) Flip a biased coin;
- (b) Flip two fair coins;
- (c) Deal a poker hand.

2. A set of possible outcomes: Ω .

- (a) $\Omega = \{H, T\}$;
- (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| = 4$;
- (c) $\Omega = \{ \underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit K\spadesuit}, \underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit Q\spadesuit}, \dots \}$
 $|\Omega| =$

Probability Space.

1. A “random experiment”:

- (a) Flip a biased coin;
- (b) Flip two fair coins;
- (c) Deal a poker hand.

2. A set of possible outcomes: Ω .

- (a) $\Omega = \{H, T\}$;
- (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| = 4$;
- (c) $\Omega = \{ \underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit K\spadesuit}, \underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit Q\spadesuit}, \dots \}$
 $|\Omega| = \binom{52}{5}$.

Probability Space.

1. A “random experiment”:

- (a) Flip a biased coin;
- (b) Flip two fair coins;
- (c) Deal a poker hand.

2. A set of possible outcomes: Ω .

- (a) $\Omega = \{H, T\}$;
- (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| = 4$;
- (c) $\Omega = \{ \underline{A\spadesuit A\heartsuit A\clubsuit A\heartsuit K\spadesuit}, \underline{A\spadesuit A\heartsuit A\clubsuit A\heartsuit Q\spadesuit}, \dots \}$
 $|\Omega| = \binom{52}{5}$.

3. Assign a probability to each outcome: $Pr : \Omega \rightarrow [0, 1]$.

- (a) $Pr[H] = p, Pr[T] = 1 - p$ for some $p \in [0, 1]$

Probability Space.

1. A “random experiment”:

- (a) Flip a biased coin;
- (b) Flip two fair coins;
- (c) Deal a poker hand.

2. A set of possible outcomes: Ω .

- (a) $\Omega = \{H, T\}$;
- (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| = 4$;
- (c) $\Omega = \{ \underline{A\spadesuit A\heartsuit A\clubsuit A\heartsuit K\spadesuit}, \underline{A\spadesuit A\heartsuit A\clubsuit A\heartsuit Q\spadesuit}, \dots \}$
 $|\Omega| = \binom{52}{5}$.

3. Assign a probability to each outcome: $Pr : \Omega \rightarrow [0, 1]$.

- (a) $Pr[H] = p, Pr[T] = 1 - p$ for some $p \in [0, 1]$
- (b) $Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}$

Probability Space.

1. A “random experiment”:

- (a) Flip a biased coin;
- (b) Flip two fair coins;
- (c) Deal a poker hand.

2. A set of possible outcomes: Ω .

- (a) $\Omega = \{H, T\}$;
- (b) $\Omega = \{HH, HT, TH, TT\}$; $|\Omega| = 4$;
- (c) $\Omega = \{ \underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit K\spadesuit}, \underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit Q\spadesuit}, \dots \}$
 $|\Omega| = \binom{52}{5}.$

3. Assign a probability to each outcome: $Pr : \Omega \rightarrow [0, 1]$.

- (a) $Pr[H] = p, Pr[T] = 1 - p$ for some $p \in [0, 1]$
- (b) $Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}$
- (c) $Pr[\underline{A\spadesuit A\diamondsuit A\clubsuit A\heartsuit K\spadesuit}] = \dots = 1 / \binom{52}{5}$

Probability Space: formalism.

Ω is the **sample space**.

Probability Space: formalism.

Ω is the **sample space**.

$\omega \in \Omega$ is a **sample point**.

Probability Space: formalism.

Ω is the **sample space**.

$\omega \in \Omega$ is a **sample point**. (Also called an **outcome**.)

Probability Space: formalism.

Ω is the **sample space**.

$\omega \in \Omega$ is a **sample point**. (Also called an **outcome**.)

Sample point ω has a probability $Pr[\omega]$ where

Probability Space: formalism.

Ω is the **sample space**.

$\omega \in \Omega$ is a **sample point**. (Also called an **outcome**.)

Sample point ω has a probability $Pr[\omega]$ where

- ▶ $0 \leq Pr[\omega] \leq 1$;

Probability Space: formalism.

Ω is the **sample space**.

$\omega \in \Omega$ is a **sample point**. (Also called an **outcome**.)

Sample point ω has a probability $Pr[\omega]$ where

- ▶ $0 \leq Pr[\omega] \leq 1$;
- ▶ $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Probability Space: formalism.


Ω is the **sample space**.

$\omega \in \Omega$ is a **sample point**. (Also called an **outcome**.)

Sample point ω has a probability $Pr[\omega]$ where

- ▶ $0 \leq Pr[\omega] \leq 1$;
- ▶ $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Sample Space



Samples (Outcomes)

$$0 \leq Pr[\omega] \leq 1$$

$$\sum_{\omega} Pr[\omega] = 1$$


Probability Space: Formalism.

In a **uniform probability space** each outcome ω is **equally probable**: $Pr[\omega] = \frac{1}{|\Omega|}$ for all $\omega \in \Omega$.

Probability Space: Formalism.

In a **uniform probability space** each outcome ω is **equally probable**: $Pr[\omega] = \frac{1}{|\Omega|}$ for all $\omega \in \Omega$.


Uniform Probability Space



Probability Space: Formalism.

In a **uniform probability space** each outcome ω is **equally probable**: $Pr[\omega] = \frac{1}{|\Omega|}$ for all $\omega \in \Omega$.

Uniform Probability Space




Examples:

- ▶ Flipping two fair coins, dealing a poker hand are uniform probability spaces.

Probability Space: Formalism.

In a **uniform probability space** each outcome ω is **equally probable**: $Pr[\omega] = \frac{1}{|\Omega|}$ for all $\omega \in \Omega$.

Uniform Probability Space



Examples:

- ▶ Flipping two fair coins, dealing a poker hand are uniform probability spaces.
- ▶ Flipping a biased coin is not a uniform probability space.

Probability Space: Formalism


Simplest physical model of a **uniform** probability space:

Probability Space: Formalism

Simplest physical model of a **uniform** probability space:



Physical experiment




Probability model

Probability Space: Formalism

Simplest physical model of a **uniform** probability space:



Physical experiment



Probability model


A bag of identical balls, except for their color (or a label).

Probability Space: Formalism

Simplest physical model of a **uniform** probability space:



Physical experiment



Probability model


A bag of identical balls, except for their color (or a label). If the bag is well shaken, every ball is equally likely to be picked.

Probability Space: Formalism

Simplest physical model of a **uniform** probability space:



Physical experiment



Probability model

A bag of identical balls, except for their color (or a label). If the bag is well shaken, every ball is equally likely to be picked.


$$\Omega = \{\text{white, red, yellow, grey, purple, blue, maroon, green}\}$$

Probability Space: Formalism

Simplest physical model of a **uniform** probability space:



Physical experiment



Probability model

A bag of identical balls, except for their color (or a label). If the bag is well shaken, every ball is equally likely to be picked.

$$\Omega = \{\text{white, red, yellow, grey, purple, blue, maroon, green}\}$$


$$Pr[\text{blue}] =$$

Probability Space: Formalism

Simplest physical model of a **uniform** probability space:



Physical experiment



Probability model

A bag of identical balls, except for their color (or a label). If the bag is well shaken, every ball is equally likely to be picked.

$$\Omega = \{\text{white, red, yellow, grey, purple, blue, maroon, green}\}$$


$$Pr[\text{blue}] = \frac{1}{8}.$$

Probability Space: Formalism


Simplest physical model of a **non-uniform** probability space:

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:




Physical experiment




Probability model

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:



Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$


$$Pr[\text{Red}] =$$

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:



Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$


$$Pr[\text{Red}] = \frac{3}{10},$$

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:



Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$


$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] =$$

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

Probability Space: Formalism

Simplest physical model of a **non-uniform** probability space:



Physical experiment



Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$


Note: Probabilities are restricted to rational numbers: $\frac{N_k}{N}$.

Probability Space: Formalism


Physical model of a general **non-uniform** probability space:

Probability Space: Formalism

Physical model of a general **non-uniform** probability space:




Physical experiment




Probability model

Probability Space: Formalism

Physical model of a general **non-uniform** probability space:



Physical experiment




Probability model


The roulette wheel stops in sector ω with probability p_ω .

Probability Space: Formalism

Physical model of a general **non-uniform** probability space:



Physical experiment




Probability model

The roulette wheel stops in sector ω with probability p_ω .


$$\Omega = \{1, 2, 3, \dots, N\},$$

Probability Space: Formalism

Physical model of a general **non-uniform** probability space:



Physical experiment



Probability model

The roulette wheel stops in sector ω with probability p_ω .

$$\Omega = \{1, 2, 3, \dots, N\}, Pr[\omega] = p_\omega.$$

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .
- ▶ In this case, it would be wrong to think that $\Omega = \{H, T\}$ and that the experiment selects two outcomes.

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .
- ▶ In this case, it would be wrong to think that $\Omega = \{H, T\}$ and that the experiment selects two outcomes.
- ▶ Why?

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .
- ▶ In this case, it would be wrong to think that $\Omega = \{H, T\}$ and that the experiment selects two outcomes.
- ▶ Why? Because this would not describe how the two coin flips are related to each other.

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .
- ▶ In this case, it would be wrong to think that $\Omega = \{H, T\}$ and that the experiment selects two outcomes.
- ▶ Why? Because this would not describe how the two coin flips are related to each other.
- ▶ For instance, say we glue the coins side-by-side so that they face up the same way.

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .
- ▶ In this case, it would be wrong to think that $\Omega = \{H, T\}$ and that the experiment selects two outcomes.
- ▶ Why? Because this would not describe how the two coin flips are related to each other.
- ▶ For instance, say we glue the coins side-by-side so that they face up the same way. Then one gets HH or TT with probability 50% each.

An important remark

- ▶ The random experiment selects **one and only one** outcome in Ω .
- ▶ For instance, when we flip a fair coin **twice**
 - ▶ $\Omega = \{HH, TH, HT, TT\}$
 - ▶ The experiment selects *one* of the elements of Ω .
- ▶ In this case, it would be wrong to think that $\Omega = \{H, T\}$ and that the experiment selects two outcomes.
- ▶ Why? Because this would not describe how the two coin flips are related to each other.
- ▶ For instance, say we glue the coins side-by-side so that they face up the same way. Then one gets HH or TT with probability 50% each. This is not captured by 'picking two outcomes.'

Events

Next idea: an event!

Set notation review

Set notation review




Figure : Two events

Set notation review




Figure : Two events




Figure : Complement
(not)

Set notation review




Figure : Two events




Figure : Union (or)




Figure : Complement
(not)

Set notation review




Figure : Two events




Figure : Union (or)




Figure : Complement
(not)




Figure : Intersection
(and)

Set notation review




Figure : Two events




Figure : Union (or)




Figure : Difference (A , not B)




Figure : Complement (not)




Figure : Intersection (and)

Set notation review




Figure : Two events




Figure : Union (or)




Figure : Difference (A , not B)




Figure : Complement (not)




Figure : Intersection (and)




Figure : Symmetric difference (only one)

Probability of exactly one 'heads' in two coin flips?

Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT, TH .

Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT , TH .

This leads to a definition!

Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT, TH .

This leads to a definition!

Definition:

Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT, TH .

This leads to a definition!

Definition:

- ▶ An **event**, E , is a subset of outcomes: $E \subset \Omega$.

Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT, TH .

This leads to a definition!

Definition:

- ▶ An **event**, E , is a subset of outcomes: $E \subset \Omega$.
- ▶ The **probability of** E is defined as $Pr[E] = \sum_{\omega \in E} Pr[\omega]$.


Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT, TH .

This leads to a definition!

Definition:

- ▶ An **event**, E , is a subset of outcomes: $E \subset \Omega$.
- ▶ The **probability of** E is defined as $Pr[E] = \sum_{\omega \in E} Pr[\omega]$.




Probability of exactly one 'heads' in two coin flips?

Idea: Sum the probabilities of all the different outcomes that have exactly one 'heads': HT, TH .


This leads to a definition!

Definition:

- ▶ An **event**, E , is a subset of outcomes: $E \subset \Omega$.
- ▶ The **probability of** E is defined as $Pr[E] = \sum_{\omega \in E} Pr[\omega]$.




Uniform Probability Space




Event: Example

Event: Example




Physical experiment




Probability model

Event: Example




Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

Event: Example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

$$Pr[\text{Red}] =$$

Event: Example




Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10},$$

Event: Example




Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] =$$

Event: Example




Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

Event: Example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

$$E = \{\text{Red, Green}\}$$

Event: Example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

$$E = \{\text{Red, Green}\} \Rightarrow Pr[E] =$$

Event: Example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

$$E = \{\text{Red, Green}\} \Rightarrow Pr[E] = \frac{3+4}{10} =$$

Event: Example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

$$E = \{\text{Red, Green}\} \Rightarrow Pr[E] = \frac{3+4}{10} = \frac{3}{10} + \frac{4}{10} =$$

Event: Example



Physical experiment



Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$
$$Pr[\text{Red}] = \frac{3}{10}, Pr[\text{Green}] = \frac{4}{10}, \text{ etc.}$$

$$E = \{\text{Red, Green}\} \Rightarrow Pr[E] = \frac{3+4}{10} = \frac{3}{10} + \frac{4}{10} = Pr[\text{Red}] + Pr[\text{Green}].$$

Probability of exactly one heads in two coin flips?

Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Uniform probability space:

$$Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}.$$


Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Uniform probability space:

$$Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}.$$

Event, E , “exactly one heads”: $\{TH, HT\}$.




Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Uniform probability space:

$$Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}.$$

Event, E , “exactly one heads”: $\{TH, HT\}$.



$$Pr[E] = \sum_{\omega \in E} Pr[\omega]$$


Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Uniform probability space:

$$Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}.$$

Event, E , “exactly one heads”: $\{TH, HT\}$.



$$Pr[E] = \sum_{\omega \in E} Pr[\omega] = \frac{|E|}{|\Omega|}$$


Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Uniform probability space:

$$Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}.$$

Event, E , “exactly one heads”: $\{TH, HT\}$.



$$Pr[E] = \sum_{\omega \in E} Pr[\omega] = \frac{|E|}{|\Omega|} = \frac{2}{4}$$


Probability of exactly one heads in two coin flips?

Sample Space, $\Omega = \{HH, HT, TH, TT\}$.

Uniform probability space:

$$Pr[HH] = Pr[HT] = Pr[TH] = Pr[TT] = \frac{1}{4}.$$

Event, E , “exactly one heads”: $\{TH, HT\}$.



$$Pr[E] = \sum_{\omega \in E} Pr[\omega] = \frac{|E|}{|\Omega|} = \frac{2}{4} = \frac{1}{2}.$$

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20};$$

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

- What is more likely?

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

- $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or
- $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Example: 20 coin tosses.

20 coin tosses

Sample space: $\Omega =$ set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer:

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why?

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why? There are many sequences of 20 tosses with ten Hs;

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why? There are many sequences of 20 tosses with ten Hs;
only one with twenty Hs.

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why? There are many sequences of 20 tosses with ten Hs;

only one with twenty Hs. $\Rightarrow Pr[E_1] = \frac{1}{|\Omega|} \ll Pr[E_2] = \frac{|E_2|}{|\Omega|}$.

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why? There are many sequences of 20 tosses with ten Hs;

only one with twenty Hs. $\Rightarrow Pr[E_1] = \frac{1}{|\Omega|} \ll Pr[E_2] = \frac{|E_2|}{|\Omega|}$.

$$|E_2| =$$

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why? There are many sequences of 20 tosses with ten Hs;

only one with twenty Hs. $\Rightarrow Pr[E_1] = \frac{1}{|\Omega|} \ll Pr[E_2] = \frac{|E_2|}{|\Omega|}$.

$$|E_2| = \binom{20}{10} =$$

Example: 20 coin tosses.

20 coin tosses

Sample space: Ω = set of 20 fair coin tosses.

$$\Omega = \{T, H\}^{20} \equiv \{0, 1\}^{20}; \quad |\Omega| = 2^{20}.$$

► What is more likely?

► $\omega_1 := (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1)$, or

► $\omega_2 := (1, 0, 1, 1, 0, 0, 0, 1, 0, 1, 0, 1, 1, 0, 1, 1, 1, 0, 0, 0)$?

Answer: Both are equally likely: $Pr[\omega_1] = Pr[\omega_2] = \frac{1}{|\Omega|}$.

► What is more likely?

(E_1) Twenty Hs out of twenty, or

(E_2) Ten Hs out of twenty?

Answer: Ten Hs out of twenty.

Why? There are many sequences of 20 tosses with ten Hs;

only one with twenty Hs. $\Rightarrow Pr[E_1] = \frac{1}{|\Omega|} \ll Pr[E_2] = \frac{|E_2|}{|\Omega|}$.

$$|E_2| = \binom{20}{10} = 184,756.$$

Probability of n heads in 100 coin tosses.

Probability of n heads in 100 coin tosses.


$$\Omega = \{H, T\}^{100};$$

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$


Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Probability of n heads in 100 coin tosses.


$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Event $E_n = 'n \text{ heads}';$

Probability of n heads in 100 coin tosses.


$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Event $E_n = \text{'}n \text{ heads'}; |E_n| =$

Probability of n heads in 100 coin tosses.


$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



$$\text{Event } E_n = \text{'n heads'}; |E_n| = \binom{100}{n}$$

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$




Event $E_n = \text{'n heads'}; |E_n| = \binom{100}{n}$

$$p_n := \Pr[E_n] =$$

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$




Event $E_n = \text{'n heads'}; |E_n| = \binom{100}{n}$

$$p_n := Pr[E_n] = \frac{|E_n|}{|\Omega|} =$$

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$




Event $E_n = \text{'n heads'}; |E_n| = \binom{100}{n}$

$$p_n := Pr[E_n] = \frac{|E_n|}{|\Omega|} = \frac{\binom{100}{n}}{2^{100}}$$

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$




Event $E_n = \text{'}n \text{ heads'}; |E_n| = \binom{100}{n}$

$$p_n := \Pr[E_n] = \frac{|E_n|}{|\Omega|} = \frac{\binom{100}{n}}{2^{100}}$$

Observe:

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Event $E_n = 'n \text{ heads}'; |E_n| = \binom{100}{n}$


$$p_n := Pr[E_n] = \frac{|E_n|}{|\Omega|} = \frac{\binom{100}{n}}{2^{100}}$$

Observe:

- Concentration around mean:

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Event $E_n = 'n \text{ heads}'; |E_n| = \binom{100}{n}$


$$p_n := Pr[E_n] = \frac{|E_n|}{|\Omega|} = \frac{\binom{100}{n}}{2^{100}}$$

Observe:

- Concentration around mean:
Law of Large Numbers;

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Event $E_n = 'n \text{ heads}'; |E_n| = \binom{100}{n}$


$$p_n := Pr[E_n] = \frac{|E_n|}{|\Omega|} = \frac{\binom{100}{n}}{2^{100}}$$

Observe:

- Concentration around mean:
Law of Large Numbers;
- Bell-shape:

Probability of n heads in 100 coin tosses.

$$\Omega = \{H, T\}^{100}; |\Omega| = 2^{100}.$$



Event $E_n = 'n \text{ heads}'; |E_n| = \binom{100}{n}$


$$p_n := Pr[E_n] = \frac{|E_n|}{|\Omega|} = \frac{\binom{100}{n}}{2^{100}}$$

Observe:

- Concentration around mean:
Law of Large Numbers;
- Bell-shape: Central Limit Theorem.

Roll a red and a blue die.

Roll a red and a blue die.



$$Pr[\text{Sum to 7}] = \frac{6}{36}$$

$$Pr[\text{Sum to 10}] = \frac{3}{36}$$

Exactly 50 heads in 100 coin tosses.

Sample space: Ω = set of 100 coin tosses

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.

$$|\Omega| = 2 \times 2 \times \cdots \times 2$$

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.
 $|\Omega| = 2 \times 2 \times \cdots \times 2 = 2^{100}$.

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.
 $|\Omega| = 2 \times 2 \times \cdots \times 2 = 2^{100}$.

Uniform probability space: $Pr[\omega] = \frac{1}{2^{100}}$.

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.
 $|\Omega| = 2 \times 2 \times \cdots \times 2 = 2^{100}$.

Uniform probability space: $Pr[\omega] = \frac{1}{2^{100}}$.

Event $E = \text{"100 coin tosses with exactly 50 heads"}$

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.
 $|\Omega| = 2 \times 2 \times \cdots \times 2 = 2^{100}$.

Uniform probability space: $Pr[\omega] = \frac{1}{2^{100}}$.

Event $E = \text{"100 coin tosses with exactly 50 heads"}$

$|E|?$

Choose 50 positions out of 100 to be heads.

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.
 $|\Omega| = 2 \times 2 \times \cdots \times 2 = 2^{100}$.

Uniform probability space: $Pr[\omega] = \frac{1}{2^{100}}$.

Event $E = \text{"100 coin tosses with exactly 50 heads"}$

$|E|$?

Choose 50 positions out of 100 to be heads.

$$|E| = \binom{100}{50}.$$

Exactly 50 heads in 100 coin tosses.

Sample space: $\Omega = \text{set of 100 coin tosses} = \{H, T\}^{100}$.
 $|\Omega| = 2 \times 2 \times \cdots \times 2 = 2^{100}$.

Uniform probability space: $Pr[\omega] = \frac{1}{2^{100}}$.

Event $E = \text{"100 coin tosses with exactly 50 heads"}$

$|E|?$

Choose 50 positions out of 100 to be heads.

$$|E| = \binom{100}{50}.$$

$$Pr[E] = \frac{\binom{100}{50}}{2^{100}}.$$

Calculation.

Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

Calculation.

Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

$$\binom{2n}{n} \approx \frac{\sqrt{4\pi n}(2n/e)^{2n}}{[\sqrt{2\pi n}(n/e)^n]^2}$$

Calculation.

Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

$$\binom{2n}{n} \approx \frac{\sqrt{4\pi n}(2n/e)^{2n}}{[\sqrt{2\pi n}(n/e)^n]^2} \approx \frac{4^n}{\sqrt{\pi n}}.$$

Calculation.

Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

$$\binom{2n}{n} \approx \frac{\sqrt{4\pi n}(2n/e)^{2n}}{[\sqrt{2\pi n}(n/e)^n]^2} \approx \frac{4^n}{\sqrt{\pi n}}.$$

$$Pr[E] = \frac{|E|}{|\Omega|} =$$

Calculation.

Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

$$\binom{2n}{n} \approx \frac{\sqrt{4\pi n}(2n/e)^{2n}}{[\sqrt{2\pi n}(n/e)^n]^2} \approx \frac{4^n}{\sqrt{\pi n}}.$$

$$Pr[E] = \frac{|E|}{|\Omega|} = \frac{|E|}{2^{2n}} =$$

Calculation.

Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

$$\binom{2n}{n} \approx \frac{\sqrt{4\pi n}(2n/e)^{2n}}{[\sqrt{2\pi n}(n/e)^n]^2} \approx \frac{4^n}{\sqrt{\pi n}}.$$

$$Pr[E] = \frac{|E|}{|\Omega|} = \frac{|E|}{2^{2n}} = \frac{1}{\sqrt{\pi n}} =$$

Calculation.


Stirling formula (for large n):

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n.$$

$$\binom{2n}{n} \approx \frac{\sqrt{4\pi n}(2n/e)^{2n}}{[\sqrt{2\pi n}(n/e)^n]^2} \approx \frac{4^n}{\sqrt{\pi n}}.$$

$$Pr[E] = \frac{|E|}{|\Omega|} = \frac{|E|}{2^{2n}} = \frac{1}{\sqrt{\pi n}} = \frac{1}{\sqrt{50\pi}} \approx .08.$$

Exactly 50 heads in 100 coin tosses.



Lecture 13: Summary

1. Random Experiment
2. Probability Space: Ω ; $Pr[\omega] \in [0, 1]$; $\sum_{\omega} Pr[\omega] = 1$.
3. Uniform Probability Space: $Pr[\omega] = 1/|\Omega|$ for all $\omega \in \Omega$.
4. Events: subsets of Ω .

$$Pr[E] = \sum_{\omega \in E} Pr[\omega].$$