

CS70: Discrete Math and Probability

Slides adopted from Satish Rao, CS70 Spring 2016

June 20, 2016

Programming Computers

Programming Computers \equiv Superpower!

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Discrete Math

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Discrete Math \implies immense application.

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Discrete Math \implies immense application.

Computers learn and interact with the world?

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Discrete Math \implies immense application.

Computers learn and interact with the world?

E.g. machine learning, data analysis.

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Discrete Math \implies immense application.

Computers learn and interact with the world?

E.g. machine learning, data analysis.

Probability!

Programming Computers \equiv Superpower!

What are your super powerful programs doing?

Logic and Proofs!

Induction \equiv Recursion.

What can computers do?

Work with discrete objects.

Discrete Math \implies immense application.

Computers learn and interact with the world?

E.g. machine learning, data analysis.

Probability!

See note 1, for more discussion.

Course Webpage: www.eecs70.org

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Questions

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Questions \implies piazza:

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Questions \implies piazza:

piazza.com/berkeley/summer2016/cs70

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Questions \implies piazza:

piazza.com/berkeley/summer2016/cs70

Assessment:

Homework: 20%

Midterm 1 (07/08): 20%

Midterm 2 (07/29): 20%

Final (08/12): 35%

Quiz: 4%

Sundry: 1%

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Questions \implies piazza:

piazza.com/berkeley/summer2016/cs70

Assessment:

Homework: 20%

Midterm 1 (07/08): 20%

Midterm 2 (07/29): 20%

Final (08/12): 35%

Quiz: 4%

Sundry: 1%

Conflicts?

Course Webpage: www.eecs70.org

Explains policies, has homework/discussion worksheets, slides, exam dates, etc.

Questions \implies piazza:

piazza.com/berkeley/summer2016/cs70

Assessment:

Homework: 20%

Midterm 1 (07/08): 20%

Midterm 2 (07/29): 20%

Final (08/12): 35%

Quiz: 4%

Sundry: 1%

Conflicts? Piazza pinned post.

Take homework seriously!

Learning tips

Take homework seriously!
Go to homework parties,

Learning tips

Take homework seriously!

Go to homework parties, study groups

Learning tips

Take homework seriously!

Go to homework parties, study groups

VERY fast paced, start early

Learning tips

Take homework seriously!

Go to homework parties, study groups

VERY fast paced, start early

Use piazza, help each other out

Learning tips

Take homework seriously!

Go to homework parties, study groups

VERY fast paced, start early

Use piazza, help each other out

Questions?

3 Co-Instructors

Just graduated,

Just graduated, from Berkeley

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Enjoy climbing, badminton, boxing,

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Enjoy climbing, badminton, boxing, also like to watch movies and games

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Enjoy climbing, badminton, boxing, also like to watch movies and games
Recently I'm climbing ...

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Enjoy climbing, badminton, boxing, also like to watch movies and games
Recently I'm climbing ... the ladder of league of legends ranking system ...

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Enjoy climbing, badminton, boxing, also like to watch movies and games
Recently I'm climbing ... the ladder of league of legends ranking system ...

Office hours: Monday 10-11, Tuesday 11-12 in Soda 611

Just graduated, from Berkeley

Been TA for CS70 for two semesters

Will start working at Google as a software engineer on September

Enjoy climbing, badminton, boxing, also like to watch movies and games
Recently I'm climbing ... the ladder of league of legends ranking system ...

Office hours: Monday 10-11, Tuesday 11-12 in Soda 611 or by appointment

Email: dinh@cs.berkeley.edu

Office Hours: M/W 3:30-5:00 (right after lecture) in 606 Soda

I just finished my first year of grad school. My research interests are numerical algorithms and complexity theory - essentially, I work on making faster algorithms for doing things like solving equations, factoring matrices, etc. (and proving that they run fast!), as well as showing that there are limits on how fast we can make these algorithms.

Also did my undergrad here at Cal - CS70 was by far my favorite lower-div.

Fun fact: I like to make ice cream.

Not here today.

Not here today. Tomorrow lecture

3 Co-Instructors

12 awesome and talented TAs.

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side,
and **mode of travel**.

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:
"If a person travels to Chicago, he/she flies."

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:
"If a person travels to Chicago, he/she flies."

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:
"If a person travels to Chicago, he/she flies."
- Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Alice	Bob	Charlie	Donna
Baltimore	drove	Chicago	flew

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:
"If a person travels to Chicago, he/she flies."
- Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Alice	Bob	Charlie	Donna
Baltimore	drove	Chicago	flew

- Which cards do you need to flip to test the theory?

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:
"If a person travels to Chicago, he/she flies."
- Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Alice	Bob	Charlie	Donna
Baltimore	drove	Chicago	flew

- Which cards do you need to flip to test the theory?

Answer:

Wason's experiment:1

Suppose we have four cards on a table:

- 1st about Alice, 2nd about Bob, 3rd Charlie, 4th Donna.
- Card contains person's **destination** on one side, and **mode of travel**.
- Consider the theory:
"If a person travels to Chicago, he/she flies."
- Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Alice	Bob	Charlie	Donna
Baltimore	drove	Chicago	flew

- Which cards do you need to flip to test the theory?

Answer: Later.

Today: Note 1.

Today: Note 1. Note 0 is background.

Today: Note 1. Note 0 is background. Do read/skim it.

Today: Note 1. Note 0 is background. Do read/skim it.

The language of proofs!

Today: Note 1. Note 0 is background. Do read/skim it.

The language of proofs!

1. Propositions.
2. Propositional Forms.
3. Implication.
4. Truth Tables
5. Quantifiers
6. More De Morgan's Laws

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$$2+2 = 4$$

$$2+2 = 3$$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$$4 + 5$$

$$x + x$$

Alice travelled to Chicago

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$$2+2 = 4$$

$$2+2 = 3$$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$$4 + 5$$

$$x + x$$

Alice travelled to Chicago

Proposition

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$$2+2 = 4$$

$$2+2 = 3$$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$$4 + 5$$

$$x + x$$

Alice travelled to Chicago

Proposition

True

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

Proposition

True

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition True

Proposition True

Proposition False

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition True

Proposition True

Proposition False

Proposition False

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

False

Not a Proposition

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

False

Not a Proposition

Proposition

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition **True**

Proposition **True**

Proposition **False**

Proposition **False**

Not a Proposition

Proposition **False**

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

False

Not a Proposition

Proposition

False

Not a Proposition.

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

False

Not a Proposition

Proposition

False

Not a Proposition.

Not a Proposition.

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

False

Not a Proposition

Proposition

False

Not a Proposition.

Not a Proposition.

Proposition.

Again: “value” of a proposition is ...

Propositions: Statements that are true or false.

$\sqrt{2}$ is irrational

$2+2 = 4$

$2+2 = 3$

826th digit of pi is 4

Stephen Curry is a good player.

All evens > 2 are sums of 2 primes

$4 + 5$

$x + x$

Alice travelled to Chicago

Proposition

True

Proposition

True

Proposition

False

Proposition

False

Not a Proposition

Proposition

False

Not a Proposition.

Not a Proposition.

Proposition.

False

Again: “value” of a proposition is ... True or False

Propositional Forms.

Put propositions together to make another...

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** .

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is True when at least one P or Q is True .

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is True when at least one P or Q is True . Else False .

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is True when at least one P or Q is True . Else False .

Negation (“not”): $\neg P$

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is True when at least one P or Q is True . Else False .

Negation (“not”): $\neg P$

“ $\neg P$ ” is True when P is False .

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is True when at least one P or Q is True . Else False .

Negation (“not”): $\neg P$

“ $\neg P$ ” is True when P is False . Else False .

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is True when both P and Q are True . Else False .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is True when at least one P or Q is True . Else False .

Negation (“not”): $\neg P$

“ $\neg P$ ” is True when P is False . Else False .

Examples:

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** . Else **False** .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is **True** when at least one P or Q is **True** . Else **False** .

Negation (“not”): $\neg P$

“ $\neg P$ ” is **True** when P is **False** . Else **False** .

Examples:

\neg “ $(2 + 2 = 4)$ ” – a proposition that is ...

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** . Else **False** .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is **True** when at least one P or Q is **True** . Else **False** .

Negation (“not”): $\neg P$

“ $\neg P$ ” is **True** when P is **False** . Else **False** .

Examples:

\neg “ $(2 + 2 = 4)$ ” – a proposition that is ... **False**

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** . Else **False** .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is **True** when at least one P or Q is **True** . Else **False** .

Negation (“not”): $\neg P$

“ $\neg P$ ” is **True** when P is **False** . Else **False** .

Examples:

\neg “ $(2 + 2 = 4)$ ” – a proposition that is ... **False**

“ $2 + 2 = 3$ ” \wedge “ $2 + 2 = 4$ ” – a proposition that is ...

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** . Else **False** .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is **True** when at least one P or Q is **True** . Else **False** .

Negation (“not”): $\neg P$

“ $\neg P$ ” is **True** when P is **False** . Else **False** .

Examples:

\neg “ $(2 + 2 = 4)$ ” – a proposition that is ... **False**

“ $2 + 2 = 3$ ” \wedge “ $2 + 2 = 4$ ” – a proposition that is ... **False**

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** . Else **False** .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is **True** when at least one P or Q is **True** . Else **False** .

Negation (“not”): $\neg P$

“ $\neg P$ ” is **True** when P is **False** . Else **False** .

Examples:

\neg “ $2 + 2 = 4$ ” – a proposition that is ... **False**

“ $2 + 2 = 3$ ” \wedge “ $2 + 2 = 4$ ” – a proposition that is ... **False**

“ $2 + 2 = 3$ ” \vee “ $2 + 2 = 4$ ” – a proposition that is ...

Propositional Forms.

Put propositions together to make another...

Conjunction (“and”): $P \wedge Q$

“ $P \wedge Q$ ” is **True** when both P and Q are **True** . Else **False** .

Disjunction (“or”): $P \vee Q$

“ $P \vee Q$ ” is **True** when at least one P or Q is **True** . Else **False** .

Negation (“not”): $\neg P$

“ $\neg P$ ” is **True** when P is **False** . Else **False** .

Examples:

\neg “ $2 + 2 = 4$ ” – a proposition that is ... **False**

“ $2 + 2 = 3$ ” \wedge “ $2 + 2 = 4$ ” – a proposition that is ... **False**

“ $2 + 2 = 3$ ” \vee “ $2 + 2 = 4$ ” – a proposition that is ... **True**

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

$P \wedge Q$...

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

$P \wedge Q \dots$ False

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

$P \wedge Q$... False

$P \vee Q$...

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

$P \wedge Q \dots$ False

$P \vee Q \dots$ True

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

$P \wedge Q$... False

$P \vee Q$... True

$\neg P$...

Propositional Forms: quick check!

$P = \text{"}\sqrt{2} \text{ is rational"}$

$Q = \text{"826th digit of pi is 2"}$

P is ...False .

Q is ...True .

$P \wedge Q$... False

$P \vee Q$... True

$\neg P$... True

Put them together..

Propositions:

C_1 - Take class 1

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Can you take class 1?

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Can you take class 1?

Can you take class 1 and class 5 together?

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Can you take class 1?

Can you take class 1 and class 5 together?

This seems ...

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Can you take class 1?

Can you take class 1 and class 5 together?

This seems ...**complicated**.

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Can you take class 1?

Can you take class 1 and class 5 together?

This seems ...**complicated**.

We can program!!!!

Put them together..

Propositions:

C_1 - Take class 1

C_2 - Take class 2

....

You can only take one of class 1, class 2 and one of class 3, class 4, or take both class 2, class 3 and take either class 5 or class 6, as long as you are not taking class 4 at the same time.

Propositional Form:

$$((C_1 \vee C_2) \wedge (C_3 \vee C_4)) \vee ((C_2 \wedge C_3) \wedge (C_5 \vee C_6) \wedge (\neg C_4))$$

Can you take class 1?

Can you take class 1 and class 5 together?

This seems ...**complicated**.

We can program!!!! We need a way to keep track!

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	
F	T	
F	F	

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	
F	F	

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	
T	F	
F	T	
F	F	

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	
F	T	
F	F	

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	
T	F		
F	T		
F	F		

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F		
F	T		
F	F		

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	
F	T		
F	F		

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T		
F	F		

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

DeMorgan's Law's for Negation: distribute and flip!

$\neg(P \wedge Q)$

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

DeMorgan's Law's for Negation: distribute and flip!

$$\neg(P \wedge Q) \equiv \neg P \vee \neg Q$$

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

DeMorgan's Law's for Negation: distribute and flip!

$$\neg(P \wedge Q) \equiv \neg P \vee \neg Q$$

$$\neg(P \vee Q)$$

Truth Tables for Propositional Forms.

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Notice: \wedge and \vee are commutative.

One use for truth tables: Logical Equivalence of propositional forms!

Example: $\neg(P \wedge Q)$ logically equivalent to $\neg P \vee \neg Q$

...because the two propositional forms have the same...

....Truth Table!

P	Q	$\neg(P \vee Q)$	$\neg P \wedge \neg Q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

DeMorgan's Law's for Negation: distribute and flip!

$$\neg(P \wedge Q) \equiv \neg P \vee \neg Q$$

$$\neg(P \vee Q) \equiv \neg P \wedge \neg Q$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$,

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

LHS: $T \wedge (Q \vee R)$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is True .

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R)$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R)$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R)$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F)$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Simplify: $T \vee Q \equiv T$,

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Simplify: $T \vee Q \equiv T$, $F \vee Q \equiv Q$.

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Simplify: $T \vee Q \equiv T$, $F \vee Q \equiv Q$.

Foil 1:

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Simplify: $T \vee Q \equiv T$, $F \vee Q \equiv Q$.

Foil 1:

$$(A \vee B) \wedge (C \vee D) \equiv (A \wedge C) \vee (A \wedge D) \vee (B \wedge C) \vee (B \wedge D)?$$

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Simplify: $T \vee Q \equiv T$, $F \vee Q \equiv Q$.

Foil 1:

$$(A \vee B) \wedge (C \vee D) \equiv (A \wedge C) \vee (A \wedge D) \vee (B \wedge C) \vee (B \wedge D)?$$

Foil 2:

Distributive?

$$P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)?$$

Simplify: $(T \wedge Q) \equiv Q$, $(F \wedge Q) \equiv F$.

Cases:

P is **True**.

$$\text{LHS: } T \wedge (Q \vee R) \equiv (Q \vee R).$$

$$\text{RHS: } (T \wedge Q) \vee (T \wedge R) \equiv (Q \vee R).$$

P is **False**.

$$\text{LHS: } F \wedge (Q \vee R) \equiv F.$$

$$\text{RHS: } (F \wedge Q) \vee (F \wedge R) \equiv (F \vee F) \equiv F.$$

$$P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)?$$

Simplify: $T \vee Q \equiv T$, $F \vee Q \equiv Q$.

Foil 1:

$$(A \vee B) \wedge (C \vee D) \equiv (A \wedge C) \vee (A \wedge D) \vee (B \wedge C) \vee (B \wedge D)?$$

Foil 2:

$$(A \wedge B) \vee (C \wedge D) \equiv (A \vee C) \wedge (A \vee D) \wedge (B \vee C) \wedge (B \vee D)?$$

Implication.

$P \implies Q$ interpreted as

Implication.

$P \implies Q$ interpreted as

If P , then Q .

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Q = "you will get wet"

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Q = "you will get wet"

Statement: "Stand in the rain"

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Q = "you will get wet"

Statement: "Stand in the rain"

Can conclude: "you'll get wet."

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Q = "you will get wet"

Statement: "Stand in the rain"

Can conclude: "you'll get wet."

Statement: If a right triangle has sidelengths $a \leq b \leq c$, then $a^2 + b^2 = c^2$.

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Q = "you will get wet"

Statement: "Stand in the rain"

Can conclude: "you'll get wet."

Statement: If a right triangle has sidelengths $a \leq b \leq c$, then $a^2 + b^2 = c^2$.

P = "a right triangle has sidelengths $a \leq b \leq c$ ",

Implication.

$P \implies Q$ interpreted as

If P , then Q .

True Statements: $P, P \implies Q$.

Conclude: Q is true.

Examples:

Statement: If you stand in the rain, then you'll get wet.

P = "you stand in the rain"

Q = "you will get wet"

Statement: "Stand in the rain"

Can conclude: "you'll get wet."

Statement: If a right triangle has sidelengths $a \leq b \leq c$, then $a^2 + b^2 = c^2$.

P = "a right triangle has sidelengths $a \leq b \leq c$ ",

Q = " $a^2 + b^2 = c^2$ ".

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True**

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

$P \implies Q$ and P are **True** *does* mean Q is **True** .

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

$P \implies Q$ and P are **True** *does* mean Q is **True** .

The chemical plant pollutes river.

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

$P \implies Q$ and P are **True** *does* mean Q is **True** .

The chemical plant pollutes river. Can we conclude fish die?

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

$P \implies Q$ and P are **True** *does* mean Q is **True** .

The chemical plant pollutes river. Can we conclude fish die?

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

$P \implies Q$ and P are **True** *does* mean Q is **True** .

The chemical plant pollutes river. Can we conclude fish die?

Some Fun: use propositional formulas to describe implication?

Non-Consequences/consequences of Implication

The statement " $P \implies Q$ "

only is **False** if P is **True** and Q is **False** .

False implies nothing

P **False** means Q can be **True** or **False**

Anything implies true.

P can be **True** or **False** when Q is **True**

If chemical plant pollutes river, fish die.

If fish die, did chemical plant pollute river?

Not necessarily.

$P \implies Q$ and Q are **True** does not mean P is **True**

Be careful!

Instead we have:

$P \implies Q$ and P are **True** *does* mean Q is **True** .

The chemical plant pollutes river. Can we conclude fish die?

Some Fun: use propositional formulas to describe implication?

$((P \implies Q) \wedge P) \implies Q$.

Implication and English.

$$P \implies Q$$

- If P , then Q .

Implication and English.

$$P \implies Q$$

- If P , then Q .
- Q if P .

Just reversing the order.

Implication and English.

$$P \implies Q$$

- If P , then Q .
- Q if P .

Just reversing the order.

$$P \implies Q$$

- If P , then Q .
- Q if P .

Just reversing the order.

- P only if Q .

Remember if P is true then Q must be true.

this suggests that P can only be true if Q is true.

since if Q is false P must have been false.

$$P \implies Q$$

- If P , then Q .

- Q if P .

Just reversing the order.

- P only if Q .

Remember if P is true then Q must be true.

this suggests that P can only be true if Q is true.

since if Q is false P must have been false.

- P is sufficient for Q .

This means that proving P allows you
to conclude that Q is true.

$$P \implies Q$$

- If P , then Q .

- Q if P .

Just reversing the order.

- P only if Q .

Remember if P is true then Q must be true.

this suggests that P can only be true if Q is true.

since if Q is false P must have been false.

- P is sufficient for Q .

This means that proving P allows you

to conclude that Q is true.

- Q is necessary for P .

For P to be true it is necessary that Q is true.

Or if Q is false then we know that P is false.

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	
F	T	
F	F	

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	
F	F	

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	
T	F	
F	T	
F	F	

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	T
T	F	
F	T	
F	F	

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	T
T	F	F
F	T	T
F	F	T

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	T
T	F	F
F	T	T
F	F	T

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	T
T	F	F
F	T	T
F	F	T

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	T
T	F	F
F	T	T
F	F	T

$$\neg P \vee Q \equiv P \implies Q.$$

Truth Table: implication.

P	Q	$P \implies Q$
T	T	T
T	F	F
F	T	T
F	F	T

P	Q	$\neg P \vee Q$
T	T	T
T	F	F
F	T	T
F	F	T

$$\neg P \vee Q \equiv P \implies Q.$$

These two propositional forms are logically equivalent!

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
- If you stand in the rain, you get wet.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
 - If you stand in the rain, you get wet.
 - If you did not stand in the rain, you did not get wet.
(not contrapositive!)

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
 - If you stand in the rain, you get wet.
 - If you did not stand in the rain, you did not get wet.
(not contrapositive!)
 - If you did not get wet, you did not stand in the rain.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
 - If you stand in the rain, you get wet.
 - If you did not stand in the rain, you did not get wet.
(not contrapositive!)
 - If you did not get wet, you did not stand in the rain.
(contrapositive.)

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
 - If you stand in the rain, you get wet.
 - If you did not stand in the rain, you did not get wet.
(not contrapositive!)
 - If you did not get wet, you did not stand in the rain.
(contrapositive.)

Logically equivalent! Notation: \equiv .

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.
 - If the plant pollutes, fish die.
 - If the fish don't die, the plant does not pollute.
(contrapositive)
 - If you stand in the rain, you get wet.
 - If you did not stand in the rain, you did not get wet.
(not contrapositive!)
 - If you did not get wet, you did not stand in the rain.
(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q$$

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.
(contrapositive)
- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.
(not contrapositive!)
- If you did not get wet, you did not stand in the rain.
(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q$$

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!)

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P$$

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!)

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!)

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

- Converse of $P \implies Q$ is $Q \implies P$.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!)

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

- Converse of $P \implies Q$ is $Q \implies P$.

If fish die the plant pollutes.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!) converse!

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

- Converse of $P \implies Q$ is $Q \implies P$.

If fish die the plant pollutes.

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!) converse!

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

- Converse of $P \implies Q$ is $Q \implies P$.

If fish die the plant pollutes.

Not logically equivalent!

Contrapositive, Converse

- Contrapositive of $P \implies Q$ is $\neg Q \implies \neg P$.

- If the plant pollutes, fish die.
- If the fish don't die, the plant does not pollute.

(contrapositive)

- If you stand in the rain, you get wet.
- If you did not stand in the rain, you did not get wet.

(not contrapositive!) converse!

- If you did not get wet, you did not stand in the rain.

(contrapositive.)

Logically equivalent! Notation: \equiv .

$$P \implies Q \equiv \neg P \vee Q \equiv \neg(\neg Q) \vee \neg P \equiv \neg Q \implies \neg P.$$

- Converse of $P \implies Q$ is $Q \implies P$.

If fish die the plant pollutes.

Not logically equivalent!

- **Definition:** If $P \implies Q$ and $Q \implies P$ is P if and only if Q or $P \iff Q$.

(Logically Equivalent: \iff .)

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}.$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}.$
- $x > 2$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = \text{"}x \text{ is even"}$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = \text{"}x \text{ is even"}$

Same as boolean valued functions from 61A or 61AS!

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$
- Remember Wason's experiment!
 $F(x) = "Person x \text{ flew}."$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$
- Remember Wason's experiment!
 $F(x) = "Person x \text{ flew}."$
 $C(x) = "Person x \text{ went to Chicago}"$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$
- Remember Wason's experiment!
 $F(x) = "Person x \text{ flew}."$
 $C(x) = "Person x \text{ went to Chicago}"$
- $C(x) \implies F(x).$

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$
- Remember Wason's experiment!
 $F(x) = "Person x \text{ flew}."$
 $C(x) = "Person x \text{ went to Chicago}"$
- $C(x) \implies F(x)$. Theory from Wason's.

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$
- Remember Wason's experiment!
 $F(x) = "Person x \text{ flew}."$
 $C(x) = "Person x \text{ went to Chicago}"$
- $C(x) \implies F(x)$. Theory from Wason's.
If person x goes to Chicago then person x flew.

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = "x \text{ is even}"$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = "\sum_{i=1}^n i = \frac{n(n+1)}{2}."$
- $R(x) = "x > 2"$
- $G(n) = "n \text{ is even and the sum of two primes}"$
- Remember Wason's experiment!
 $F(x) = "Person x \text{ flew}."$
 $C(x) = "Person x \text{ went to Chicago}"$
- $C(x) \implies F(x)$. Theory from Wason's.
If person x goes to Chicago then person x flew.

Next:

Variables.

Propositions?

- $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.
- $x > 2$
- n is even and the sum of two primes

No. They have a free variable.

We call them predicates, e.g., $Q(x) = \text{"}x \text{ is even"}$

Same as boolean valued functions from 61A or 61AS!

- $P(n) = \text{"}\sum_{i=1}^n i = \frac{n(n+1)}{2}\text{"}$.
- $R(x) = \text{"}x > 2\text{"}$
- $G(n) = \text{"}n \text{ is even and the sum of two primes"}$
- Remember Wason's experiment!
 $F(x) = \text{"Person } x \text{ flew."}$
 $C(x) = \text{"Person } x \text{ went to Chicago"}$
- $C(x) \implies F(x)$. Theory from Wason's.
If person x goes to Chicago then person x flew.

Next: Statements about boolean valued functions!!

Quantifiers..

There exists quantifier:

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0)$ "

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1)$ "

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4)$ "

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S) (P(x))$. means "For all x in S , we have $P(x)$ is **True**."

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S) (P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S) (P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

"Adding 1 makes a bigger number."

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S)(P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

"Adding 1 makes a bigger number." $(\forall x \in \mathbb{N})(x + 1 > x)$

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S) (P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

"Adding 1 makes a bigger number." $(\forall x \in \mathbb{N}) (x + 1 > x)$

"the square of a number is always non-negative"

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S)(P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

"Adding 1 makes a bigger number." $(\forall x \in \mathbb{N})(x + 1 > x)$

"the square of a number is always non-negative" $(\forall x \in \mathbb{N})(x^2 \geq 0)$

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S)(P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

"Adding 1 makes a bigger number." $(\forall x \in \mathbb{N})(x + 1 > x)$

"the square of a number is always non-negative" $(\forall x \in \mathbb{N})(x^2 \geq 0)$

Wait!

There exists quantifier:

$(\exists x \in S)(P(x))$ means "There exists an x in S where $P(x)$ is true."

For example:

$$(\exists x \in \mathbb{N})(x = x^2)$$

Equivalent to " $(0 = 0) \vee (1 = 1) \vee (2 = 4) \vee \dots$ "

Much shorter to use a quantifier!

For all quantifier;

$(\forall x \in S) (P(x))$. means "For all x in S , we have $P(x)$ is **True**."

Examples:

"Adding 1 makes a bigger number." $(\forall x \in \mathbb{N}) (x + 1 > x)$

"the square of a number is always non-negative" $(\forall x \in \mathbb{N})(x^2 \geq 0)$

Wait! What is \mathbb{N} ?

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**:

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Quantifiers: universes.

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

Quantifiers: universes.

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

- $\mathbb{N} = \{0, 1, \dots\}$ (natural numbers).

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

- $\mathbb{N} = \{0, 1, \dots\}$ (natural numbers).
- $\mathbb{Z} = \{\dots, -1, 0, \dots\}$ (integers)

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

- $\mathbb{N} = \{0, 1, \dots\}$ (natural numbers).
- $\mathbb{Z} = \{\dots, -1, 0, \dots\}$ (integers)
- \mathbb{Z}^+ (positive integers)

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

- $\mathbb{N} = \{0, 1, \dots\}$ (natural numbers).
- $\mathbb{Z} = \{\dots, -1, 0, \dots\}$ (integers)
- \mathbb{Z}^+ (positive integers)
- \mathbb{R} (real numbers)

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

- $\mathbb{N} = \{0, 1, \dots\}$ (natural numbers).
- $\mathbb{Z} = \{\dots, -1, 0, \dots\}$ (integers)
- \mathbb{Z}^+ (positive integers)
- \mathbb{R} (real numbers)
- Any set: $S = \{Alice, Bob, Charlie, Donna\}$.

Proposition: “For all natural numbers n , $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.”

Proposition has **universe**: “the natural numbers”.

Universe examples include..

- $\mathbb{N} = \{0, 1, \dots\}$ (natural numbers).
- $\mathbb{Z} = \{\dots, -1, 0, \dots\}$ (integers)
- \mathbb{Z}^+ (positive integers)
- \mathbb{R} (real numbers)
- Any set: $S = \{Alice, Bob, Charlie, Donna\}$.
- See note 0 for more!

Back to: Wason's experiment:1

Theory:

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago."

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x)$

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** .

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** .

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B)$

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(Bob)$ must be **False** .

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(Bob)$ must be **False** .

$P(C)$ = **True** .

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(Bob)$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(\text{Bob})$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(\text{Bob})$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes. $P(C) \implies Q(C)$ means $Q(C)$ must be true.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(\text{Bob})$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes. $P(C) \implies Q(C)$ means $Q(C)$ must be true.

$Q(D)$ = **True** .

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(Bob)$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes. $P(C) \implies Q(C)$ means $Q(C)$ must be true.

$Q(D)$ = **True** . Do we care about $P(D)$?

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(\text{Bob})$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes. $P(C) \implies Q(C)$ means $Q(C)$ must be true.

$Q(D)$ = **True** . Do we care about $P(D)$?

No.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(Bob)$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes. $P(C) \implies Q(C)$ means $Q(C)$ must be true.

$Q(D)$ = **True** . Do we care about $P(D)$?

No. $P(D) \implies Q(D)$ holds whatever $P(D)$ is when $Q(D)$ is true.

Back to: Wason's experiment:1

Theory:

"If a person travels to Chicago, he/she flies."

Suppose you see that Alice went to Baltimore, Bob drove, Charlie went to Chicago, and Donna flew.

Which cards do you need to flip to test the theory?

$P(x)$ = "Person x went to Chicago." $Q(x)$ = "Person x flew"

Statement/theory: $\forall x \in \{A, B, C, D\}, P(x) \implies Q(x)$

$P(A)$ = **False** . Do we care about $Q(A)$?

No. $P(A) \implies Q(A)$, when $P(A)$ is **False** , $Q(A)$ can be anything.

$Q(B)$ = **False** . Do we care about $P(B)$?

Yes. $P(B) \implies Q(B) \equiv \neg Q(B) \implies \neg P(B)$.

So $P(Bob)$ must be **False** .

$P(C)$ = **True** . Do we care about $P(C)$?

Yes. $P(C) \implies Q(C)$ means $Q(C)$ must be true.

$Q(D)$ = **True** . Do we care about $P(D)$?

No. $P(D) \implies Q(D)$ holds whatever $P(D)$ is when $Q(D)$ is true.

Only have to turn over cards for Bob and Charlie.

More for all quantifiers examples.

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x)$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in \mathbb{N}) (2x > x) \quad \text{False}$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$(\forall x \in \mathbb{N}) (2x > x)$ **False** Consider $x = 0$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x)$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)(x > 5$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)(x > 5 \implies$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)(x > 5 \implies x^2 > 25).$$

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)(x > 5 \implies x^2 > 25).$$

Idea alert:

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)(x > 5 \implies x^2 > 25).$$

Idea alert: Restrict domain using implication.

More for all quantifiers examples.

- “doubling a natural number always makes it larger”

$$(\forall x \in N) (2x > x) \quad \text{False Consider } x = 0$$

Can fix statement...

$$(\forall x \in N) (2x \geq x) \quad \text{True}$$

- “Square of any natural number greater than 5 is greater than 25.”

$$(\forall x \in N)(x > 5 \implies x^2 > 25).$$

Idea alert: Restrict domain using implication.

Note that we may omit universe if clear from context.

Quantifiers..not commutative.

- In English: “there is a natural number that is the square of every natural number”.

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in N)$$

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in N) (\forall x \in N)$$

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in N) (\forall x \in N) (y = x^2)$$

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in \mathbb{N}) (\forall x \in \mathbb{N}) (y = x^2) \quad \text{False}$$

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in \mathbb{N}) (\forall x \in \mathbb{N}) (y = x^2) \quad \text{False}$$

- In English: “the square of every natural number is a natural number.”

Quantifiers..not commutative.

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in \mathbb{N}) (\forall x \in \mathbb{N}) (y = x^2) \quad \text{False}$$

- In English: “the square of every natural number is a natural number.”

$$(\forall x \in \mathbb{N})$$

Quantifiers..not commutative.

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in N) (\forall x \in N) (y = x^2) \quad \text{False}$$

- In English: “the square of every natural number is a natural number.”

$$(\forall x \in N)(\exists y \in N)$$

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in N) (\forall x \in N) (y = x^2) \quad \text{False}$$

- In English: “the square of every natural number is a natural number.”

$$(\forall x \in N)(\exists y \in N) (y = x^2)$$

Quantifiers..not commutative.

- In English: “there is a natural number that is the square of every natural number”.

$$(\exists y \in N) (\forall x \in N) (y = x^2) \quad \text{False}$$

- In English: “the square of every natural number is a natural number.”

$$(\forall x \in N)(\exists y \in N) (y = x^2) \quad \text{True}$$

Consider

$$\neg(\forall x \in S)(P(x)),$$

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

For **False**, find x , where $\neg P(x)$.

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

For **False**, find x , where $\neg P(x)$.

Counterexample.

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

For **False**, find x , where $\neg P(x)$.

Counterexample.

Bad input.

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

For **False**, find x , where $\neg P(x)$.

Counterexample.

Bad input.

Case that illustrates bug.

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

For **False**, find x , where $\neg P(x)$.

- Counterexample.

- Bad input.

- Case that illustrates bug.

For **True** : prove claim.

Quantifiers....negation...DeMorgan again.

Consider

$$\neg(\forall x \in S)(P(x)),$$

English: there is an x in S where $P(x)$ does not hold.

That is,

$$\neg(\forall x \in S)(P(x)) \iff \exists(x \in S)(\neg P(x)).$$

What we do in this course! We consider claims.

Claim: $(\forall x) P(x)$ “For all inputs x the program works.”

For **False**, find x , where $\neg P(x)$.

Counterexample.

Bad input.

Case that illustrates bug.

For **True** : prove claim. Next lectures...

Negation of exists.

Consider

Negation of exists.

Consider

$$\neg(\exists x \in S)(P(x))$$

Negation of exists.

Consider

$$\neg(\exists x \in S)(P(x))$$

English: means that for all x in S , $P(x)$ does not hold.

Negation of exists.

Consider

$$\neg(\exists x \in S)(P(x))$$

English: means that for all x in S , $P(x)$ does not hold.

That is,

$$\neg(\exists x \in S)(P(x)) \iff \forall(x \in S)\neg P(x).$$

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

$(\forall n \in \mathbb{N});$

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

$(\forall n \in \mathbb{N}); \neg(\exists a, b, c \in \mathbb{N});$

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

$$(\forall n \in \mathbb{N}); \neg(\exists a, b, c \in \mathbb{N}); (n \geq 3 \implies a^n + b^n = c^n)$$

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

$$(\forall n \in \mathbb{N}); \neg(\exists a, b, c \in \mathbb{N}); (n \geq 3 \implies a^n + b^n = c^n)$$

Using implication to state edge case restrictions (for any integer strictly greater than two)

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

$$(\forall n \in \mathbb{N}); \neg(\exists a, b, c \in \mathbb{N}); (n \geq 3 \implies a^n + b^n = c^n)$$

Using implication to state edge case restrictions (for any integer strictly greater than two)

DeMorgan Restatement:

Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer n strictly larger than two.

Which Theorem?

Fermat's Last Theorem!

How to express this theorem using propositions?

$$(\forall n \in \mathbb{N}); \neg(\exists a, b, c \in \mathbb{N}); (n \geq 3 \implies a^n + b^n = c^n)$$

Using implication to state edge case restrictions (for any integer strictly greater than two)

DeMorgan Restatement:

$$\text{Theorem: } \neg(\exists n \in \mathbb{N}) (\exists a, b, c \in \mathbb{N}) (n \geq 3 \implies a^n + b^n = c^n)$$

Summary.

Propositions are statements that are true or false.

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems!

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: “Flip and Distribute negation”

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: “Flip and Distribute negation”

$$\neg(P \vee Q) \iff$$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: “Flip and Distribute negation”

$$\neg(P \vee Q) \iff (\neg P \wedge \neg Q)$$

$$\neg \forall x P(x) \iff$$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: “Flip and Distribute negation”

$$\neg(P \vee Q) \iff (\neg P \wedge \neg Q)$$

$$\neg \forall x P(x) \iff \exists x \neg P(x).$$

$$\neg \exists x P(x) \iff$$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: “Flip and Distribute negation”

$$\neg(P \vee Q) \iff (\neg P \wedge \neg Q)$$

$$\neg \forall x P(x) \iff \exists x \neg P(x).$$

$$\neg \exists x P(x) \iff \forall x \neg P(x).$$

Summary.

Propositions are statements that are true or false.

Propositional forms use \wedge, \vee, \neg .

Propositional forms correspond to truth tables.

Logical equivalence of forms means same truth tables.

Implication: $P \implies Q \iff \neg P \vee Q$.

Contrapositive: $\neg Q \implies \neg P$

Converse: $Q \implies P$

Predicates: Statements with “free” variables.

Quantifiers: $\forall x P(x), \exists y Q(y)$

Now can state theorems! And disprove false ones!

DeMorgans Laws: “Flip and Distribute negation”

$$\neg(P \vee Q) \iff (\neg P \wedge \neg Q)$$

$$\neg \forall x P(x) \iff \exists x \neg P(x).$$

$$\neg \exists x P(x) \iff \forall x \neg P(x).$$

Next Time: proofs!