CS70: Counting

Alex Psomas

July 7, 2016

Today:

- Balls and bins.
- Sum rule.
- Combinatorial proofs.
- Maybe start review?

What we've learned so far

Sample k items out of n.

	With Replacement	Without Replacement
Order matters	n ^k	$\frac{n!}{(n-k)!}$
Order doesn't matter	$\binom{n+k-1}{n-1}$	$\binom{n}{k}$

Hats!

Hats! Say I have 10 different hats.

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

► Case 1: After I wear a hat I can wear it again.

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

Case 1: After I wear a hat I can wear it again. (Sampling with replacement)

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday.

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh.

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 10 options for Monday,

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday...

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷.

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing.

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)

How many samples?

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)
 - How many samples? a day is a sample, so 7.

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)
 - How many samples? a day is a sample, so 7. From how big of a set?

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)
 - How many samples? a day is a sample, so 7. From how big of a set? 10 hats.

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)

$$\binom{n+k-1}{n-1}$$

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)

$$\binom{n+k-1}{n-1} = \binom{10+7-1}{10-1}$$

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)

$$\binom{n+k-1}{n-1} = \binom{10+7-1}{10-1} = \binom{16}{9}$$

Hats! Say I have 10 different hats. I'm thinking of how many different outfits I have for one week:

- Case 1: After I wear a hat I can wear it again. (Sampling with replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. duh. (Order matters.)
 - 10 options for Monday, 10 options for Tuesday... 10⁷
 - Subcase: I don't care about which day I wore what, I just care which hats I ending up wearing. (Order doesn't matter.)

$$\binom{n+k-1}{n-1} = \binom{10+7-1}{10-1} = \binom{16}{9} = 11440$$

Case 2: After I wear a hat I destroy it.

 Case 2: After I wear a hat I destroy it. (Sampling without replacement)

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday,

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday...

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday...

```
\frac{n!}{(n-k)!}
```

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday...

$$\frac{n!}{(n-k)!} = \frac{10!}{3!}$$

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday...

$$\frac{n!}{(n-k)!} = \frac{10!}{3!} = 604800.$$

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday... $\frac{n!}{(n-k)!} = \frac{10!}{3!} = 604800.$
 - Subcase: I don't care about which day I wore what. (Order doesn't matter.)

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday... $\frac{n!}{(n-k)!} = \frac{10!}{3!} = 604800.$
 - Subcase: I don't care about which day I wore what. (Order doesn't matter.)

- Case 2: After I wear a hat I destroy it. (Sampling without replacement)
 - Subcase: I care about which day I wore what: blue hat on Monday is different than blue hat on Tuesday. (Order matters.)
 - 10 options for Monday, 9 options for Tuesday...

$$\frac{n!}{(n-k)!} = \frac{10!}{3!} = 604800.$$

Subcase: I don't care about which day I wore what. (Order doesn't matter.)

$$\binom{10}{7} = 120$$

How many (non-negative) solutions to x + y = 10?

How many (non-negative) solutions to x + y = 10? Easy:

How many (non-negative) solutions to x + y = 10? Easy: x = 0, y = 10,

How many (non-negative) solutions to x + y = 10? Easy: x = 0, y = 10, x = 1, y = 9,

How many (non-negative) solutions to x + y = 10? Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions.

How many (non-negative) solutions to x + y = 10? Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions. Same as 10 stars, and 1 bar.

How many (non-negative) solutions to x + y = 10?

Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions.

Same as 10 stars, and 1 bar.

$$x = 3, y = 7$$
: $\star \star \star | \star \star \star \star \star \star \star$

How many (non-negative) solutions to x + y = 10?

Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions.

Same as 10 stars, and 1 bar.

x = 3, y = 7: $\star \star \star | \star \star \star \star \star \star \star$

Think of a star as the number 1.

How many (non-negative) solutions to x + y = 10?

Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions.

Same as 10 stars, and 1 bar.

$$x = 3, y = 7$$
: $\star \star \star | \star \star \star \star \star \star \star$

Think of a star as the number 1.

How many ways to make an 8 problem midterm such the total points add up to 100?

How many (non-negative) solutions to x + y = 10?

Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions.

Same as 10 stars, and 1 bar.

$$x = 3, y = 7$$
: $\star \star \star | \star \star \star \star \star \star \star$

Think of a star as the number 1.

How many ways to make an 8 problem midterm such the total points add up to 100? 100 stars,

How many (non-negative) solutions to x + y = 10?

Easy: x = 0, y = 10, x = 1, y = 9, x = 2, y = 8, ..., x = 10, y = 0. So 11 solutions.

Same as 10 stars, and 1 bar.

$$x = 3, y = 7:$$
 $\star \star \star | \star \star \star \star \star \star \star$

Think of a star as the number 1.

How many ways to make an 8 problem midterm such the total points add up to 100?

100 stars, 7 bars.





"k Balls in n bins" \equiv "k samples from n possibilities."



"k Balls in n bins" \equiv "k samples from n possibilities."

"indistinguishable balls" \equiv "order doesn't matter"




"k Balls in n bins" \equiv "k samples from n possibilities." "indistinguishable balls" \equiv "order doesn't matter" "only one ball in each bin" \equiv "without replacement"


How many 5 digit numbers?

How many 5 digit numbers? Throwing 5 numbered balls in 10 (numbered) bins:

How many 5 digit numbers? Throwing 5 numbered balls in 10 (numbered) bins:




How many 5 digit numbers? Throwing 5 numbered balls in 10 (numbered) bins:



Picture has number 62280.

How many 5 digit numbers?
Throwing 5 numbered balls in 10 (numbered) bins:




Picture has number 62280.

5 samples from 10 possibilities with replacement (order matters): 10⁵


How many 5 digit numbers without repeating a digit?

How many 5 digit numbers without repeating a digit? Throwing 5 numbered balls in 10 (numbered) bins, at most one ball per bin:

How many 5 digit numbers without repeating a digit? Throwing 5 numbered balls in 10 (numbered) bins, at most one ball per bin:




How many 5 digit numbers without repeating a digit? Throwing 5 numbered balls in 10 (numbered) bins, at most one ball per bin:



Picture has number 62480.

How many 5 digit numbers without repeating a digit? Throwing 5 numbered balls in 10 (numbered) bins, at most one ball per bin:



Picture has number 62480.


5 samples from 10 possibilities without replacement (order matters):

10! 5!


How many 3 card poker hands?

How many 3 card poker hands? Throwing 3 indistinguishable balls in 52 (numbered) bins. At most 1 ball per bin:

How many 3 card poker hands? Throwing 3 indistinguishable balls in 52 (numbered) bins. At most 1 ball per bin:




How many 3 card poker hands? Throwing 3 indistinguishable balls in 52 (numbered) bins. At most 1 ball per bin:



Picture has cards 1,3 and 52.

How many 3 card poker hands?

Throwing 3 indistinguishable balls in 52 (numbered) bins. At most 1 ball per bin:




Picture has cards 1,3 and 52.

3 samples from 52 possibilities without replacement (order doesn't matter): $\binom{52}{3}$


Dividing 5 dollars among Alice, Bob and Eve.

Dividing 5 dollars among Alice, Bob and Eve. 5 indistinguishable balls into 3 (numbered) bins:

Dividing 5 dollars among Alice, Bob and Eve. 5 indistinguishable balls into 3 (numbered) bins:




Dividing 5 dollars among Alice, Bob and Eve. 5 indistinguishable balls into 3 (numbered) bins:



Picture: Alice 3, Bob 0, Eve 2.

Dividing 5 dollars among Alice, Bob and Eve. 5 indistinguishable balls into 3 (numbered) bins:



Picture: Alice 3, Bob 0, Eve 2. 5 samples from 3 possibilities with replacement (order doesn't matter): $\binom{7}{2}$

Sum Rule

Two indistinguishable jokers in 54 card deck. How many 5 card poker hands?

Two indistinguishable jokers in 54 card deck. How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

Two indistinguishable jokers in 54 card deck. How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

(52) 5

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4}$$

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck.

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}.$$

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)?

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$${52 \choose 5} + {52 \choose 4} + {52 \choose 3}.$$

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} +$$

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\left(\begin{smallmatrix}52\\5\end{smallmatrix}\right) + 2*\left(\begin{smallmatrix}52\\4\end{smallmatrix}\right) +$$

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$${52 \choose 5} + {52 \choose 4} + {52 \choose 3}.$$

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$${52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}$$

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$${52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}$$

Wait a minute!

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$${52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}$$

Wait a minute! Same as

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2 * \binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2*\binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54 or

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2 * \binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54 or

$$\binom{54}{5}$$

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2 * \binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54 or

$$\binom{54}{5}$$

Theorem: $\binom{54}{5}$

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2*\binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54 or

$$\binom{54}{5}$$

Theorem: $\binom{54}{5} = \binom{52}{5} + 2 * \binom{52}{4} + \binom{52}{3}$.

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2*\binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54 or

$$\binom{54}{5}$$

Theorem: $\binom{54}{5} = \binom{52}{5} + 2 * \binom{52}{4} + \binom{52}{3}$.

Two indistinguishable jokers in 54 card deck.

How many 5 card poker hands?

Sum rule: Can sum over disjoint sets.

No jokers or exactly one joker or exactly two jokers

$$\binom{52}{5} + \binom{52}{4} + \binom{52}{3}$$
.

Two distinguishable jokers in 54 card deck. How many 5 card poker hands (distinguishable jokers)? No jokers or exactly one of two jokers or exactly two jokers

$$\binom{52}{5} + 2*\binom{52}{4} + \binom{52}{3}$$

Wait a minute! Same as choosing 5 cards from 54 or

$$\binom{54}{5}$$

Theorem: $\binom{54}{5} = \binom{52}{5} + 2 * \binom{52}{4} + \binom{52}{3}$. **Proof:** Above is combinatorial proof.

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$$\binom{54}{5} = \tfrac{54!}{5!49!}$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$$\binom{54}{5} = \frac{54!}{5!49!} \ , \quad \ \binom{52}{5} = \frac{52!}{5!47!}$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = \frac{54!}{5!49!} \ , \quad {52 \choose 5} = \frac{52!}{5!47!} \ , \quad {52 \choose 4} = \frac{52!}{4!48!}$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = \frac{54!}{5!49!} \ , \quad {52 \choose 5} = \frac{52!}{5!47!} \ , \quad {52 \choose 4} = \frac{52!}{4!48!} \ , \quad {52 \choose 3} = \frac{52!}{3!49!}$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

Proof:

$$\binom{54}{5} = \frac{54!}{5!49!} \ , \qquad \binom{52}{5} = \frac{52!}{5!47!} \ , \qquad \binom{52}{4} = \frac{52!}{4!48!} \ , \qquad \binom{52}{3} = \frac{52!}{3!49!}$$

RHS

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

Proof:

 $54!47!4!48!3! \stackrel{?}{=} 52!(4!48!3!49! + 2*5!47!3!49! + 5!47!4!48!)$ I tried this for a while...

$${54 \choose 5} = {52 \choose 5} + 2*{52 \choose 4} + {52 \choose 3}.$$

Proof:

49! and 5! cancel out. Cross multiply and get:

 $54!47!4!48!3! \stackrel{?}{=} 52!(4!48!3!49! + 2*5!47!3!49! + 5!47!4!48!)$

I tried this for a while......

Theorem:
$$\binom{n}{k} = \binom{n}{n-k}$$

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size *k*?

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k?

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k? Choose a subset of size n-k

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k? Choose a subset of size n-kand what's left out

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k?

Choose a subset of size n - kand what's left out is a subset of size k.

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k? Choose a subset of size n-k and what's left out is a subset of size k.

Choosing a subset of size k is same

Theorem: $\binom{n}{k} = \binom{n}{n-k}$

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k? Choose a subset of size n-k

and what's left out is a subset of size k.

Choosing a subset of size k is same as choosing n - k elements to not take.

```
Theorem: \binom{n}{k} = \binom{n}{n-k}
```

Proof: How many subsets of size k? $\binom{n}{k}$

How many subsets of size k?

Choose a subset of size n-k and what's left out is a subset of size k.

Choosing a subset of size k is same as choosing n - k elements to not take.

 $\implies \binom{n}{n-k}$ subsets of size k.

```
Theorem: \binom{n}{k} = \binom{n}{n-k}
Proof: How many subsets of size k? \binom{n}{k}
How many subsets of size k?
Choose a subset of size n-k
and what's left out is a subset of size k.
Choosing a subset of size k is same
as choosing n-k elements to not take.
\implies \binom{n}{n-k} subsets of size k.
```

1 1 1

```
1
1 1
1 2 1
```

```
1
1 1
1 2 1
1 3 3
```

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
```

```
1 1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 1 5 10 10 5 1
```

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1

Row n: coefficients of (1+x)^n = (1+x)(1+x)\cdots(1+x).
```

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1

Row n: coefficients of (1+x)^n = (1+x)(1+x)\cdots(1+x).

Zero-th row: (1+x)^0 = 1

First row: (1+x)^1 = x+1. Coefficients: 1 and 1
```

```
\begin{array}{c} 1 & 1 \\ 1 & 2 & 1 \\ 1 & 3 & 3 & 1 \\ 1 & 4 & 6 & 4 & 1 \\ 1 & 5 & 10 & 10 & 5 & 1 \\ \end{array} Row n: coefficients of (1+x)^n=(1+x)(1+x)\cdots(1+x). Zero-th row: (1+x)^0=1 First row: (1+x)^1=x+1. Coefficients: 1 and 1 Second row: (1+x)^2=1+2x+x^2. Coefficients: 1,2 and 1 Third row: (1+x)^3=1
```

```
1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 5 10 10 5 1 Row n: coefficients of (1+x)^n=(1+x)(1+x)\cdots(1+x). Zero-th row: (1+x)^0=1 First row: (1+x)^1=x+1. Coefficients: 1 and 1 Second row: (1+x)^2=1+2x+x^2. Coefficients: 1,2 and 1 Third row: (1+x)^3=1+3x+3x^2+x^3. Coefficients: 1,3,3 and 1
```

```
Row n: coefficients of (1+x)^n = (1+x)(1+x)\cdots(1+x).
Zero-th row: (1 + x)^0 = 1
First row: (1+x)^1 = x+1. Coefficients: 1 and 1
Second row: (1+x)^2 = 1+2x+x^2. Coefficients: 1,2 and 1
Third row: (1+x)^3 = 1 + 3x + 3x^2 + x^3. Coefficients: 1,3,3 and 1
```

```
Row n: coefficients of (1+x)^n = (1+x)(1+x)\cdots(1+x).
Zero-th row: (1 + x)^0 = 1
First row: (1+x)^1 = x+1. Coefficients: 1 and 1
Second row: (1+x)^2 = 1+2x+x^2. Coefficients: 1,2 and 1
Third row: (1+x)^3 = 1 + 3x + 3x^2 + x^3. Coefficients: 1,3,3 and 1
Foil??
```

```
Row n: coefficients of (1+x)^n = (1+x)(1+x)\cdots(1+x).
Zero-th row: (1 + x)^0 = 1
First row: (1+x)^1 = x+1. Coefficients: 1 and 1
Second row: (1+x)^2 = 1+2x+x^2. Coefficients: 1,2 and 1
Third row: (1+x)^3 = 1 + 3x + 3x^2 + x^3. Coefficients: 1,3,3 and 1
Foil?? I hate this already...
```

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k .

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:

```
1 1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product.

```
1 1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x):

```
1 1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from:

```
1 1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term

```
1 1 1 1 1 2 1 1 3 3 1 1 4 6 4 1 1 1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term , first and third term

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term , first and third term , first and fourth

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and fourth, third and fourth.

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and third, second and fourth, third and fourth. 6 of them

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and third, second and fourth, third and fourth. 6 of them, aka $\binom{4}{2}$.

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and third, second and fourth, third and fourth. 6 of them, aka $\binom{4}{2}$.

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product. (1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and third, second and fourth, third and fourth. 6 of them, aka $\binom{4}{2}$.

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} 2 \\ 0 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} \begin{pmatrix} 2 \\ 2 \end{pmatrix}$$

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k . Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product.

(1+x)(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and third, second and fourth, third and fourth. 6 of them, aka $\binom{4}{2}$.

 $\begin{pmatrix} \binom{0}{0} \\ \binom{1}{0} & \binom{1}{1} \\ \binom{2}{0} & \binom{2}{1} & \binom{2}{2} \\ \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \binom{3}{3} \end{pmatrix}$

```
1
1 1
1 2 1
1 3 3 1
1 4 6 4 1
1 5 10 10 5 1
```

Simplify: collect all terms corresponding to x^k .

Coefficient of x^k is $\binom{n}{k}$:choose k factors where x is in product.

(1+x)(1+x)(1+x): Coefficients of x^2 come from: first and second term, first and third term, first and fourth, second and third, second and fourth, third and fourth. 6 of them, aka $\binom{4}{2}$.

$$\begin{pmatrix} \binom{0}{0} \\ \binom{1}{0} & \binom{1}{1} \\ \binom{2}{0} & \binom{2}{1} & \binom{2}{2} \\ \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \binom{3}{3} \end{pmatrix}$$

Pascal's rule
$$\implies \binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$$
.

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size k subsets of n+1?

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size k subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size k subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1? The ones that contain the first element

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first.

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

$$\implies \binom{n}{k-1}$$

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

$$\implies \binom{n}{k-1}$$

How many don't contain the first element?

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

$$\implies \binom{n}{k-1}$$

How many don't contain the first element?

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

$$\implies \binom{n}{k-1}$$

How many don't contain the first element?

$$\implies \binom{n}{k}$$

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

$$\implies \binom{n}{k-1}$$

How many don't contain the first element?

$$\Longrightarrow \binom{n}{k}$$

So, $\binom{n}{k-1} + \binom{n}{k}$

Theorem: $\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}$.

Proof: How many size *k* subsets of n+1? $\binom{n+1}{k}$.

How many size k subsets of n+1?

The ones that contain the first element plus the ones that don't contain the first element.

How many contain the first element?

Pick the first. Then I need to choose k-1 more from remaining n elements.

$$\implies \binom{n}{k-1}$$

How many don't contain the first element?

$$\implies \binom{n}{k}$$

So,
$$\binom{n}{k-1} + \binom{n}{k} = \binom{n+1}{k}$$
.

Theorem:
$$\binom{n}{k} = \binom{n-1}{k-1} + \cdots + \binom{k-1}{k-1}$$
.

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$. **Proof:**

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \cdots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \cdots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

Theorem:
$$\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$$
. **Proof:**

Left Hand Side (LHS): Size k subsets of n.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements.

Theorem:
$$\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$$
.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements.

$$\implies \binom{n-i}{k-1}$$
 such subsets.

Theorem:
$$\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$$
. Proof:

Left Hand Side (LHS): Size k subsets of n.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements.

 $\implies \binom{n-i}{k-1}$ such subsets.

1 is smallest element chosen:

Theorem:
$$\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$$
.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements.

$$\implies \binom{n-i}{k-1}$$
 such subsets.

1 is smallest element chosen: $\binom{n-1}{k-1}$ choices for the rest.

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements.

 $\implies \binom{n-i}{k-1}$ such subsets.

1 is smallest element chosen: $\binom{n-1}{k-1}$ choices for the rest.

2 is smallest element chosen:

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements.

 $\implies \binom{n-i}{k-1}$ such subsets.

1 is smallest element chosen: $\binom{n-1}{k-1}$ choices for the rest.

2 is smallest element chosen: $\binom{n-2}{k-1}$ choices for the rest.

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{\textit{i}},\ldots,\underline{\textit{n}}\}$$

Must choose k-1 elements from n-i remaining elements.

 $\implies \binom{n-i}{k-1}$ such subsets.

1 is smallest element chosen: $\binom{n-1}{k-1}$ choices for the rest.

2 is smallest element chosen: $\binom{n-2}{k-1}$ choices for the rest. and so on.

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \dots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size *k* subset where *i* is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements. $\Rightarrow \binom{n-i}{k-1}$ such subsets.

1 is smallest element chosen: $\binom{n-1}{k-1}$ choices for the rest.

2 is smallest element chosen: $\binom{n-2}{k-1}$ choices for the rest. and so on.

Add them up to get the total number of subsets of size k

Combinatorial Proof.

Theorem: $\binom{n}{k} = \binom{n-1}{k-1} + \cdots + \binom{k-1}{k-1}$.

Proof:

Left Hand Side (LHS): Size *k* subsets of *n*.

Consider size k subset where i is the smallest element chosen.

$$\{1,\ldots,\underline{i},\ldots,\underline{n}\}$$

Must choose k-1 elements from n-i remaining elements. $\implies \binom{n-i}{k-1}$ such subsets.

1 is smallest element chosen: $\binom{n-1}{k-1}$ choices for the rest.

2 is smallest element chosen: $\binom{n-2}{k-1}$ choices for the rest. and so on.

Add them up to get the total number of subsets of size *k* which is also $\binom{n}{k}$.

Theorem:
$$2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$$

Theorem: $2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$

Proof: How many subsets of $\{1,...,n\}$?

Theorem: $2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices:

Theorem:
$$2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$$

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of n choices:

element *i* is in or is not in the subset: 2 possibilities.

Theorem:
$$2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$$

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices: element *i* **is in** or **is not** in the subset: 2 possibilities.

First rule of counting: $2 \times 2 \cdots \times 2 = 2^n$ subsets.

Theorem:
$$2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$$

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices: element *i* **is in** or **is not** in the subset: 2 possibilities.

First rule of counting: $2 \times 2 \cdots \times 2 = 2^n$ subsets.

How many subsets of $\{1, \ldots, n\}$?

```
Theorem: 2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}
```

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices: element *i* **is in** or **is not** in the subset: 2 possibilities.

First rule of counting: $2 \times 2 \cdots \times 2 = 2^n$ subsets.

How many subsets of $\{1,\ldots,n\}$?

$$\binom{n}{i} =$$

```
Theorem: 2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}
```

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices: element *i* **is in** or **is not** in the subset: 2 possibilities.

First rule of counting: $2 \times 2 \cdots \times 2 = 2^n$ subsets.

How many subsets of $\{1, ..., n\}$? $\binom{n}{i} = \text{subsets of size } i.$

Theorem:
$$2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$$

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices: element *i* **is in** or **is not** in the subset: 2 possibilities.

First rule of counting: $2 \times 2 \cdots \times 2 = 2^n$ subsets.

How many subsets of $\{1, \ldots, n\}$?

 $\binom{n}{i}$ = subsets of size *i*.

A subset has size either 0, or 1, or 2, ..., or n

Theorem:
$$2^n = \binom{n}{n} + \binom{n}{n-1} + \cdots + \binom{n}{0}$$

Proof: How many subsets of $\{1, ..., n\}$?

Construct a subset with sequence of *n* choices: element *i* **is in** or **is not** in the subset: 2 possibilities.

First rule of counting: $2 \times 2 \cdots \times 2 = 2^n$ subsets.

How many subsets of $\{1, \ldots, n\}$?

 $\binom{n}{i}$ = subsets of size *i*.

A subset has size either 0, or 1, or 2, ..., or n

Sum over *i* to get total number of subsets.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first i elements.)

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit. $|S| = 10^9$

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit. $|S| = 10^9$

T = phone numbers with 7 as second digit.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit. $|S| = 10^9$

T = phone numbers with 7 as second digit. $|T| = 10^9$.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit. $|S| = 10^9$

T = phone numbers with 7 as second digit. $|T| = 10^9$.

 $S \cap T$ = phone numbers with 7 as first and second digit.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit. $|S| = 10^9$

T = phone numbers with 7 as second digit. $|T| = 10^9$.

 $S \cap T$ = phone numbers with 7 as first and second digit. $|S \cap T| = 10^8$.

Sum Rule: For disjoint sets S and T, $|S \cup T| = |S| + |T|$ Used to reason about all subsets by adding number of subsets of size 0, 1, 2, 3,...

Also reasoned about subsets that contained or didn't contain an element. (E.g., first element, first *i* elements.)

Inclusion/Exclusion Rule: For any S and T, $|S \cup T| = |S| + |T| - |S \cap T|$.

Example: How many 10-digit phone numbers have 7 as their first or second digit?

S = phone numbers with 7 as first digit. $|S| = 10^9$

T = phone numbers with 7 as second digit. $|T| = 10^9$.

 $S \cap T$ = phone numbers with 7 as first and second digit. $|S \cap T| = 10^8$.

Answer: $|S| + |T| - |S \cap T| = 10^9 + 10^9 - 10^8$.

Inclusion/Exclusion: two sets of objects.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Pascal's Triangle Example: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Pascal's Triangle Example: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$.

RHS: Number of subsets of n+1 items size k.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Pascal's Triangle Example: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$.

RHS: Number of subsets of n+1 items size k.

LHS: $\binom{n}{k-1}$ counts subsets of n+1 items with first item.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Pascal's Triangle Example: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$.

RHS: Number of subsets of n+1 items size k.

LHS: $\binom{n}{k-1}$ counts subsets of n+1 items with first item.

 $\binom{n}{k}$ counts subsets of n+1 items without first item.

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Pascal's Triangle Example: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$.

RHS: Number of subsets of n+1 items size k.

LHS: $\binom{n}{k-1}$ counts subsets of n+1 items with first item.

 $\binom{n}{k}$ counts subsets of n+1 items without first item.

Disjoint

Inclusion/Exclusion: two sets of objects.

Add number of each subtract intersection of sets.

Sum Rule: If disjoint just add.

Combinatorial Proofs: Identity from counting same in two ways.

Pascal's Triangle Example: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$.

RHS: Number of subsets of n+1 items size k.

LHS: $\binom{n}{k-1}$ counts subsets of n+1 items with first item.

 $\binom{n}{k}$ counts subsets of n+1 items without first item.

Disjoint – so add!