

Alex Psomas: Lecture 14.

Events, Conditional Probability, Independence, Bayes' Rule

Alex Psomas: Lecture 14.

Events, Conditional Probability, Independence, Bayes' Rule

1. Probability Basics Review
2. Conditional Probability
3. Independence of Events
4. Bayes' Rule

Probability Basics Review

Probability Basics Review

Setup:

Probability Basics Review

Setup:

- ▶ Random Experiment.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 $Pr[HH] = \dots = Pr[TT] = 1/4$

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 $Pr[HH] = \dots = Pr[TT] = 1/4$
 1. $0 \leq Pr[\omega] \leq 1$.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 $Pr[HH] = \dots = Pr[TT] = 1/4$
 1. $0 \leq Pr[\omega] \leq 1$.
 2. $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 $Pr[HH] = \dots = Pr[TT] = 1/4$
 1. $0 \leq Pr[\omega] \leq 1$.
 2. $\sum_{\omega \in \Omega} Pr[\omega] = 1$.
- ▶ Event.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 $Pr[HH] = \dots = Pr[TT] = 1/4$
 1. $0 \leq Pr[\omega] \leq 1$.
 2. $\sum_{\omega \in \Omega} Pr[\omega] = 1$.
- ▶ Event. Set of the outcomes.

Probability Basics Review

Setup:

- ▶ Random Experiment.
Flip a fair coin twice.
- ▶ Probability Space.
 - ▶ **Sample Space:** Set of outcomes, Ω .
 $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
 - ▶ **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 $Pr[HH] = \dots = Pr[TT] = 1/4$
 1. $0 \leq Pr[\omega] \leq 1$.
 2. $\sum_{\omega \in \Omega} Pr[\omega] = 1$.
- ▶ Event. Set of the outcomes.

Probability is Additive

Theorem

Probability is Additive

Theorem

(a) If events A and B are disjoint,

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$,

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint,

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint,
i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$,

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are **pairwise** disjoint,
i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] =$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] =$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] +$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega]$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega] = Pr[A] + Pr[B]$$

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega] = Pr[A] + Pr[B]$$

Can I instead say that $|A \cup B| = |A| + |B|$?

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega] = Pr[A] + Pr[B]$$

Can I instead say that $|A \cup B| = |A| + |B|$?

No!

Probability is Additive

Theorem

(a) If events A and B are disjoint, i.e., $A \cap B = \emptyset$, then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events A_1, \dots, A_n are pairwise disjoint, i.e., $A_k \cap A_m = \emptyset, \forall k \neq m$, then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega] = Pr[A] + Pr[B]$$

Can I instead say that $|A \cup B| = |A| + |B|$?

No! We don't know if the sample space is uniform.

Consequences of Additivity

Theorem

Consequences of Additivity

Theorem

$$(a) \Pr[A \cup B] = \Pr[A] + \Pr[B] - \Pr[A \cap B];$$

Consequences of Additivity

Theorem

$$(a) \Pr[A \cup B] = \Pr[A] + \Pr[B] - \Pr[A \cap B];$$

(inclusion-exclusion property)

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B];$
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n];$
(union bound)

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;
(union bound)
- (c) If A_1, \dots, A_N are a partition of Ω ,

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;
(union bound)
- (c) If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$,

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;
(union bound)
- (c) If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then
$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B];$
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n];$
(union bound)
- (c) If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then
$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

(law of total probability)

Consequences of Additivity

Theorem

- (a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
(inclusion-exclusion property)
- (b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;
(union bound)
- (c) If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then
$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

(law of total probability)

Proof:

Consequences of Additivity

Theorem

(a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B];$
(inclusion-exclusion property)

(b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n];$
(union bound)

(c) If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then

$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

(law of total probability)

Proof:

(b) is obvious.

Consequences of Additivity

Theorem

(a) $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B];$
(inclusion-exclusion property)

(b) $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n];$
(union bound)

(c) If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then

$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

(law of total probability)

Proof:

(b) is obvious.


See next two slides for (a) and (c).

Inclusion/Exclusion

$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$

Inclusion/Exclusion


$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$



$$\begin{aligned}Pr[A] &= x + y \\Pr[B] &= y + z \\Pr[A \cap B] &= y \\Pr[A \cup B] &= x + y + z\end{aligned}$$

Inclusion/Exclusion

$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$




$$\begin{aligned} Pr[A] &= x + y \\ Pr[B] &= y + z \\ Pr[A \cap B] &= y \\ Pr[A \cup B] &= x + y + z \end{aligned}$$

Can I instead say that $|A \cup B| = |A| + |B| - |A \cap B|$?

Inclusion/Exclusion

$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$




$$\begin{aligned}Pr[A] &= x + y \\Pr[B] &= y + z \\Pr[A \cap B] &= y \\Pr[A \cup B] &= x + y + z\end{aligned}$$

Can I instead say that $|A \cup B| = |A| + |B| - |A \cap B|$?

No!

Inclusion/Exclusion

$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$




$$\begin{aligned} Pr[A] &= x + y \\ Pr[B] &= y + z \\ Pr[A \cap B] &= y \\ Pr[A \cup B] &= x + y + z \end{aligned}$$

Can I instead say that $|A \cup B| = |A| + |B| - |A \cap B|$?

No! We don't know if the sample space is uniform.


Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .



Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .




Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .




Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Indeed, B is the union of the disjoint sets $A_n \cap B$ for $n = 1, \dots, N$.


Roll a Red and a Blue Die.

Roll a Red and a Blue Die.



$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$


Roll a Red and a Blue Die.



$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

E_1 = 'Red die shows 6';


Roll a Red and a Blue Die.



$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

E_1 = 'Red die shows 6'; E_2 = 'Blue die shows 6'

Roll a Red and a Blue Die.




$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

E_1 = 'Red die shows 6'; E_2 = 'Blue die shows 6'

$E_1 \cup E_2$ = 'At least one die shows 6'

Roll a Red and a Blue Die.




$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

E_1 = 'Red die shows 6'; E_2 = 'Blue die shows 6'

$E_1 \cup E_2$ = 'At least one die shows 6'

$$Pr[E_1] = \frac{6}{36},$$

Roll a Red and a Blue Die.




$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

E_1 = 'Red die shows 6'; E_2 = 'Blue die shows 6'

$E_1 \cup E_2$ = 'At least one die shows 6'

$$Pr[E_1] = \frac{6}{36}, Pr[E_2] = \frac{6}{36},$$

Roll a Red and a Blue Die.



$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

E_1 = 'Red die shows 6'; E_2 = 'Blue die shows 6'

$E_1 \cup E_2$ = 'At least one die shows 6'

$$Pr[E_1] = \frac{6}{36}, Pr[E_2] = \frac{6}{36}, Pr[E_1 \cup E_2] = \frac{11}{36}.$$

Conditional probability: example.

Two coin flips (fair coin).

Conditional probability: example.

Two coin flips (fair coin). First flip is heads.

Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$$\Omega = \{HH, HT, TH, TT\};$$

Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads:

Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.


Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform




Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ;


Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ; uniform still.


Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ; uniform still.




Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ; uniform still.



Event B = two heads.


Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ; uniform still.



Event B = two heads.

The probability of two heads if the first flip is heads.


Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ; uniform still.



Event B = two heads.

The probability of two heads if the first flip is heads.

The probability of B given A


Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$; Uniform probability space.

Event A = first flip is heads: $A = \{HH, HT\}$.

Ω : uniform



New sample space: A ; uniform still.



Event B = two heads.

The probability of two heads if the first flip is heads.

The probability of B given A is $1/2$.

A similar example.

Two coin flips(fair coin).

A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$$\Omega = \{HH, HT, TH, TT\};$$

A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads.

A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.


A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.




A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ;


A similar example.

Two coin flips(fair coin). At least one of the flips is heads.

→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ; uniform still.


A similar example.

Two coin flips(fair coin). At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ; uniform still.




A similar example.

Two coin flips(fair coin). At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ; uniform still.



Event B = two heads.


A similar example.

Two coin flips(fair coin). At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ; uniform still.



Event B = two heads.

The probability of two heads if at least one flip is heads.


A similar example.

Two coin flips(fair coin). At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ; uniform still.



Event B = two heads.

The probability of two heads if at least one flip is heads.

The probability of B given A


A similar example.

Two coin flips(fair coin). At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$; uniform.

Event A = at least one flip is heads. $A = \{HH, HT, TH\}$.



New sample space: A ; uniform still.




Event B = two heads.

The probability of two heads if at least one flip is heads.


The probability of B given A is $1/3$.

Conditional Probability: A non-uniform example

Conditional Probability: A non-uniform example




Physical experiment




Probability model

Conditional Probability: A non-uniform example




Physical experiment




Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

Conditional Probability: A non-uniform example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

$$Pr[\text{Red} | \text{Red or Green}] =$$

Conditional Probability: A non-uniform example



Physical experiment




Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

$$Pr[\text{Red} | \text{Red or Green}] = \frac{3}{7} =$$

Conditional Probability: A non-uniform example



Physical experiment



Probability model

$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

$$Pr[\text{Red} | \text{Red or Green}] = \frac{3}{7} = \frac{Pr[\text{Red} \cap (\text{Red or Green})]}{Pr[\text{Red or Green}]}$$

Another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Another non-uniform example


Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Let $A = \{3, 4\}$, $B = \{1, 2, 3\}$.

Another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.


Let $A = \{3, 4\}$, $B = \{1, 2, 3\}$.



Another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Let $A = \{3, 4\}, B = \{1, 2, 3\}$.




$$Pr[A|B] =$$

Another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Let $A = \{3, 4\}, B = \{1, 2, 3\}$.



$$Pr[A|B] = \frac{p_3}{p_1 + p_2 + p_3} = \frac{Pr[A \cap B]}{Pr[B]}.$$

Yet another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Yet another non-uniform example


Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Let $A = \{2, 3, 4\}$, $B = \{1, 2, 3\}$.

Yet another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.


Let $A = \{2, 3, 4\}$, $B = \{1, 2, 3\}$.



Yet another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Let $A = \{2, 3, 4\}$, $B = \{1, 2, 3\}$.




$$Pr[A|B] =$$

Yet another non-uniform example

Consider $\Omega = \{1, 2, \dots, N\}$ with $Pr[n] = p_n$.

Let $A = \{2, 3, 4\}$, $B = \{1, 2, 3\}$.




$$Pr[A|B] = \frac{p_2 + p_3}{p_1 + p_2 + p_3} = \frac{Pr[A \cap B]}{Pr[B]}.$$

Conditional Probability.

Definition: The **conditional probability** of B given A is


$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$




In A !

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$




In A !

In B ?

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$




In A !
In B ?
Must be in $A \cap B$.

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In A !

In B ?

Must be in $A \cap B$.

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

More fun with conditional probability.


Toss a red and a blue die, sum is 4,

More fun with conditional probability.

Toss a red and a blue die, sum is 4,
What is probability that red is 1?


More fun with conditional probability.

Toss a red and a blue die, sum is 4,
What is probability that red is 1?



More fun with conditional probability.


Toss a red and a blue die, sum is 4,
What is probability that red is 1?



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{3};$$

More fun with conditional probability.


Toss a red and a blue die, sum is 4,
What is probability that red is 1?



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{3}; \text{ versus } Pr[B] = 1/6.$$

More fun with conditional probability.

Toss a red and a blue die, sum is 4,
What is probability that red is 1?



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{3}; \text{ versus } Pr[B] = 1/6.$$


B is more likely given A .

Yet more fun with conditional probability.

Toss a red and a blue die, sum is 7,
what is probability that red is 1?


Yet more fun with conditional probability.

Toss a red and a blue die, sum is 7,
what is probability that red is 1?



Yet more fun with conditional probability.


Toss a red and a blue die, sum is 7,
what is probability that red is 1?



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{6};$$

Yet more fun with conditional probability.


Toss a red and a blue die, sum is 7,
what is probability that red is 1?



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{6}; \text{ versus } Pr[B] = \frac{1}{6}.$$

Yet more fun with conditional probability.

Toss a red and a blue die, sum is 7,
what is probability that red is 1?



$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{6}; \text{ versus } Pr[B] = \frac{1}{6}.$$

Observing A does not change your mind about the likelihood of B .

Emptiness..

Suppose I toss 3 balls into 3 bins.

Emptiness..

Suppose I toss 3 balls into 3 bins.

$A = \text{"1st bin empty"};$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = “1st bin empty”; B = “2nd bin empty.”

Emptiness..


Suppose I toss 3 balls into 3 bins.

A = “1st bin empty”; B = “2nd bin empty.” What is $Pr[A|B]$?

Emptiness..

Suppose I toss 3 balls into 3 bins.


A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?



Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$Pr[B]$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] =$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] =$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B]$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B] = Pr[(3, 3, 3)] =$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?




$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$


$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B]$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$


$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$


$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]} = \frac{(1/27)}{(8/27)} = 1/8;$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$


$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]} = \frac{(1/27)}{(8/27)} = 1/8; \text{ vs. } Pr[A] = \frac{8}{27}.$$

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = “1st bin empty”; B = “2nd bin empty.” What is $Pr[A|B]$?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$


$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]} = \frac{(1/27)}{(8/27)} = 1/8; \text{ vs. } Pr[A] = \frac{8}{27}.$$

A is less likely given B :

Emptiness..

Suppose I toss 3 balls into 3 bins.

A = "1st bin empty"; B = "2nd bin empty." What is $Pr[A|B]$?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]} = \frac{(1/27)}{(8/27)} = 1/8; \text{ vs. } Pr[A] = \frac{8}{27}.$$

A is less likely given B : If second bin is empty the first is more likely to have balls in it.

Gambler's fallacy.

Flip a fair coin 51 times.

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

$A = \{HH \dots HT, HH \dots HH\}$

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Uniform probability space.

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Uniform probability space.

$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{2}.$$

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Uniform probability space.

$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{2}.$$

Same as $Pr[B]$.

Gambler's fallacy.

Flip a fair coin 51 times.

A = “first 50 flips are heads”

B = “the 51st is heads”

$Pr[B|A]$?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Uniform probability space.

$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{2}.$$

Same as $Pr[B]$.

The likelihood of 51st heads does not depend on the previous flips.

Product Rule

Recall the definition:

Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Hence,

$$Pr[A \cap B] = Pr[A]Pr[B|A].$$

Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Hence,

$$Pr[A \cap B] = Pr[A]Pr[B|A].$$

Consequently,

$$Pr[A \cap B \cap C] = Pr[(A \cap B) \cap C]$$

Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Hence,

$$Pr[A \cap B] = Pr[A]Pr[B|A].$$

Consequently,

$$\begin{aligned} Pr[A \cap B \cap C] &= Pr[(A \cap B) \cap C] \\ &= Pr[A \cap B]Pr[C|A \cap B] \end{aligned}$$

Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Hence,

$$Pr[A \cap B] = Pr[A]Pr[B|A].$$

Consequently,

$$\begin{aligned} Pr[A \cap B \cap C] &= Pr[(A \cap B) \cap C] \\ &= Pr[A \cap B]Pr[C|A \cap B] \\ &= Pr[A]Pr[B|A]Pr[C|A \cap B]. \end{aligned}$$

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof:

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof: By induction.

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof: By induction.

Assume the result is true for n .

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof: By induction.

Assume the result is true for n . (It holds for $n = 2$.)

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof: By induction.

Assume the result is true for n . (It holds for $n = 2$.) Then,

$$\begin{aligned} Pr[A_1 \cap \dots \cap A_n \cap A_{n+1}] \\ = Pr[A_1 \cap \dots \cap A_n]Pr[A_{n+1}|A_1 \cap \dots \cap A_n] \end{aligned}$$

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof: By induction.

Assume the result is true for n . (It holds for $n = 2$.) Then,

$$\begin{aligned} Pr[A_1 \cap \dots \cap A_n \cap A_{n+1}] \\ &= Pr[A_1 \cap \dots \cap A_n]Pr[A_{n+1}|A_1 \cap \dots \cap A_n] \\ &= Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]Pr[A_{n+1}|A_1 \cap \dots \cap A_n], \end{aligned}$$

Product Rule

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Proof: By induction.

Assume the result is true for n . (It holds for $n = 2$.) Then,

$$\begin{aligned} Pr[A_1 \cap \dots \cap A_n \cap A_{n+1}] \\ &= Pr[A_1 \cap \dots \cap A_n]Pr[A_{n+1}|A_1 \cap \dots \cap A_n] \\ &= Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]Pr[A_{n+1}|A_1 \cap \dots \cap A_n], \end{aligned}$$

so that the result holds for $n + 1$. □

Correlation

An example.

Correlation

An example.

Random experiment: Pick a person at random.

Correlation

An example.

Random experiment: Pick a person at random.

Event A : the person has lung cancer.

Correlation

An example.

Random experiment: Pick a person at random.

Event A : the person has lung cancer.

Event B : the person is a heavy smoker.

Correlation

An example.

Random experiment: Pick a person at random.

Event A : the person has lung cancer.

Event B : the person is a heavy smoker.

$$Pr[A|B] = 1.17 \times Pr[A].$$

Correlation

An example.

Random experiment: Pick a person at random.

Event A : the person has lung cancer.

Event B : the person is a heavy smoker.

$$Pr[A|B] = 1.17 \times Pr[A].$$

Conclusion:

- ▶ Smoking increases the probability of lung cancer by 17%.

Correlation

An example.

Random experiment: Pick a person at random.

Event A : the person has lung cancer.

Event B : the person is a heavy smoker.

$$Pr[A|B] = 1.17 \times Pr[A].$$

Conclusion:

- ▶ Smoking increases the probability of lung cancer by 17%.
- ▶ Smoking causes lung cancer.

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Note that

$$Pr[A|B] = 1.17 \times Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A]$$

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A]Pr[B] \end{aligned}$$

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A]Pr[B] \\ &\Leftrightarrow Pr[B|A] = 1.17 \times Pr[B]. \end{aligned}$$

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A]Pr[B] \\ &\Leftrightarrow Pr[B|A] = 1.17 \times Pr[B]. \end{aligned}$$

Conclusion:

- ▶ Lung cancer increases the probability of smoking by 17%.

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A]Pr[B] \\ &\Leftrightarrow Pr[B|A] = 1.17 \times Pr[B]. \end{aligned}$$

Conclusion:

- ▶ Lung cancer increases the probability of smoking by 17%.
- ▶ Lung cancer causes smoking.

Correlation

Event A : the person has lung cancer. Event B : the person is a heavy smoker. $Pr[A|B] = 1.17 \times Pr[A]$.

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A]Pr[B] \\ &\Leftrightarrow Pr[B|A] = 1.17 \times Pr[B]. \end{aligned}$$

Conclusion:

- ▶ Lung cancer increases the probability of smoking by 17%.
- ▶ Lung cancer causes smoking. Really?

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Other examples:

- ▶ Tesla owners are more likely to be rich.

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.
- ▶ People who go to the opera are more likely to have a good career.

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.
- ▶ People who go to the opera are more likely to have a good career. That does not mean that going to the opera will improve your career.

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.
- ▶ People who go to the opera are more likely to have a good career. That does not mean that going to the opera will improve your career.
- ▶ Rabbits eat more carrots and do not wear glasses.

Causality vs. Correlation

Events A and B are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

A and B being positively correlated does not mean that A causes B or that B causes A .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.
- ▶ People who go to the opera are more likely to have a good career. That does not mean that going to the opera will improve your career.
- ▶ Rabbits eat more carrots and do not wear glasses. Are carrots good for eyesight?

Proving Causality

Proving causality is generally difficult.

Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

- ▶ A and B may be positively correlated because they have a common cause.

Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

- ▶ A and B may be positively correlated because they have a common cause. (E.g., being a rabbit.)

Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

- ▶ A and B may be positively correlated because they have a common cause. (E.g., being a rabbit.)
- ▶ If B precedes A , then B is more likely to be the cause. (E.g., smoking.)

Proving Causality

Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

- ▶ A and B may be positively correlated because they have a common cause. (E.g., being a rabbit.)
- ▶ If B precedes A , then B is more likely to be the cause. (E.g., smoking.) However, they could have a common cause that induces B before A .

Proving Causality


Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

- ▶ A and B may be positively correlated because they have a common cause. (E.g., being a rabbit.)
- ▶ If B precedes A , then B is more likely to be the cause. (E.g., smoking.) However, they could have a common cause that induces B before A . (E.g., smart, CS70, Tesla.)


Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .



Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .




Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .




Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Indeed, B is the union of the disjoint sets $A_n \cap B$ for $n = 1, \dots, N$.

Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .



Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$


Indeed, B is the union of the disjoint sets $A_n \cap B$ for $n = 1, \dots, N$.

Thus,

$$Pr[B] = Pr[A_1]Pr[B|A_1] + \dots + Pr[A_N]Pr[B|A_N].$$

Total probability

Assume that Ω is the union of the disjoint sets A_1, \dots, A_N .



$$Pr[B] = Pr[A_1]Pr[B|A_1] + \dots + Pr[A_N]Pr[B|A_N].$$

Independence

Definition: Two events A and B are **independent** if

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent;

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent;
- ▶ When flipping coins, $A = \text{coin 1 yields heads}$ and $B = \text{coin 2 yields tails}$ are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent;
- ▶ When flipping coins, $A = \text{coin 1 yields heads}$ and $B = \text{coin 2 yields tails}$ are independent;

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent;
- ▶ When flipping coins, $A = \text{coin 1 yields heads}$ and $B = \text{coin 2 yields tails}$ are independent;
- ▶ When throwing 3 balls into 3 bins, $A = \text{bin 1 is empty}$ and $B = \text{bin 2 is empty}$ are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent;
- ▶ When flipping coins, $A = \text{coin 1 yields heads}$ and $B = \text{coin 2 yields tails}$ are independent;
- ▶ When throwing 3 balls into 3 bins, $A = \text{bin 1 is empty}$ and $B = \text{bin 2 is empty}$ are **not** independent;

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed:

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed: $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$, so that

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed: $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$, so that

$$Pr[A|B] = Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = Pr[A]$$

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed: $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$, so that

$$Pr[A|B] = Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = Pr[A] \Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B].$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair',

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] =$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2, P[B|\bar{A}] =$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$,

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] =$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A = \text{'coin is fair'}$, $B = \text{'outcome is heads'}$

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$Pr[B] = Pr[A \cap B] + Pr[\bar{A} \cap B] =$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$Pr[B] = Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$\begin{aligned} Pr[B] &= Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}] \\ &= (1/2)(1/2) + (1/2)0.6 = 0.55. \end{aligned}$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$\begin{aligned} Pr[B] &= Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}] \\ &= (1/2)(1/2) + (1/2)0.6 = 0.55. \end{aligned}$$

Thus,


$$Pr[A|B] = \frac{Pr[A]Pr[B|A]}{Pr[B]} = \frac{(1/2)(1/2)}{(1/2)(1/2) + (1/2)0.6} \approx 0.45.$$

Is your coin loaded?

A picture:


Is your coin loaded?

A picture:



Is your coin loaded?

A picture:




Imagine 100 situations, among which

$m := 100(1/2)(1/2)$ are such that A and B occur and

$n := 100(1/2)(0.6)$ are such that \bar{A} and B occur.

Is your coin loaded?

A picture:



Imagine 100 situations, among which


$m := 100(1/2)(1/2)$ are such that A and B occur and

$n := 100(1/2)(0.6)$ are such that \bar{A} and B occur.

Thus, among the $m + n$ situations where B occurred, there are m where A occurred.

Is your coin loaded?

A picture:



Imagine 100 situations, among which

$m := 100(1/2)(1/2)$ are such that A and B occur and


$n := 100(1/2)(0.6)$ are such that \bar{A} and B occur.

Thus, among the $m + n$ situations where B occurred, there are m where A occurred.


Hence,

$$Pr[A|B] = \frac{m}{m+n} = \frac{(1/2)(1/2)}{(1/2)(1/2) + (1/2)0.6}.$$


Why do you have a fever?



Why do you have a fever?




Why do you have a fever?



Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

Why do you have a fever?




Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

Why do you have a fever?




Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

Why do you have a fever?



Using Bayes' rule, we find


$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

These are the **posterior probabilities**.

Why do you have a fever?



Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$


$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$


These are the **posterior probabilities**. One says that 'Flu' is the **Most Likely a Posteriori** (MAP) cause of the high fever.

Bayes' Rule Operations

Bayes' Rule Operations



Bayes' Rule Operations



Bayes' Rule is the canonical example of how information changes our opinions.

Thomas Bayes

Thomas Bayes




Portrait used of Bayes in a 1936 book,^[1] but it is doubtful whether the portrait is actually of him.^[2]

No earlier portrait or claimed portrait survives.

Born	c. 1701 London, England
Died	7 April 1761 (aged 59) Tunbridge Wells, Kent, England
Residence	Tunbridge Wells, Kent, England
Nationality	English
Known for	Bayes' theorem

Thomas Bayes



A Bayesian picture of Thomas Bayes.

Testing for disease.

Let's watch TV!!

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test*, *disease*)

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test*, *disease*)

A - prostate cancer.

B - positive PSA test.

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.

- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.

- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

From http://www.cpcn.org/01_psa_tests.htm and

<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.

- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

From http://www.cpcn.org/01_psa_tests.htm and
<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)

Positive PSA test (*B*). Do I have disease?

Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.


- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

From http://www.cpcn.org/01_psa_tests.htm and
<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)


Positive PSA test (*B*). Do I have disease?

$$Pr[A|B]???$$

Bayes Rule.




Bayes Rule.



Using Bayes' rule, we find


Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10}$$


Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

Bayes Rule.




Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test.

Bayes Rule.




Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test. !!!!

Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test. !!!!
!!!!

Monty Hall.

Summary

Events, Conditional Probability, Independence, Bayes' Rule

Summary

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

Summary

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.

Summary

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.
- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

Summary

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.
- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

$Pr[A_n|B]$ = posterior probability; $Pr[A_n]$ = prior probability .

Summary

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.
- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

$Pr[A_n|B]$ = posterior probability; $Pr[A_n]$ = prior probability .

- ▶ All these are possible:

$$Pr[A|B] < Pr[A]; Pr[A|B] > Pr[A]; Pr[A|B] = Pr[A].$$