

Chapter 4 Linear Models

LECTURER: ZHIHUA JIANG

A solid green horizontal bar spanning the width of the slide at the bottom.

Content

3.1 Linear model

3.2 Linear regression

3.3 Logistic regression


3.4 Linear discriminant analysis (LDA)

3.1 Linear model

- Linear model: linear function of attributes

$$f(\mathbf{x}) = w_1x_1 + w_2x_2 + \dots + w_dx_d + b$$

- Matrix form: $f(\mathbf{x}) = \mathbf{w}^T \mathbf{x} + b$


$$\beta = (\mathbf{w}; b), \quad \hat{\mathbf{x}} = (\mathbf{x}; 1)$$
$$\mathbf{w}^T \mathbf{x} + b \longrightarrow \beta^T \hat{\mathbf{x}}$$

- Advantages:
simple model; basic model; good interpretability

3.2 Linear regression

- **Univariate linear regression**

$$f(x_i) = wx_i + b \text{ such that } f(x_i) \approx y_i$$

Where x_i is a scalar

- **Multivariate linear regression**

$$f(\mathbf{x}_i) = \mathbf{w}^T \mathbf{x}_i + b \text{ such that } f(\mathbf{x}_i) \approx y_i$$

Where \mathbf{x}_i is a vector

- **Generalized linear model**

$$y = g^{-1}(\mathbf{w}^T \mathbf{x} + b)$$

Where $g(\cdot)$ is a monotone differentiable function

Univariate linear regression

- How to determine w and b ?
 - a) To minimize the *MSE*

$$\begin{aligned}(w^*, b^*) &= \arg \min_{(w, b)} \sum_{i=1}^m (f(x_i) - y_i)^2 \\ &= \arg \min_{(w, b)} \sum_{i=1}^m (y_i - wx_i - b)^2 .\end{aligned}$$

Univariate linear regression

- How to determine w and b ? (cont.)

b) Parameter estimation based on least square method

$$E_{(w,b)} = \sum_{i=1}^m (y_i - wx_i - b)^2$$

- Differentiate with w and b respectively

$$\frac{\partial E_{(w,b)}}{\partial w} = 2 \left(w \sum_{i=1}^m x_i^2 - \sum_{i=1}^m (y_i - b) x_i \right) ,$$

$$\frac{\partial E_{(w,b)}}{\partial b} = 2 \left(mb - \sum_{i=1}^m (y_i - wx_i) \right) ,$$



$$w = \frac{\sum_{i=1}^m y_i (x_i - \bar{x})}{\sum_{i=1}^m x_i^2 - \frac{1}{m} \left(\sum_{i=1}^m x_i \right)^2} ,$$
$$b = \frac{1}{m} \sum_{i=1}^m (y_i - wx_i)$$

Multivariate linear regression

- Given $D = \{(x_1, y_1), \dots, (x_m, y_m)\}$,

$x_i = (x_{i1}; \dots; x_{id}), y_i \in \mathbb{R} \rightarrow m \times (d+1)$ matrix X

$$\hat{w} = (w; b)$$

$$X = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1d} & 1 \\ x_{21} & x_{22} & \dots & x_{2d} & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ x_{m1} & x_{m2} & \dots & x_{md} & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{x}_1^T & 1 \\ \mathbf{x}_2^T & 1 \\ \vdots & \vdots \\ \mathbf{x}_m^T & 1 \end{pmatrix}$$

$$\mathbf{y} = (y_1; y_2; \dots; y_m)$$

Multivariate linear regression

- Similarly, to minimize the MSE

$$\hat{\mathbf{w}}^* = \arg \min_{\hat{\mathbf{w}}} (\mathbf{y} - \mathbf{X}\hat{\mathbf{w}})^T (\mathbf{y} - \mathbf{X}\hat{\mathbf{w}})$$

$$E_{\hat{\mathbf{w}}} = (\mathbf{y} - \mathbf{X}\hat{\mathbf{w}})^T (\mathbf{y} - \mathbf{X}\hat{\mathbf{w}})$$

- Differentiate with $\hat{\mathbf{w}}$

$$\frac{\partial E_{\hat{\mathbf{w}}}}{\partial \hat{\mathbf{w}}} = 2 \mathbf{X}^T (\mathbf{X}\hat{\mathbf{w}} - \mathbf{y}) \quad \begin{array}{c} \text{full-rank matrix, or} \\ \text{positive definite matrix} \end{array} \quad \hat{\mathbf{w}}^* = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$

Generalized linear model

- Problem: how to let the linear prediction to approximate some function of real labels?


$$y = g^{-1}(w^T x + b)$$

Where $g(\cdot)$ is the link function (*monotone & differentiable*),

$g^{-1}(\cdot)$ is the inverse function

- Example: $\ln y = w^T x + b$ \Rightarrow $y = e^{w^T x + b}$ (对数线性回归)

Example: log-linear regression




3.3 *logistic regression*

- Problem: use the linear regression for the classification problem?
- Solution: *Generalized linear model*
- Candidates of $g(\cdot)$:
 - a) Unit-step function (单位阶跃函数)
 - b) Logistic function (对数几率函数)

Class label

Unit-step
function

Logistic function



Logistic function

- Odds(几率)

y : probability of x being a positive example

$1-y$: probability of x being a negative example

$y/(1-y)$: relative probability of x being a positive example

- Logistic function: any order differentiable

mathematical function

$$y = \frac{1}{1 + e^{-z}} \xrightarrow{\text{ties}} y = \frac{1}{1 + e^{-(\mathbf{w}^T \mathbf{x} + b)}} \xrightarrow{\text{mathematical}} \ln \frac{y}{1-y} = \mathbf{w}^T \mathbf{x} + b$$

Logistic function

- Posterior probability estimation

$$\ln \frac{y}{1-y} = \mathbf{w}^T \mathbf{x} + b \quad \Rightarrow \quad \ln \frac{p(y=1 | \mathbf{x})}{p(y=0 | \mathbf{x})} = \mathbf{w}^T \mathbf{x} + b$$



$$p(y=1 | \mathbf{x}) = \frac{e^{\mathbf{w}^T \mathbf{x} + b}}{1 + e^{\mathbf{w}^T \mathbf{x} + b}},$$

$$p(y=0 | \mathbf{x}) = \frac{1}{1 + e^{\mathbf{w}^T \mathbf{x} + b}}.$$

maximum likelihood method



$$\prod_{i=1}^m p(y_i | x_i; \mathbf{w}, b)$$

Logistic function

- Log likelihood function:

$$\ell(\mathbf{w}, b) = \sum_{i=1}^m \ln p(y_i \mid \mathbf{x}_i; \mathbf{w}, b)$$



$$\boldsymbol{\beta} = (\mathbf{w}; b), \quad \hat{\mathbf{x}} = (\mathbf{x}; 1)$$

$$\mathbf{w}^T \mathbf{x} + b \longrightarrow \boldsymbol{\beta}^T \hat{\mathbf{x}}$$

$$p_1(\hat{\mathbf{x}}; \boldsymbol{\beta}) = p(y = 1 \mid \hat{\mathbf{x}}; \boldsymbol{\beta})$$

$$p_0(\hat{\mathbf{x}}; \boldsymbol{\beta}) = p(y = 0 \mid \hat{\mathbf{x}}; \boldsymbol{\beta}) = 1 - p_1(\hat{\mathbf{x}}; \boldsymbol{\beta})$$

$$p(y_i \mid \mathbf{x}_i; \mathbf{w}, b) = y_i p_1(\hat{\mathbf{x}}_i; \boldsymbol{\beta}) + (1 - y_i) p_0(\hat{\mathbf{x}}_i; \boldsymbol{\beta})$$

Logistic function

- Likelihood function (cont.)

$$p(y_i | \mathbf{x}_i; \mathbf{w}, b) = y_i p_1(\hat{\mathbf{x}}_i; \boldsymbol{\beta}) + (1 - y_i) p_0(\hat{\mathbf{x}}_i; \boldsymbol{\beta})$$

$$\mathbf{w}^T \mathbf{x} + b \longrightarrow \boldsymbol{\beta}^T \hat{\mathbf{x}}$$

$$\text{Max } l(\mathbf{w}; b) \leftrightarrow \text{Min } l(\boldsymbol{\beta})$$

$$\ell(\boldsymbol{\beta}) = \sum_{i=1}^m \left(-y_i \boldsymbol{\beta}^T \hat{\mathbf{x}}_i + \ln \left(1 + e^{\boldsymbol{\beta}^T \hat{\mathbf{x}}_i} \right) \right)$$

Newton method (update for the $t+1$ iteration)

$$\boldsymbol{\beta}^{t+1} = \boldsymbol{\beta}^t - \left(\frac{\partial^2 \ell(\boldsymbol{\beta})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^T} \right)^{-1} \frac{\partial \ell(\boldsymbol{\beta})}{\partial \boldsymbol{\beta}}$$

$$\ell(\beta) = \sum_{i=1}^m \left(-y_i \beta^T \hat{x}_i + \ln(1 + e^{\beta^T \hat{x}_i}) \right)$$

Max $l(w, b) \leftrightarrow$ Min $l(\beta)$

$$l(w, b) = \sum_{i=1}^m \ln p(y_i | x_i; w, b)$$

$$= \sum_{i=1}^m \ln [y_i p_1(\hat{x}_i; \beta) + (1 - y_i) p_0(\hat{x}_i; \beta)]$$

$$= \sum_{i=1}^m \ln \left[y_i \frac{e^{\beta^T \hat{x}_i}}{1 + e^{\beta^T \hat{x}_i}} + (1 - y_i) \frac{1}{1 + e^{\beta^T \hat{x}_i}} \right]$$

$$= \sum_{i=1}^m \ln \frac{y_i e^{\beta^T \hat{x}_i} + (1 - y_i)}{1 + e^{\beta^T \hat{x}_i}}$$


$$= \begin{cases} y_i = 1, & \sum_{i=1}^m \ln \frac{e^{\beta^T \hat{x}_i}}{1 + e^{\beta^T \hat{x}_i}} = \sum_{i=1}^m (\beta^T \hat{x}_i - \ln(1 + e^{\beta^T \hat{x}_i})) \\ y_i = 0, & \sum_{i=1}^m \ln \frac{1}{1 + e^{\beta^T \hat{x}_i}} = \sum_{i=1}^m (-\ln(1 + e^{\beta^T \hat{x}_i})) \end{cases}$$

3.4 Linear discriminant analysis (LDA)

- Idea:

- Cast the samples onto a straight line
- Project the similar samples as close as possible
- Project the dissimilar samples as far as possible
- For a new sample, determine the class according to the relative position of its projection point.

3.4 Linear discriminant analysis (LDA)



Goal of LDA

- Given dataset $D = \{(x_i, y_i)\}_{i=1}^m, y_i \in \{0, 1\}$
 - X_i : sample set of the i^{th} class
 - μ_i : mean vector of the i^{th} class
 - Σ_i : covariance matrix of the i^{th} class
 - Projection points of the two centers in the line: $w^T \mu_0$ and $w^T \mu_1$
 - Covariance: $w^T \Sigma_0 w$ and $w^T \Sigma_1 w$
- Similar samples as close as possible $\rightarrow w^T \Sigma_0 w + w^T \Sigma_1 w$ as small as possible
- Dissimilar samples as far as possible \rightarrow as large as possible

$$\|w^T \mu_0 - w^T \mu_1\|_2^2$$

$$\|x\|_2 := \sqrt{x_1^2 + \cdots + x_n^2}.$$

If the entries in the column vector

$$\mathbf{X} = \begin{bmatrix} X_1 \\ \vdots \\ X_n \end{bmatrix}$$

are random variables, each with finite variance, then the covariance matrix Σ is the matrix whose (i, j) entry is the covariance

$$\Sigma_{ij} = \text{cov}(X_i, X_j) = E[(X_i - \mu_i)(X_j - \mu_j)] = E[X_i X_j] - \mu_i \mu_j,$$

where the operator E denotes the expected (mean) value of its argument, and

$$\mu_i = E(X_i)$$

is the expected value of the i -th entry in the vector \mathbf{X} . In other words,

$$\Sigma = \begin{bmatrix} E[(X_1 - \mu_1)(X_1 - \mu_1)] & E[(X_1 - \mu_1)(X_2 - \mu_2)] & \cdots & E[(X_1 - \mu_1)(X_n - \mu_n)] \\ E[(X_2 - \mu_2)(X_1 - \mu_1)] & E[(X_2 - \mu_2)(X_2 - \mu_2)] & \cdots & E[(X_2 - \mu_2)(X_n - \mu_n)] \\ \vdots & \vdots & \ddots & \vdots \\ E[(X_n - \mu_n)(X_1 - \mu_1)] & E[(X_n - \mu_n)(X_2 - \mu_2)] & \cdots & E[(X_n - \mu_n)(X_n - \mu_n)] \end{bmatrix}.$$

3.4 Goal of LDA

- Goal: maximize

$$J = \frac{\|w^T \mu_0 - w^T \mu_1\|_2^2}{w^T \Sigma_0 w + w^T \Sigma_1 w}$$
$$= \frac{w^T (\mu_0 - \mu_1) (\mu_0 - \mu_1)^T w}{w^T (\Sigma_0 + \Sigma_1) w}$$



$$J = \frac{w^T S_b w}{w^T S_w w}$$

- Within-class scatter matrix

$$S_w = \Sigma_0 + \Sigma_1$$
$$= \sum_{x \in X_0} (x - \mu_0) (x - \mu_0)^T + \sum_{x \in X_1} (x - \mu_1) (x - \mu_1)^T$$

- Between-class scatter matrix

$$S_b = (\mu_0 - \mu_1) (\mu_0 - \mu_1)^T$$

3.4 Goal of LDA

- How to solve w ?

$$L(w, \lambda) = -w^T S_b w + \lambda(w^T S_w w - 1)$$

$$\frac{\partial L}{\partial w} = -2S_b w + 2\lambda S_w w = 0 \Rightarrow S_b w = \lambda S_w w$$

Lagrange
multipliers

$$J = \frac{w^T S_b w}{w^T S_w w} \xrightarrow{\text{set } w^T S_w w = 1} \min_w -w^T S_b w \xrightarrow{\text{Lagrange multipliers}} S_b w = \lambda S_w w$$

s.t. $w^T S_w w = 1$

[推导]: 由公式 (3.36) 可得拉格朗日函数为

$$L(w, \lambda) = -w^T S_b w + \lambda(w^T S_w w - 1)$$

对 w 求偏导可得

$$\begin{aligned}\frac{\partial L(w, \lambda)}{\partial w} &= -\frac{\partial(w^T S_b w)}{\partial w} + \lambda \frac{\partial(w^T S_w w - 1)}{\partial w} \\ &= -(S_b + S_b^T)w + \lambda(S_w + S_w^T)w\end{aligned}$$

由于 $S_b = S_b^T, S_w = S_w^T$, 所以

$$\frac{\partial L(w, \lambda)}{\partial w} = -2S_b w + 2\lambda S_w w$$

令上式等于 0 即可得

$$-2S_b w + 2\lambda S_w w = 0$$

$$S_b w = \lambda S_w w$$

由于我们要求解的只有 w , 而 λ 这个拉格朗乘子具体取值多少都无所谓, 因此我们可以任意设定 λ 来配合我们求解 w 。我们注意到

$$S_b w = (\mu_0 - \mu_1)(\mu_0 - \mu_1)^T w$$

如果我们令 λ 恒等于 $(\mu_0 - \mu_1)^T w$, 那么上式即可改写为

$$S_b w = \lambda(\mu_0 - \mu_1)$$

将其代入 $S_b w = \lambda S_w w$ 即可解得

$$w = S_w^{-1}(\mu_0 - \mu_1)$$


Lagrange multipliers (拉格朗日乘子法)

- Idea:

d variables, k constraints $\rightarrow d+k$ variables, 0 constraint

- Goal:

\mathbf{x} is d -vector, minimize $f(\mathbf{x})$ s.t. $g(\mathbf{x})=0$

- $L(\mathbf{x}, \lambda) = f(\mathbf{x}) + \lambda g(\mathbf{x})$, n :  $\left\{ \begin{array}{l} \nabla f(\mathbf{x}^*) + \lambda \nabla g(\mathbf{x}^*) = 0 \\ g(\mathbf{x}) = 0. \end{array} \right.$
Set partial derivative to be 0