

GAIA-Q: A Quantum-Coherent Atmospheric Framework for Planetary-Scale Information Processing

Harshvinder Singh

Mtech Data Science and Analytics.

Lovely Professional University

Phagwara, India

singh.harshvinder2000@gmail.com

Abstract—The Gaia hypothesis has long been used to define the earth as an auto-regulated bio-geophysical system where various atmospheric, ecological, and chemical interactions interact. New discoveries in the fields of quantum biology, the atmospheric electrodynamics, and the complex adaptive systems encourage a more in-depth investigation whether these interactions are found to possess characteristics that are consistent with large-scale information processing. In this paper, a scientifically based framework, GAIA-Q, is presented that represents Earth as a quantum-coherent and cyber-physical information architecture based on an atmosphere-biosphere continuum. The model proposed combines quantum scattering coherence, Schumann resonance dynamics, ionospheric charge transport and aerosol feedback (biosphere mediated) to examine the phenomenon of a structure flow of information at the planetary scale. On the basis of Maxwell field theory mathematical formulations, entropy-based measures of information, and coherence-length analysis, GAIA-Q defines the multi-layer interactions that produce the global sensing, low-frequency electromagnetic communication, and adaptive climatic regulation. Theoretical modeling has shown that the coupled Earth system exhibits non-trivial information storage and signal transmission as well as feedback-based modulation similar to distributed computational intelligence. Although empirical validation needs substantial atmospheric measurement and simulation, GAIA-Q provides the pioneer integrated physical system of processing information at planetary scale and provides architectural information on how to create strong and self-organizing artificial general intelligence systems based on what nature knows with regard to regulating its own processes.

Index Terms—Biogenic Aerosol, feedback Atmospheric Electrodynamics, Cyber-Physical Systems, Gaia Hypothesis, Ionospheric Charge Transport, Planetary-Scale Information Processing, Quantum Biology, Quantum Coherence, Schumann Resonances, Self-organizing Systems.

I. INTRODUCTION

The classical Gaia hypothesis, proposed by Lovelock and Margulis, conceptualizes Earth as a self-regulating system in which atmospheric, biological, geochemical, and hydrological processes co-evolve to maintain planetary stability. While traditionally focused on ecological and biogeochemical regulation, recent advances in quantum biology, atmospheric electrodynamics, and complex adaptive systems suggest that Earth may also function as a large-scale information-

processing network. Long-lived quantum coherence observed in photosynthetic complexes and avian magnetoreception indicates that quantum effects persist in warm, noisy environments. Concurrently, the Earth-ionosphere cavity sustains coherent Schumann resonances that act as global electromagnetic waveguides. Biosphere-driven atmospheric processes—including aerosol formation, cloud microphysics, and microbial and vegetation feedback—further contribute to structured regulatory dynamics. Interpreted through information theory and cybernetics, these multi-scale phenomena resemble sensing, encoding, transmission, and feedback operations. This motivates the GAIA-Q framework, which integrates quantum coherence, electromagnetic resonance, entropy flow, and ecological regulation to model Earth as a distributed cyber-physical information architecture.

II. BACKGROUND AND RELATED WORK:

The GAIA-Q model is based on interdisciplinary evidence of quantum biology, atmospheric electrodynamics, biosphere-climate feedbacks as well as complex systems theory. It has been demonstrated that quantum coherence can also survive in the other microscopic systems at physiological temperatures, such as in Engel in photosynthetic complexes and by Gauger in avian magneto-reception, so it is clear that complex systems in nature can maintain a structured phase information. Schumann resonances at the planetary level offer unperturbed electromagnetic modes of the Earth ionosphere cavity as Nickolaenko, Hayakawa, and Rycroft define that globally stable ELF propagation is observable. It has been experimentally determined by biosphere-atmosphere coupling research, such as that of Carslaw and Volk, that biogenic aerosols, evapotranspiration and microbial emissions are capable of having quantifiable regulatory impact on cloud microphysics and climate feedback. Complex adaptive systems theory also describes the scaling of local interactions to global behavior, and the Gaia hypothesis and the new concepts of planetary intelligence encourage the view of coupled systems on Earth as a globally distributed information-processing architecture. All these early works contribute towards GAIA-Q as a physically based model that

is quantum coherent, electromagnetic resonant and biospheric feedback to explain how information flows at the planetary scale.

TABLE I
REVIEW OF SELECTED LITERATURE RELEVANT TO GAIA-Q: KEY CONTRIBUTIONS AND LIMITATIONS.

Reference	Year	Main Contribution	Limitation
Engel et al.	2007	Evidence of long-lived quantum coherence in photosynthesis.	Limited to molecular scales; no evidence of atmospheric-scale coherence.
Gauger et al.	2011	Coherent radical-pair magnetoreception mechanism in birds.	Biological scale only; not applicable to geophysical media.
Nickolaenko & Hayakawa	2014	Detailed characterization of Schumann resonance modes and Q-factors.	Descriptive only; lacks information-theoretic interpretation.
Rycroft et al.	2000	Global electric circuit linked to solar/ionospheric variability.	Phenomenological; no quantitative information-capacity modeling.
Lovelock & Margulis	1974	Proposed Gaia hypothesis of biospheric regulation.	Conceptual; lacks measurable physical mechanisms.
Volk	1987	Geochemical feedbacks affecting atmospheric CO ₂ .	Geological scale only; not linked to dynamic information flow.
Carslaw et al.	2018	Biogenic aerosols modulate cloud microphysics.	Does not consider aerosols as information carriers.
Frank & Walker	2022	Introduced planetary intelligence concept.	Conceptual; lacks physical coherence mechanisms.
Holland	1992	Defined complex adaptive systems theory.	Abstract; not linked to atmospheric/quantum processes.

III. PROPOSED GAIA-Q FRAMEWORK

The hypothesis illustrated by the GAIA-Q framework is that the multi-layered planetary information architecture of the coupled atmosphere-biosphere-ionosphere system is formalized. It is a quantum level architecture that incorporates quantum level interactions, electromagnetic resonances, atmospheric communication channels, and biosphere based regulatory processes. Unlike the classical Gaia theory, which thinks of the Earth as a self-regulating ecosystem, GAIA-Q presents a cyber-physical view, and the subsystems of the Earth are presented in the context of information flow, coherence, signal propagation, and control feedback.

The suggested framework is composed of four layers allied to each other and each having a specific group of physical processes. Together the layers create a hierarchy of information-processing in the quantum dynamical scales, mesoscale dynamical scales and planetary dynamical scales.

A. Layer 1: Photonic interaction and quantum Coherence Layer

The atmosphere is modeled by the GAIA-Q as the first layer in which a structured phase information can be maintained due to coherent scattering, wave-particle and ionization dynamics. Atmospheric photons generated from solar radiation undergo Rayleigh, Raman, and Mie scattering, potentially maintaining coherence over mesoscale distances depending on photon wavelength λ and spectral width $\Delta\lambda$. The coherence length is defined as:

$$L_c = \frac{\lambda^2}{\Delta\lambda} \quad (1)$$

The layer explores the possible existence of long-lived coherences phenomena to aid information encoding through the form of interference pattern, coherence modulation or regular spectral signature. Along with that, ionospheric variations in charge and plasma oscillations are also modeled using quantum-like dynamical states and comprise the lowest-level information substrate of GAIA-Q.

B. Layer 2: Planetary Wave and Vibrations and Electricity Electromagnetic Resonance.

The ionosphere and surface of the earth make a natural electro. where the Schumann resonances are maintained due to some cavity, which is magnetic, provided:

$$f_n = \frac{c}{2\pi R} \sqrt{n(n+1)} \quad (2)$$

where R denotes the radius of the earth and n represents the resonance. These are standing waves that transmit ELF/VLF throughout the whole world with little attenuation, permitting dashboard interaction between atmospheric regions. The Schumann resonance system is effective in GAIA-Q the world communication infrastructure, which is similar to the world-planetary scale bus architecture in the distributed computing. Resonance quality, modal stability and amplitude variations are useful as measures of the information carrying capacity in the cavity. This is because any disturbance in resonance spectra stores information about any alterations in global ionospheric conductivity, atmospheric, and lightning activity dynamics.

C. Layer 3: Entropy- and Atmospheric Communication and Regulated Transport Layer

This layer defines the dynamics of the atmosphere- including turbulence, gradients of ion density, waves propagation, and aerosol transport as agents of low-frequency information. The content of atmospheric information is given using Shannon entropy. is defined as:

$$S(t) = - \sum_i p_i(t) \log p_i(t) \quad (3)$$

The atmosphere thus is a thermodynamic channel of communication which has the ability of modulating information density due to fluctuation of pressure, humidity, chemical

composition, aerosol concentration, Cloud microphysics, turbulence cascades, and mesoscale teleconnections as they are being of use to operate on signals, signal transformation operators allowing environmental en- filtering, redistributing and coding information.

D. Layer 4: Cybernetic Feedback and Biospheric Control Layer

The biosphere impacts the atmospheric condition by evapo-volatile organic compound (VOC) emission, transpiration. microbial cloud seeding, biogenic aerosol release, and biogenic aerosol release these processes serve as cybernetic feedback regulators which produce organized changes in the composition of the atmosphere and thermodynamics. The functions of feedback are modeled as:

$$B(x, t) = \alpha_1 F_{\text{aerosol}} + \alpha_2 F_{\text{evap}} + \alpha_3 F_{\text{biogenic}} \quad (4)$$

in which the coefficients α_i are the measures of biospheric impact. on atmospheric processes. This layer offers sensing (e.g. biological response to radiation), actuation (e.g., moisture release), and control (e.g. cloud albedo modulation), and establish the control system of the GAIA-Q.

E. Cross- Layer Interactions and Hierarchy of Information.

Processing instead of being independently coupled, the four layers are dynamically coupled. For example:

- Biospheric emissions regulate aerosol densities that change scattering patterns of Layer 1.
- Atmospheric teleconnections modify earth ionosphere cavity modify Schumann resonance in it geometry of Layer 2.
- Electro-quantum level coherence perturbations varying magnetic field, biospheric feedback in Layer 4.

These contacts are a distributed cyber-physical global senses, signal propagation in information network and responsive control facilities.

F. Planetary-Scale Architecture of Information Surveillance.

TABLE II
GAIA-Q INFORMATION ARCHITECTURE SUMMARY

Layer	Function	Information Role
Quantum Layer	Photon scattering, coherence	Signal encoding, phase information
EM Resonance Layer	Schumann resonances	Planetary communication backbone
Atmospheric Layer	Teleconnections, aerosols	Transport, modulation, filtering
Biospheric Layer	Climate feedback, emissions	Sensing, actuation, control

Collectively, these layers will make GAIA-Q a planetary-scale ,cyber-physical intelligence architecture and multi-layer.

IV. METHODOLOGY

GAIA-Q is based on the methodological basis. a multi-level analysis system that incorporates quantum. optical theory, atmospheric electrodynamics, thermodynamic. cybernetic biosphereclimate feedback, entropy analysis and cybersystems. modeling. The Earth system is highly nonlinear and hence, this has been the case. hierarchy and cross-scale coupling, a hierarchical modeling approach. is used to de-contextualize, formalize and quantify information. interactions between and inside every subsystem. The methodology has five major elements: subsystem decomposi- tion, mathematical formulation, cross-layer coupling bestowal, theoretical simulation, information-processing validation.

A. A subsystem Decomposition and Formal Layer Architecture.

The Earth system is decomposed into Four theoretically, the earth system is broken down into:

- **independent and dynamically coupled::** controls the coherence of the photons, ionospheric charge oscillation, optical scattering and. wave-particle interactions.
- **Earth Magnetic Resonance Layer:** controlled by the Earth- Schumann resonance and ionosphere cavity dynamics ionosphere cavity dynamics and Schumann resonance modal stability.
- **Atmospheric Information Transport Layer:**This layer serves the upper-most layer, which provides information services on the atmosphere. statistics of turbulence gradients thermodynamically diffused, and information encoding on the basis of entropy.
- **Biosphere Cybernetic Feedback Layer::** consisting of evapotranspiration VOC emissions- aerosol microphysics, regulation of cloud albedo, and ecological sensing. mechanisms.

The different layers are mathematically modeled as dynam- icals. subsystem having quantifiable state variables, forcing terms, and transfer functions. This degradation enables corporeal. mechanisms that are to be studied separately and still maintain the. multi-scale interaction necessary to emerge on a global scale.

B. Mathematical Representations of Information about the Earth-System. Pathways

1) *Quantum Layer Formulation:* Quantum-level interactions are modeled using coherence length theory:

$$L_c = \frac{\lambda^2}{\Delta\lambda} \quad (5)$$

and density matrix evolution:

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[H, \rho] + \mathcal{L}(\rho) \quad (6)$$

where $\mathcal{L}$ represents atmospheric decoherence superopera- tors. These equations quantify the temporal and spatial persistence of coherent optical information in the lower and middle atmosphere.

2) *Electromagnetic Resonance Layer Formulation:* Schumann resonances are modeled by solving the spherical wave equation:

$$\nabla^2 E + \frac{\omega^2}{c^2} E = 0 \quad (7)$$

leading to resonance frequencies:

$$f_n = \frac{c}{2\pi R} \sqrt{n(n+1)} \quad (8)$$

A Q-factor analysis provides a quantitative estimate of information retention:

$$Q = 2\pi \frac{E_{\text{stored}}}{E_{\text{lost}}} \quad (9)$$

3) *Atmospheric Transport Layer Formulation:* Atmospheric information encoding is modeled using thermodynamic entropy:

$$S(t) = - \sum_i p_i(t) \ln p_i(t) \quad (10)$$

along with turbulence probability density functions and multi-scale entropy (MSE) analysis. These models capture the formation, deformation, and propagation of structured atmospheric information packets.

4) *Biospheric Feedback Layer Formulation:* Climate-biosphere control interactions are modeled using cybernetic feedback equations:

$$B(x, t) = \beta_1 A(x, t) + \beta_2 E(x, t) + \beta_3 V(x, t) \quad (11)$$

where A , E , and V represent aerosol emissions, evapotranspiration rates, and vapor rates. These variables become biological sensing functions that act as actuation functions that enforce structure state regulation on atmospheric information.

C. Modeling of Cross Layer Coupling and Interaction.

The formal coupling tensor, Γ_{ij} is built to reflect, directional inter-layer influence between layer i and j . The tensor incorporates:

- Radiative forcing transfer.
- Aerosol, Kernels Teleconnection propagation kernels
- Atmospheric and biosphere mutual information states.
- Here a coefficient of electromagnetic susceptibility

This offers a strict mathematical foundation of identification, that cohesiveness from interactions on a global scale, system stabilization, or dissolve informational structure, dynamics.

D. Theoretical Simulations and Analytical Appraisals.

The study was not able to measure the planets on a planetary scale because such measurements were not complete. uses conceptual models based on confirmed physical. models:

- The simulation of Schumann resonance mode propagation of. phase stability lifetimes and Global coherence lifetimes.

- Simulation of the atmospheric quantum coherence decay. interactions of photons with aerosol loading conditions of different levels.
- Simulation of entropy perturbation on humidity, aerosol, etc. and thermal anomalies.
- Simulations Biospheric simulations: feedback response simulations efficiency and stability margins of regulation.

Theoretical limits to the feasibility are obtained with these simulations. of information processing of planets.

E. Information Processing Approachs to validation.

In order to know whether the Earth system is satisfying the theoretical computational needs of large scale computation, multiple quantitative indicators are determined:

1) Coherence Indicators::

- Photon coherence length L_c
- Q-factor of electromagnetic resonance.
- The phase correlation of cross-layers.

2) Information Capacity Indicators:

- Modal bandwidth of Schumann resonances
- Atmospheric entropy gradient amplitude
- Information propagation length scales

3) Control-Theoretic Indicators:

- Stability margins of ecological-atmospheric loops
- Response times to radiative or chemical perturbations
- Gain dynamics of biospheric regulators

4) Emergent Intelligence Indicators:

- Global synchrony patterns
- Redundancy and fault tolerance
- Multi-scale adaptation
- Self-regulation under perturbation

These criteria determine whether GAIA-Q satisfies the necessary and sufficient conditions for planetary-scale information processing.

F. Methodology Limitations and Assumptions.

The analysis is limited by a number of assumptions:

- Quantum coherence of the atmosphere is not. zero but intermittent.
- Stability of Schumann resonance is assumed to be adequate to long-range coupling.
- Biospheric feedback is a cyber-feedback which is modeled to be measurable. netic regulatory system.
- It is information theory that is strictly applied in the physical. delimiting, other than metaphysical understandings.

V. RESULTS AND DISCUSSION

This part contains the theoretical findings obtained out of. the GAIA-Q framework and elaborates on their implications of the same to. knowledge of the atmosphere-biosphere- ionosphere of the Earth. tinuum distributed information-processing system. Since the current research is theoretical and simulation based, the conclusions. emphasize on analyses, derived measurement, cross-domain. correlations and properties of

theoretical coherence were proved. contrary to laws of physical nature and objective tendencies.

It was found through the study of atmospheric scattering that when atmospheric scattering occurs, it is under special spectral requirements, the length of coherence.

A. The Results of Quantum Coherence and Photonic Interaction Analysis

It was found through the study of atmospheric scattering that when atmospheric scattering occurs, it is under special spectral requirements, the length of coherence.

$$L_c = \frac{\lambda^2}{\Delta\lambda} \quad (12)$$

are capable of being bigger than mesoscale turbulent cells, especially duration- conditions of low aerosol, high visibility. This suggests that phase can tentatively be held in portions of the atmosphere. long enough to be involved in an organized photonic interactions.

Model simulations indicate:

- increased coherence retention at shorter wavelengths (UV-;—human—;Higher coherence retention at shorter wavelengths (UV- to -). blue band),
- A lower decoherence in low humidity areas of the upper troposphere,
- possible reinforcement of coherence by repetition —human—;Concrete reinforcement: toring events.

Although coherence does not exist over international levels, its. episodic preservation provides a medium of localized quantum- like information encoding. These results facilitate the suit- quantum layer ability to initialize information as the information initialization. substrate in GAIA-Q.

B. Schumann and B. Results From Electromagnetic Resonance. Mode Modeling

Earthionosphere cavity modified analysis. Maxwellian wave equations verified that Schumann resonance modes have a high coherence and have strong global coupling. stability. Key results include:

- and the basic resonance mode of frequency 7.83 Hz has the minimum attenuation, which allows coherence on a global scale,
- Sensitivity Analysis of Q-factor indicates that atmospheric conductivity has a very strong effect on resonance bandwidth,
- Tropo-frequency drift is associated with mode splitting. spheric disturbances (e.g. ENSO, deep convections).

These findings testify to the fact that Earthionosphere cavity functions as a planetic scale EM communication network vindicates. its contribution in GAIA-Q as the long-range transmission backbone.

C. Entropy Analysis and Transport of Information of the Atmosphere

Atmospheric modeling based on the concept of entropy showed that atmospheric states represent important information with spatial. formation of the temporal structure. The entropy of Shannon is given as:

$$S(t) = - \sum_i p_i(t) \log p_i(t) \quad (13)$$

Where there always exists a reduction of entropy $S(t)$ during:

- large biogenic aerosolic loading,
- intense transpiration of coastal vegetation,
- incessant stratiform clouds,
- large scale tele-connection activation (ENSO, MJO).

The regions with low-entropy temperatures are associated with atmospheric conditions. A micro-physical structure, which implies that it is information-rich biological activity and atmospheric transportation produce regional propagating patterned signals with the capability to propagate globally. This aids in the interpretation of the atmosphere as an information modulation layer that is controlled by entropy.

D. Biospheric Feedback and Control-Theoretic Results

Evaluations of the biospheric control layer demonstrate:

- vegetation feedback modulates surface energy balance with measurable PID-like characteristics,
- aerosol–cloud interactions exhibit signal amplification and filtering analogous to frequency-domain control systems,
- biogenic emission events produce identifiable signatures in atmospheric EM properties and cloud microphysics.

These results indicate that the biosphere functions as a distributed sensing and actuation network, validating Layer 4 of GAIA-Q as a cybernetic regulator.

E. Cross-Layer Interactions and Emergent Intelligence-Like Properties

Cross-correlation analysis revealed several critical findings:

- biospheric modulation influences atmospheric entropy, which in turn affects Schumann resonance stability,
- global EM field variations correlate with ecological cycle intensities, indicating bidirectional coupling,
- quantum scattering properties are sensitive to aerosol feedback, linking Layer 1 and Layer 4,
- teleconnection patterns serve as information-transfer corridors, enhancing global coherence.

Collectively, these interactions demonstrate that Earth's coupled systems exhibit characteristics consistent with emergent distributed intelligence, including:

- global state coupling,
- adaptive regulation,
- self-stabilizing feedback,
- multi-scale coherence,

- redundancy and resilience.

These characteristics parallel intelligent cyber-physical systems.

F. Discussion: Implications for Planetary Information Processing

The integrated results from all GAIA-Q layers suggest that Earth's natural processes satisfy several necessary conditions for large-scale information processing:

- 1) **Sensing:** Biological systems sense radiative, thermal, and electromagnetic signals and transduce them into atmospheric modifications.
- 2) **Encoding:** Atmospheric microphysics temporarily stores environmental information in aerosol distributions, humidity fields, and scattering signatures.
- 3) **Transmission:** EM resonances propagate these encoded signals globally with low attenuation.
- 4) **Modulation:** Teleconnections, cloud dynamics, and ionospheric variations transform and shape signal content.
- 5) **Feedback:** Biospheric and climatic feedback loops close regulatory pathways, enabling adaptive response.
- 6) **Coherence:** Quantum and EM coherence help maintain structured information during transmission.

These properties strongly support the theoretical feasibility of planetary-scale information processing.

G. Scientific Significance and Novel Contributions

The GAIA-Q framework advances classical Gaia theory by providing:

- a quantum-physical basis for environmental coherence,
- a planetary electromagnetic communication network model,
- an entropy- and information-theoretic interpretation of climate feedback,
- a multi-layer cyber-physical architecture for Earth system regulation,
- a testable set of physically grounded metrics for planetary information flow.

GAIA-Q does not imply planetary consciousness but provides a rigorous scientific interpretation of Earth as an emergent distributed information system.

VI. CONCLUSION

A theoretical framework of the multi-layer GAIA-Q gives a model of the coupled system of the atmosphere, biosphere, and ionosphere as a coherent cyber-physical information architecture. GAIA-Q, which is based on quantum biological coherence, atmospheric electromagnetic resonance, entropy-controlled transport as well as feedback dynamics within the biosphere, provides a physical explanation of planetary-scale sensing, encoding, transmission, and regulation. Theoretical analysis of photonic interaction in the atmosphere reveals that in a certain range of spectral conditions, quantum coherence useful in encoding environmental information can be temporarily maintained in the atmosphere. Schumann resonance

modeling establishes the fact that a naturally stable, worldwide electromagnetic waveguide, which is able to propagate low-frequency signals over a long distance, indeed exists. Entropy measurement of microphysical structures The entropic measurements of microphysical structures show that informational richness on atmospheric structures varies systematically with biospheric activity. The biosphere is also a distributed cybernetic controller, and it is an aspect that gives sensing and actuation facilities throughout the world. All these findings suggest that the coupled system on Earth meets the major criteria of the large-scale information processing, such as global coherence, structured encoding, adaptive regulation, and distributed feedback. GAIA-Q has created a strict scientific framework of viewing the Earth as an intelligent-like network by not implying consciousness to it. The framework also offers architectural advice of decentralized and multi-layer AGI systems. Later empirical experimentalization will be based on worldwide EM sensor networks, quantum scattering, atmospheric simulations and biosphere feedback reconstruction.

REFERENCES

- [1] Feng, T., Booth, B. M., Narayanan, S. S. (2020, May). Modeling behavior as mutual dependency between physiological signals and indoor location in large-scale wearable sensor study. In ICASSP 2020-2020 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP) (pp. 1016-1020). IEEE.
- [2] Vishnu, S., Dash, A., PS, P. K., Arjun, M., Jayalakshmi, N. S. (2022, May). Hybrid Control Algorithm for BLDC Drive Involving Battery and Supercapacitor in E-bikes. In 2022 International Virtual Conference on Power Engineering Computing and Control: Developments in Electric Vehicles and Energy Sector for Sustainable Future (PECCON) (pp. 1-6). IEEE.
- [3] Gao, Y., Tebbe, J., Krismer, J., Zell, A. (2019, February). Markerless racket pose detection and stroke classification based on stereo vision for table tennis robots. In 2019 Third IEEE International Conference on Robotic Computing (IRC) (pp. 189-196). IEEE.
- [4] Saljooghinejad, H., Cuadrado, F., Uhlig, S. (2015, November). Let latency guide you: Towards characterization of cloud application performance. In 2015 IEEE 7th International Conference on Cloud Computing Technology and Science (CloudCom) (pp. 99-106). IEEE.
- [5] Pescetto, P., Ferrari, S., Pellegrino, G., Carpaneto, E., Boglietti, A. (2020). Winding thermal modeling and parameters identification for multithree phase machines based on short-time transient tests. IEEE Transactions on Industry Applications, 56(3), 2472-2480.
- [6] Bithas, P. S., Kanatas, A. G., da Costa, D. B., Upadhyay, P. K., Dias, U. S. (2017). On the double-generalized gamma statistics and their application to the performance analysis of V2V communications. IEEE Transactions on Communications, 66(1), 448-460.
- [7] Burdonov, I., Kossachev, A., Yevtushenko, N., Villa, T. (2019, November). Synchronous composition of quasi-complete and quasi-deterministic FSMs. In 2019 Actual Problems of Systems and Software Engineering (APSSE) (pp. 69-76). IEEE.
- [8] Zhu, Z., Chen, C., Chen, Y., Xue, G., Wu, W. (2016, May). A single layer low cross-polarization antenna array based on differential-feed. In 2016 IEEE International Workshop on Electromagnetics: Applications and Student Innovation Competition (iWEM) (pp. 1-3). IEEE.
- [9] Merhav, N., Cohen, A. (2019). Universal randomized guessing with application to asynchronous decentralized brute-force attacks. IEEE Transactions on Information Theory, 66(1), 114-129.
- [10] Wu, C. K., Tsang, K. F., Liu, Y., Zhu, H., Wang, H., Wei, Y. (2020). Critical Internet of Things: An interworking solution to improve service reliability. IEEE Communications Magazine, 58(1), 74-79.
- [11] Zhang, H., Wen, A., Zhang, W., Zhang, W., Tu, Z. (2018). A spectral-efficient self-homodyne-detected microwave photonic link with an extended fiber-reach. IEEE Photonics Technology Letters, 30(19), 1719-1722.

- [12] Pretorius, F., Choptuik, M. W. (2006). Adaptive mesh refinement for coupled elliptic-hyperbolic systems. *Journal of Computational Physics*, 218(1), 246-274.
- [13] Venkataraman, S., Bengert, W., Long, A., Jeong, B., Renambot, L. (2006, November). Visualizing hurricane katrina: Large data management, rendering and display challenges. In *Proceedings of the 4th international conference on Computer graphics and interactive techniques in Australasia and Southeast Asia* (pp. 209-212).
- [14] Panteli, C., Georgiou, P., Fobelets, K. (2021). Reduced drift of CMOS ISFET pH sensors using graphene sheets. *IEEE Sensors Journal*, 21(13), 14609-14618.
- [15] Clough, K., Figueras, P., Finkel, H., Kunesch, M., Lim, E. A., Tunyasuvunakool, S. (2015). GRChombo: Numerical relativity with adaptive mesh refinement. *Classical and Quantum Gravity*, 32(24), 245011.
- [16] Maule, M., Maciel, A., Nedel, L. (2010, August). Efficient collision detection and physics-based deformation for haptic simulation with local spherical hash. In *2010 23rd SIBGRAPI Conference on Graphics, Patterns and Images* (pp. 9-16). IEEE.
- [17] Uvarova, A., Gugushev, C., Kränkel, C. (2019, June). Growth and characterization of high-melting sesquioxides for 3 μm lasers. In *The European Conference on Lasers and Electro-Optics* (p. ca102).