

Unadjusted Effects of Risk Factors on Binary Stunting Outcomes

Jade Benjamin-Chung, Andrew N. Mertens, Esther Chung, Jeremy Coyle, Oleg Sofrygin, Nima Hejazi, Ivana Malenica, Alan E. Hubbard, Mark J. van der Laan, John M. Colford, Jr., Benjamin F. Arnold

2018-06-14


Contents

Relative risk pooled across cohorts	1
Point prevalence of stunting	1
Appendix 1 - Relative risk stratified by cohort	19
Point prevalence of stunting	19
Appendix 2 - Population Attributable Risk stratified by cohort	54
Point prevalence of stunting	54
Appendix 3 - Population Attributable Fraction stratified by cohort	89
Point prevalence of stunting	89


Relative risk pooled across cohorts

Point prevalence of stunting


Child characteristics




Birth length
Estimates from 7 cohorts in Asia, Africa




Birthweight
Estimates from 2 cohorts in Asia, Africa




Calendar month of birth
Estimates from 10 cohorts in Asia, Africa, Europe




Calendar month of measurement
Estimates from 11 cohorts in Asia, Africa, Europe




Child delivered at home
Estimates from 4 cohorts in Asia




Child's sex
Estimates from 18 cohorts in Asia, Latin America, Africa, Europe




Enrolled wasted
Estimates from 16 cohorts in Latin America, Asia, Africa, Europe




Gestational age at birth in days
Estimates from 9 cohorts in Asia, Latin America, Africa, Europe




Percent diarrhea in first 24 months
Estimates from 8 cohorts in Africa, Latin America, Asia




Percent diarrhea in first 6 months
Estimates from 1 cohorts in Asia




Percent exclusive breastfeeding
Estimates from 11 cohorts in Asia, Africa, Latin America




Persistent wasting in first 6 months
Estimates from 2 cohorts in Asia





Stunted at enrollment
Estimates from 20 cohorts in Asia, Africa, Latin America, Europe




Vaginal delivery
Estimates from 9 cohorts in Asia, Europe, Africa




Household characteristics




Improved floor
Estimates from 8 cohorts in Asia, Latin America




Improved sanitation
Estimates from 11 cohorts in Latin America, Asia




Number of children in the household under 5 years
Estimates from 2 cohorts in Asia




Number of people in the household
Estimates from 6 cohorts in Asia, Africa




Number of rooms in household
Estimates from 12 cohorts in Asia, Latin America, Africa




Safe water
Estimates from 2 cohorts in Asia, Africa




Treats water
Estimates from 1 cohorts in Asia




USAID Household Food Insecurity Access Scale (HFIAS) category
Estimates from 5 cohorts in Asia, Latin America




Maternal characteristics




Maternal height
Estimates from 5 cohorts in Latin America, Asia




Maternal parity
Estimates from 7 cohorts in Latin America, Asia, Africa




Maternal weight
Estimates from 9 cohorts in Africa, Asia, Latin America, Europe



Mother is unmarried
Estimates from 6 cohorts in Latin America, Asia, Africa




Mother's education
Estimates from 15 cohorts in Asia, Latin America, Africa, Europe




Paternal characteristics


Father age
Estimates from 2 cohorts in Latin America, Asia



Father height
Estimates from 5 cohorts in Latin America, Africa, Europe, Asia




Father's education
Estimates from 4 cohorts in Asia, Africa





Appendix 1 - Relative risk stratified by cohort

Point prevalence of stunting


Child characteristics









Calendar month of birth
reference level: 1







Child delivered at home
reference level:0




Child's sex
reference level:Female







Gestational age at birth in days
reference level:[273–287)




Percent diarrhea in first 24 months
reference level:Q1




Percent diarrhea in first 6 months
reference level:Q1







Persistent wasting in first 6 months
reference level:0




Stunted at enrollment
reference level:0








Household characteristics




Clean cook
reference level:0









Number of children in the household under 5 years
reference level:1




Number of people in the household
reference level:3 or less




Number of rooms in household
reference level:1







Treats water
reference level:0




USAID Household Food Insecurity Access Scale (HFIAS) category
reference level: Food Secure




Maternal characteristics




Maternal BMI
reference level:[18.5–25)




Maternal height
reference level: ≥ 160







Maternal weight
reference level: ≥ 57.5




Mother is unmarried
reference level:0




Mother's education
reference level:Q4




Paternal characteristics






Father's education
reference level:Q4





Appendix 2 - Population Attributable Risk stratified by cohort

Point prevalence of stunting


Child characteristics









Calendar month of birth
reference level:1







Child delivered at home
reference level:0




Child's sex
reference level:Female







Gestational age at birth in days
reference level:[273–287)




Percent diarrhea in first 24 months
reference level:Q1




Percent diarrhea in first 6 months
reference level:Q1







Persistent wasting in first 6 months
reference level:0




Stunted at enrollment
reference level:0




Vaginal delivery
reference level:0





Household characteristics




Clean cook
reference level:0









Number of children in the household under 5 years
reference level:1





Number of people in the household
reference level:3 or less




Number of rooms in household
reference level:1









USAID Household Food Insecurity Access Scale (HFIAS) category
reference level: Food Secure





Maternal characteristics




Maternal BMI
reference level:[18.5–25)












Mother is unmarried
reference level:0





Mother's education
reference level:Q4



Paternal characteristics










Appendix 3 - Population Attributable Fraction stratified by cohort

Point prevalence of stunting


Child characteristics









Calendar month of birth
reference level:1







Child delivered at home
reference level:0




Child's sex
reference level:Female




Enrolled wasted
reference level:0




Gestational age at birth in days
reference level:[273–287)




Percent diarrhea in first 24 months
reference level:Q1




Percent diarrhea in first 6 months
reference level:Q1







Persistent wasting in first 6 months
reference level:0




Stunted at enrollment
reference level:0








Household characteristics




Clean cook
reference level:0










Number of children in the household under 5 years
reference level:1




Number of people in the household
reference level:3 or less












USAID Household Food Insecurity Access Scale (HFIAS) category
reference level: Food Secure





Maternal characteristics




Maternal BMI
reference level:[18.5–25)











Mother is unmarried
reference level:0



Mother's education
reference level:Q4



Paternal characteristics

