

**1**

(a) Show that every totally separated space has a continuous injection into  $\{0, 1\}^J$  for some index set  $J$ .

*Proof.* Let  $X$  be a totally separated space. For every pair of distinct points  $x, y \in X$  there is a clopen subset  $A \subseteq X$  separating  $x$  and  $y$ , i.e.,  $x \in A$  and  $y \in A^c$ . Let  $J$  be a separating collection of clopen subsets of  $X$ , i.e., every distinct pair of points in  $X$  is separated by some clopen set in  $J$ . (We could take  $J$  to be the collection of all clopen subsets of  $X$ , but this is not necessary.)

For each  $A \in J$ , consider the indicator/characteristic function  $\chi_A : X \rightarrow \{0, 1\}$  defined by  $\chi_A(x) = 1$  if and only if  $x \in A$ . We check that  $\chi_A$  is continuous; there are four preimages:

$$\chi_A^{-1}(\emptyset) = \emptyset, \quad \chi_A^{-1}(\{0\}) = A^c, \quad \chi_A^{-1}(\{1\}) = A, \quad \chi_A^{-1}(\{0, 1\}) = X.$$

Since  $A$  is clopen, all four sets are open in  $X$ , so in fact  $\chi_A$  is continuous.

By the universal property of the product, there is a unique continuous map  $f : X \rightarrow \{0, 1\}^J$  such that  $\pi_A \circ f = \chi_A$  for all  $A \in J$ . By the assumption on  $J$ , given a pair of distinct points  $x, y \in X$ , there is a clopen set  $A \in J$  separating  $x$  and  $y$ . Then  $\chi_A(x) = 1$  and  $\chi_A(y) = 0$ , implying that  $f(x) \neq f(y)$ . Hence,  $f$  is injective.  $\square$

(b) Show that every second-countable totally separated space has a continuous injection into  $\{0, 1\}^{\mathbb{N}}$ .

*Proof.* Let  $X$  be a totally separated space with countable basis  $\mathcal{B}$ . Define the set:

$$\mathcal{S} = \{(U, V) \in \mathcal{B}^2 : U \subseteq A \subseteq V^c \text{ for some clopen } A \subseteq X\}.$$

In other words,  $\mathcal{S}$  is the pairs of basis elements which are separated by some clopen set. Note that  $\mathcal{S}$  is countable, so we can enumerate its elements by  $\mathcal{S} = \{(U_k, V_k)\}_{k \in \mathbb{N}}$ . For each  $k \in \mathbb{N}$ , choose a representative clopen set  $A_k \subseteq X$  separating  $U_k$  and  $V_k$ .

Then  $J = \{A_k\}_{k \in \mathbb{N}}$  is a countable collection of clopen subsets of  $X$ . In order to apply part (a), we must show that  $J$  separates points in  $X$ . Given distinct points  $x, y \in X$ , there is a clopen set  $A \subseteq X$  separating  $x$  and  $y$ . Since  $A$  and  $A^c$  are open neighborhoods of  $x$  and  $y$ , respectively, there are basis elements  $U, V \in \mathcal{B}$  such that  $x \in U \subseteq A$  and  $y \in V \subseteq A^c$ . In particular,  $A$  is a clopen set separating  $U$  and  $V$ , implying  $(U, V) \in \mathcal{S}$ . By construction, there is some  $A_k \in J$  separating  $U_k$  and  $V_k$ . Hence,  $J$  is a countable collection of clopen sets which separates points in  $X$ .

By part (a), there is a continuous injection  $X \rightarrow \{0, 1\}^J$ . Since  $J$  is countable, there is a natural homeomorphism  $\{0, 1\}^J \xrightarrow{\sim} \{0, 1\}^{\mathbb{N}}$  by identifying the indexing elements  $A_k \in J$  and  $k \in \mathbb{N}$ . Composing, we obtain a continuous injection  $X \rightarrow \{0, 1\}^{\mathbb{N}}$ .  $\square$

(c) Show that  $\{0, 1\}^{\mathbb{N}}$  is homeomorphic to the *middle-thirds Cantor set*: the set of numbers in  $[0, 1]$  which have a base 3 expansion consisting only of 0's and 2's, with the subspace topology inherited from  $\mathbb{R}$ .

*Proof.* Let  $C \subseteq [0, 1]$  be the middle-thirds Cantor set. Define the map  $f : \{0, 1\}^{\mathbb{N}} \rightarrow C$  by

$$(a_n)_{n \in \mathbb{N}} \mapsto \sum_{n \in \mathbb{N}} \frac{2a_n}{3^n}.$$

In other words,  $f$  maps a sequence of 0's and 1's to the corresponding base 3 decimal expansions of 0's and 2's. It is clear that  $f$  is bijective since elements of  $C$  are uniquely determined by their base 3 expansion and all base 3 expansions are described by a sequence of 0's and 1's.

The balls  $B_{2/3^k}(x)$ —for  $x \in \mathbb{R}$ ,  $k \in \mathbb{N}$ —generate the usual topology on  $\mathbb{R}$ . Therefore, it suffices to check the continuity of  $f$  on these balls. Fix a point  $x \in C$  and define  $a = f^{-1}(x)$ , then we have the decimal expansion

$$x = \sum_{n \in \mathbb{N}} \frac{2a_n}{3^n}.$$

Fix some  $k \in \mathbb{N}$  and consider the ball

$$B_{2/3^k}(x) = \{y \in C : |x - y| < 2/3^k\}.$$

Given  $y \in C$  define  $b = f^{-1}(y)$ , then

$$|x - y| = \sum_{n \in \mathbb{N}} \frac{2|a_n - b_n|}{3^n}.$$

For  $m \in \mathbb{N}$ , we have

$$\sum_{n \geq m} \frac{2|a_n - b_n|}{3^n} \leq \sum_{n \geq m} \frac{2}{3^n} = \frac{1}{3^{m-1}}.$$

From this, we deduce that  $|x - y| < 2/3^k$  if and only if  $a_n = b_n$  for all  $n \leq k$ . In other words,

$$f^{-1}(B_{2/3^k}(x)) = \bigcap_{n=1}^k \pi_n^{-1}(a_n),$$

where  $\pi_n : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}$  is the projection map to the  $n$ th coordinate. As the finite intersection of subbasis elements,  $f^{-1}(B_{2/3^k}(x))$  is open in  $\{0, 1\}^{\mathbb{N}}$ .

The preimages  $\pi_k^{-1}(t)$ —for  $t \in \{0, 1\}$ ,  $k \in \mathbb{N}$ —generate the product topology on  $\{0, 1\}^{\mathbb{N}}$ . Therefore, it suffices to check the continuity of  $f^{-1}$  on these sets. Fix values  $t \in \{0, 1\}$  and  $k \in \mathbb{N}$ , and consider the subbasis set

$$U = \pi_k^{-1}(t) = \{a \in \{0, 1\}^{\mathbb{N}} : a_k = t\}.$$

Let  $A \subseteq \{0, 1\}^{\mathbb{N}}$  be a set containing a choice of representative  $a \in \{0, 1\}^{\mathbb{N}}$  for each combination of  $a_1, \dots, a_{k-1}$  and  $a_k = t$ . (Note that  $|A| = 2^{k-1} < \infty$ .) Then we can write

$$U = \bigcup_{a \in A} \bigcap_{n=1}^k \pi_n^{-1}(a_n) = \bigcup_{a \in A} f^{-1}(B_{2/3^k}(a)),$$

so

$$f(U) = \bigcup_{a \in A} B_{2/3^k}(a).$$

As the union of finitely many open balls,  $f(U)$  is open in  $C$ . We conclude that  $f$  is an open mapping, hence a homeomorphism.  $\square$

**2** Let  $X$  be a locally compact Hausdorff space.

(a) Show that if  $K$  is compact in  $X$  and  $U$  is an open set containing  $K$ , then there is a function  $f : X \rightarrow [0, 1]$  which is supported on  $U$  and such that  $f(K) = 1$ .

**Hint:** First find a larger compact set whose interior contains  $K$ .

**Lemma 1.** Open subspaces of  $X$  are locally compact.

*Proof.* Let  $V \subseteq X$  be an open subset and  $x \in V$ . (The result is trivial if  $V$  is empty.) Since  $X$  is locally compact,  $x$  has a compact neighborhood  $L \subseteq X$ . Since  $X$  is Hausdorff,  $L$  is closed in  $X$ , so  $E = (X \setminus V) \cap L$  is also closed. Then  $E$  is compact as a closed subset of the compact set  $L$ .

(It is easy to show that disjoint compact subsets of a Hausdorff space are separated by disjoint open sets. Moreover, the condition that two sets  $A, B$  are separated is equivalent to the existence of an open set  $W$  such that  $A \subseteq W$  and  $\overline{W} \subseteq B^c$ .)

Since  $X$  is Hausdorff, there is an closed set  $F \subseteq X$  such that  $x \in \text{int } F$  and  $F \subseteq X \setminus E$ . Note

$$X \setminus E = (X \setminus (X \setminus V)) \cup (X \setminus L) = V \cup (X \setminus L),$$

so

$$F \cap L \subseteq (X \setminus E) \cap L = V.$$

Moreover,

$$x \in \text{int } F \cap \text{int } L = \text{int}(F \cap L),$$

so  $F \cap L$  is a compact neighborhood of  $x$  contained in  $V$ . □

We now prove the main result.

*Proof.* Applying Lemma 1 to each  $x \in K \subseteq U$ , there is a compact set  $L_x \subseteq X$  such that  $x \in \text{int } L_x$  and  $L_x \subseteq U$ . Then the collection of interiors  $\{\text{int } L_x\}_{x \in K}$  is an open cover of the compact set  $K$ , so there is a finite subcover  $\{\text{int } L_{x_i}\}_{i=1}^n$ . Then the union  $L = \bigcup_{i=1}^n L_{x_i}$  is a compact subset of  $U$  with

$$K \subseteq \bigcup_{i=1}^n \text{int } L_{x_i} \subseteq \text{int } L.$$

Now  $L$  is compact and Hausdorff—therefore normal. By Urysohn's lemma, there is a continuous function  $f : L \rightarrow [0, 1]$  such that  $f|_K = 1$  and  $f|_{L \setminus \text{int } L} = 0$ . We can extend  $f$  to a function  $X \rightarrow [0, 1]$  by defining  $f|_{X \setminus L} = 0$ .

By construction, the support of  $f$  is contained in  $\text{int } L \subseteq U$ . We have that  $f|_L$  is continuous, but we must check that  $f$  is continuous on all of  $X$ . Given an open set  $U \subseteq [0, 1]$  there are two cases we must consider:  $0 \in U$  and  $0 \notin U$ .

If  $0 \notin U$ , then  $f^{-1}(U) \subseteq \text{int } L$ . Since  $f|_{\text{int } L}$  is continuous,  $f^{-1}(U)$  is an open subset of the open subspace  $\text{int } L \subseteq X$  and—therefore—also an open subset of  $X$ .

If  $0 \in U$ , then we write

$$f^{-1}(U) = f|_L^{-1}(U) \cup (X \setminus L).$$

Since  $f|_L$  is continuous,  $f|_L^{-1}(U)$  is open in the subspace  $L \subseteq X$ , i.e., there is an open set  $V \subseteq X$  such that  $f|_L^{-1}(U) = V \cap L$ . Then we have

$$f^{-1}(U) = (V \cap L) \cup (X \setminus L) = V \cup (X \setminus L).$$

Since  $L$  is compact and  $X$  is Hausdorff,  $L$  is closed in  $X$ . Therefore  $X \setminus L$  is open in  $X$ , implying  $f^{-1}(U)$  is also open in  $X$ . Hence  $f$  is continuous.  $\square$

**(b)** Define the subspaces  $C_c(X) \subseteq C_0(X) \subseteq C_B(X)$  as follows:

- A function is in  $C_c(X)$  if it is *compactly supported*, i.e. it is zero outside a compact set.
- A function  $f \in C_0(X)$  if it “vanishes at infinity”, i.e. for every  $\varepsilon > 0$ ,  $\{x \in X : |f(x)| \geq \varepsilon\}$  is compact.

Show that  $C_0(X)$  is the closure of  $C_c(X)$  in the sup norm topology.

*Proof.* Let  $f \in C_0(X)$ . For  $\varepsilon > 0$  consider the compactly supported supremum norm ball

$$B_\varepsilon(f) = \{g \in C_0(X) : \|f - g\|_\infty < \varepsilon\}.$$

We want to show that  $B_\varepsilon(f) \cap C_c(X) \neq \emptyset$ .

Define  $\delta = \varepsilon/2$  and the function  $g : X \rightarrow \mathbb{R}$  by

$$g(x) = \max\{0, f(x) - \delta\} + \min\{0, f(x) + \delta\}.$$

The operations preserve continuity and boundedness, so  $g \in C_B(X)$ . Moreover, this definition gives us  $\|f - g\|_\infty \leq \delta < \varepsilon$ , implying  $g \in B_\varepsilon(f)$ .

Since  $f \in C_0(X)$ , the set  $K = \{x \in X : |f(x)| \geq \delta\}$  is compact. For every  $x \in X \setminus K$ , we have  $|f(x)| < \delta$ . Note that  $f(x) \leq 0$  implies  $f(x) > -\delta$ , and  $f(x) \geq 0$  implies  $f(x) < \delta$ . In either case,  $g(x) = 0$ . Hence, the support of  $g$  is contained in  $K$ , so in fact  $g \in C_c(X)$ .  $\square$

**3** A set  $A \subset X$  is a *retract* if there is a continuous map  $X \rightarrow A$  which is the identity on  $A$ .

- Convince yourself that  $\{0, 1\}$  is not a retract of  $[0, 1]$ .
- Convince yourself that the two obvious circles are (separately) retracts of the torus.

**(a)** Show that if  $A$  is a retract of a Hausdorff space  $X$ , then  $A$  is closed in  $X$ .

*Proof.* Let  $r : X \rightarrow A$  be a retraction map, i.e.,  $r$  is continuous and  $r|_A = \text{id}_A$ . We will show that  $A^c$  is open in  $X$ . Given a point  $x \in A^c$ , we know  $x \neq r(x)$ . Since  $X$  is Hausdorff, there are disjoint open sets  $U, V \subseteq X$  such that  $x \in U$  and  $r(x) \in V$ . Then  $V \cap A$  is open in the subspace  $A \subseteq X$ . Since  $r$  is continuous, the preimage  $r^{-1}(V \cap A)$  is open in  $X$ . Then the intersection  $W = U \cap r^{-1}(V \cap A)$  is an open neighborhood of  $x$ .

Since  $r$  restricts to the identity on  $A$ , we know that

$$r^{-1}(V \cap A) \cap A = V \cap A.$$

Since  $U$  is disjoint from  $V$ , it is also disjoint from  $V \cap A$ . Therefore,

$$W \cap A = U \cap r^{-1}(V \cap A) \cap A = U \cap V \cap A = \emptyset,$$

so  $W \subseteq A^c$ , hence  $A^c$  is open in  $X$ . □

**(b)** A space  $Y$  is an *absolute retract* if whenever  $X$  is a normal space which contains a closed set  $Y_0$  homeomorphic to  $Y$ , then  $X$  retracts to  $Y_0$ . Show that  $\mathbb{R}^J$  is an absolute retract, for any index set  $J$ .

*Proof.* Let  $X$  be a normal space with a closed subspace  $Y$  homeomorphic to  $\mathbb{R}^J$  for some index set  $J$ . For  $j \in J$ , let  $\pi_j : \mathbb{R}^J \rightarrow \mathbb{R}$  be the projection to the  $j$ th coordinate. The homeomorphism  $f : Y \xrightarrow{\sim} \mathbb{R}^J$  is characterized by its components  $f_j = \pi_j \circ f : Y \rightarrow \mathbb{R}$ .

By Tietze, there is a continuous extension  $F_j : X \rightarrow \mathbb{R}$  such that  $F_j|_Y = f_j$ . By the universal property of the topological product, there is a unique continuous map  $F : X \rightarrow \mathbb{R}^J$  such that  $F_j = \pi_j \circ F$ . Moreover, this construction gives us  $F|_Y = f$ .

Define the continuous map  $r = f^{-1} \circ F : X \rightarrow Y$ . Then  $r|_Y = f^{-1} \circ F|_Y = \text{id}_Y$ , so  $r$  describes a retraction of  $X$  onto  $Y$ . □

4 Show that a metric space  $X$  is compact if and only if every continuous function  $f : X \rightarrow \mathbb{R}$  is bounded.

*Proof.* If  $X$  is compact and  $f : X \rightarrow \mathbb{R}$  is continuous, then the image  $f(X) \subseteq \mathbb{R}$  is compact. In particular,  $f(X)$  is a bounded subset of  $\mathbb{R}$ , implying that  $f$  is a bounded function.

If  $X$  is not compact, then it is not sequentially compact, i.e., we can find a sequence  $(x_n)_{n \in \mathbb{N}}$  of points in  $X$  with no convergent subsequence. Without loss of generality, we may assume that the  $x_n$ 's are distinct (since there must be no infinitely recurring terms, otherwise they would form a trivially convergent subsequence).

In particular, each  $x_n$  is not a limit point of the image of the sequence, so we can find a radius  $r_n > 0$  such that the ball  $B_{r_n}(x_n)$  contains no other points of the sequence. Define the smaller ball  $B_n = B_{\varepsilon_n/2}(x_n)$ , which is an open neighborhood of  $x_n$ . For  $n \neq m$ , we know that  $d(x_n, x_m) < \max\{r_n, r_m\}$ , which implies that  $B_n \cap B_m = \emptyset$ .

For each  $n \in \mathbb{N}$ , we use Urysohn's lemma to construct a continuous map  $f_n : X \rightarrow [0, 1]$  such that  $f_n(x_n) = 1$  and  $f_n|_{B_n^c} = 0$ . Define  $f = \sum_{n \in \mathbb{N}} n f_n$ ; since the supports of the  $f_n$ 's are disjoint, this sum is well-defined. However,  $f(x_n) = n$  for all  $n \in \mathbb{N}$ , so this function is unbounded.  $\square$