

## Notation

Let  $K$  denote an algebraically closed ground field.

Let  $K[x_1, \dots, x_n]$  to be the  $K$ -algebra of polynomials, graded by degree. We will mostly focus on  $K[x, y]$ .

For  $n \in \mathbb{N}$ , we call  $\mathbb{A}^n = \mathbb{A}_K^n$  the **affine  $n$ -space** over  $K$ .

For  $S \subseteq K[x_1, \dots, x_n]$ , call

$$V(S) = \{x \in \mathbb{A}^n : f(x) = 0 \text{ for all } f \in S\}$$

the (affine) **zero locus** of  $S$ . Subsets of  $\mathbb{A}^n$  of this form are called **affine varieties**.

**Definition 1.** (Affine Curves)

- (a) An **(affine plane algebraic) curve** is a nonconstant polynomial  $F \in K[x, y]$  modulo units, i.e., modulo the equivalence relation  $F \sim G$  if  $F = \lambda G$  for some  $\lambda \in K^\times$ .

Call  $V(F) = \{P \in \mathbb{A}^2 : F(P) = 0\}$  the **set of points** of  $F$ .

- (b) The **degree** of a curve is its degree as a polynomial, denoted  $\deg F$ .
- (c) A curve  $F$  is called **irreducible** if it is as a polynomial, and **reducible** otherwise. Similarly, if  $F = F_1^{d_1} \cdots F_k^{d_k}$  is the irreducible decomposition of  $F$  as a polynomial, we will also call this the **irreducible decomposition** of the curve  $F$ . The curves  $F_1, \dots, F_k$  are called the **(irreducible) components** of  $F$  and  $d_1, \dots, d_k$  their multiplicities.

**Lemma 1.** Let  $F$  be an affine curve.

- (a) If  $K$  is algebraically closed then  $V(F)$  is infinite.
- (b) If  $K$  is infinite then  $\mathbb{A}_K^2 \setminus V(F)$  is infinite.

**Proposition 1.** If two curves  $F$  and  $G$  have no common component then their intersection  $V(F, G)$  is finite.

**Corollary 1.** Let  $F$  be a curve over an algebraically closed field. Then for any irreducible curve  $G$  we have  $G \mid F$  if and only if  $V(G) \subseteq V(F)$ .

In particular, the irreducible components of  $F$  (but not their multiplicities) can be recovered from  $V(F)$ .

**Notation 1.** Due to the above correspondence between a curve  $F$  and its set of points  $V(F)$ , we will sometimes write

- (a)  $P \in F$  instead of  $P \in V(F)$ , i.e.,  $F(P) = 0$ ;
- (b)  $F \cap G$  instead of  $V(F, G)$  for the points that lie on both  $F$  and  $G$ ;
- (c)  $F \cup G$  for the curve  $FG$ ;
- (d)  $G \subseteq F$  instead of  $G \mid F$ .

**Definition 2.** Let  $a \in \mathbb{A}^2$  be a point.

(a) The **local ring** of  $\mathbb{A}^2$  at  $P$  is defined as

$$\mathcal{O}_a = \mathcal{O}_{\mathbb{A}^2, a} = \left\{ \frac{g}{f} : f, g \in K[x, y] \text{ with } f(a) \neq 0 \right\} \subseteq K(x, y)$$

(b) It admits a well-defined ring homomorphism

$$\mathcal{O}_a \rightarrow K, \frac{g}{f} \mapsto \frac{g(a)}{f(a)}$$

which we call the **evaluation map**. Its kernel will be denoted by

$$I_a = I_{\mathbb{A}^2, a} = \{\varphi \in \mathcal{O}_a \mid \varphi(a) = 0\}$$

which is the unique maximal ideal in  $\mathcal{O}_a$ .

**Definition 3.** For a point  $a \in \mathbb{A}^2$  and two curves  $F$  and  $G$  we define the **intersection multiplicity** of  $F$  and  $G$  at  $a$  to be

$$\mu_a(F, G) = \dim \mathcal{O}_a / \langle F, G \rangle \in \mathbb{N} \cup \{\infty\},$$

where  $\dim$  denotes the dimension as a vector space over  $K$ .

**Lemma 2.** Let  $a \in \mathbb{A}^2$  and let  $F$  and  $G$  be two curves. We have

- (a)  $\mu_a(F, G) \geq 1$  if and only if  $a \in F \cap G$ ;
- (b)  $\mu_a(F, G) = 1$  if and only if  $\langle F, G \rangle = I_a$  in  $\mathcal{O}_a$ .

**Notation 2.** For a polynomial  $F \in K[x, y]$  of degree  $d$  and  $i = 0, \dots, d$ , we define the **degree- $i$  part** of  $F$  to be the sum of all terms of  $F$  of degree  $i$ . Hence all  $F_i$  are homogeneous, and we have  $F = F_0 + \dots + F_d$ . We call  $F_0$  the **constant** part,  $F_1$  the **linear** part, and  $F_d$  the **leading** part of  $F$ .

**Proposition 2.** Let  $F$  and  $G$  be two curves through the origin. Then  $\mu_0(F, G) = 1$  if and only if the linear parts of  $F$  and  $G$  are linearly independent.

## 1 projective

**Definition 4.** For  $n \in \mathbb{N}$ , we define the **projective  $n$ -space** over  $K$  as the set of all 1-dimensional linear subspaces of  $K^{n+1}$ . It is denoted by  $\mathbb{P}_K^n$  or simply  $\mathbb{P}^n$ .

In other words, we have

$$\mathbb{P}^n = (K^{n+1} \setminus \{0\}) / \sim$$

with the equivalence relation  $x \sim y$  if and only if  $x = \lambda y$  for some  $\lambda \in K^\times$ . We denote the equivalence class of  $(x_0, \dots, x_n)$  by  $[x_0 : \dots : x_n] \in \mathbb{P}^n$ . Call  $x_0, \dots, x_n$  the **homogeneous** or **projective coordinate** of the point  $[x_0 : \dots : x_n]$ .

For a subset  $S \subseteq K[x_0, \dots, x_n]$  of homogeneous polynomials we call

$$V(S) = \{P \in \mathbb{P}^n : f(P) = 0 \text{ for all } f \in S\} \subseteq \mathbb{P}^n$$

the projective **zero locus** of  $S$ . Subsets of  $\mathbb{P}^n$  of this form are called **projective varieties**.

**Definition 5.** (Projective curves) A **(projective plane algebraic) curve** is a nonconstant homogeneous polynomial  $F \in K[x, y, z]$  modulo units. We call  $V(F) = \{P \in \mathbb{P}^2 : F(P) = 0\}$  is **set of points**.

The **degree** of a projective curve is its degree as a polynomial.

The notions of irreducible/reducible/reduced curves, as well as irreducible components and their multiplicities, are defined in the same way as for affine curves.

**Construction 1.** For  $P \in \mathbb{P}^2$  we define the **local ring** of  $\mathbb{P}^2$  at  $P$  as

$$\mathcal{O}_P = \mathcal{O}_{\mathbb{P}^2, P} = \left\{ \frac{g}{f} : f, g \in K[x, y, z] \text{ homogeneous of the same degree with } f(P) \neq 0 \right\}$$

and the unique maximal ideal

$$I_P = I_{\mathbb{P}^2, P} = \{\varphi \in \mathcal{O}_P : \varphi(P) = 0\}.$$

There is an isomorphism  $\mathcal{O}_{\mathbb{P}^2, [x_0:y_0:1]} \xrightarrow{\sim} \mathcal{O}_{\mathbb{A}^2, (x_0, y_0)}$  given by  $\varphi \mapsto \varphi^i$ , which then restricts to  $I_{\mathbb{P}^2, [x_0:y_0:1]} \xrightarrow{\sim} I_{\mathbb{A}^2, (x_0, y_0)}$ .

**Construction 2.** Note that the local ring  $\mathcal{O}_{\mathbb{P}^2, P}$  does not contain  $K[x, y, z]$  as a subring. But for  $F_1, \dots, F_k$  homogeneous there is still a generated ideal

$$\langle F_1, \dots, F_k \rangle = \left\{ \frac{a_1}{b_1} F_1 + \dots + \frac{a_k}{b_k} F_k : a_i = 0 \text{ or } a_i b_i \text{ homogeneous with } \deg(a_i F_i) = \deg b_i \text{ for all } i \right\}$$

in  $\mathcal{O}_P$ . As in the affine case we can therefore define **intersection multiplicity** of two curves  $F$  and  $G$  at a point  $P \in \mathbb{P}^2$  as

$$\mu_P(F, G) = \dim \mathcal{O}_P / \langle F, G \rangle.$$