

Cite 5 dark matter detection papers from consultation, provide information on the most recent method (2018 theoretical coupling), and create a bar chart of publication year distribution (2005: 1, 2011: 1, 2012: 1, 2013: 1, 2018: 1)

HEP Multi-agent System
Argonne Computational Sciences Division

February 4, 2026

This report was autonomously generated by an LLM-based multi-agent system. All content, analysis, and conclusions should be independently verified.

1 Query

Search arxiv for 5 papers on dark matter detection, Briefly provide information on the most recent method. then create a bar chart showing the publication year distribution of the papers found.

2 Answer

Found 5 dark matter detection papers from arXiv spanning 2005-2018. The most recent method (2018) involves theoretical coupling mechanisms between dark matter and scalar fields, specifically examining non-minimal coupling between scalar fields and the Ricci scalar for renormalization in curved space-time. Created bar chart showing uniform distribution across publication years.

3 Execution Log

The research worker executed arXiv searches targeting specific years (2005, 2011, 2012, 2013, 2018) for dark matter detection papers. Initial broad searches returned mixed results including unrelated papers, requiring refinement with specific terms like "dark energy survey", "directional detection", and "direct detection". The worker successfully identified and retrieved metadata for all 5 target papers, then cited each using the cite() function to build references.bib.

The compute worker processed publication year data by extracting years [2005, 2011, 2012, 2013, 2018] with counts [1, 1, 1, 1, 1] and saved this as a JSON dictionary to publication_year_distribution.json for visualization.

The visualization worker encountered initial file path issues when attempting to create the bar chart. The worker listed available files, discovered the data file existed, but needed to reformat the JSON structure from {years: [...], counts: [...]} to {labels: [...], values: [...]} format. After reformatting, successfully generated the bar chart showing publication year distribution.

4 Results

The system identified **5 papers** spanning **13 years** (2005-2018) covering diverse dark matter detection approaches:

- **2005:** Astrophysical surveys using weak gravitational lensing [arXiv:astro-ph/0510346]
- **2011:** Directional detection exploiting Solar system rotation [arXiv:1110.1056]
- **2012:** Direct detection targeting 10^{-48} cm^2 [arXiv:1211.7222]
- **2013:** Four-pronged complementary detection strategy [arXiv:1305.1605]
- **2018:** Theoretical coupling mechanisms with scalar fields [arXiv:1809.09971]

The most recent method (2018) focuses on "non-minimal coupling between a scalar field and the Ricci scalar, which is necessary for renormalization of the scalar field in curved space-time" and proposes that "dark matter is uniquely connected to the inflaton" [arXiv:1809.09971].




Figure 1: Publication year distribution of dark matter detection papers showing uniform distribution across the surveyed timespan (2005-2018).

5 Discussion

Physical Plausibility: The progression from astrophysical surveys (2005) to theoretical coupling mechanisms (2018) reflects the field's evolution from observational to increasingly sophisticated theoretical approaches. The 2012 target sensitivity of 10^{-48} cm^2 represents the theoretical limit before neutrino backgrounds dominate, which

Key takeaways: The field demonstrates methodological diversity, with each approach addressing different aspects of the dark matter problem. The 2013 complementary strategy paper correctly identifies that "diversity of possible dark matter candidates requires a balanced program drawing from all four approaches" [arXiv:1305.1605].

Limitations: The search was constrained to 5 papers from specific years, potentially missing significant developments in intervening years. The uniform year distribution (1 paper per year) may not reflect actual research intensity patterns.

Gaps: No papers from 2014-2017 were included, missing potential methodological advances. The theoretical 2018 paper lacks experimental validation discussed in the abstract.

6 Developer Notes

No issues or recommendations reported by workers. The system successfully handled file path resolution and data format conversion automatically. All three workers (research, compute, visualization) completed their tasks without requiring manual intervention.

7 References

LLM-generated references. Quotes validated against source text.

✓ The Dark Energy Survey (2005)

The Dark Energy Survey Collaboration – arXiv:astro-ph/0510346

“optical-near infrared survey of 5000 sq. deg of the South Galactic Cap”

“measure the dark energy and dark matter densities”

“galaxy clusters, weak gravitational lensing tomography, galaxy angular clustering, and supernova distances”

✓ Directional detection of Dark Matter (2011)

F. Mayet and J. Billard and D. Santos – arXiv:1110.1056

“Taking advantage on the rotation of the Solar system around the galactic center through the Dark Matter halo”

“simultaneous measurement of the energy and the 3D track of low energy recoils”

“direction dependence of WIMP events”

✓ Direct dark matter detection: the next decade (2012)

Laura Baudis – arXiv:1211.7222

“ultimate WIMP detectors, capable of probing spin-independent interactions down to the unimaginably low cross section of $1e-48$ cm²”

“before the irreducible neutrino background takes over”

“constrain WIMP properties such as its mass, scattering cross section and possibly spin”

✓ Dark Matter in the Coming Decade: Complementary Paths to Discovery and Beyond (2013)

Daniel Bauer and James Buckley and Matthew Cahill-Rowley and Randel Cotta and Alex Drlica-Wagner and Jonathan L. Feng and Stefan Funk and JoAnne Hewett and Dan Hooper and Ahmed Ismail and Manoj Kaplinghat and Alexander Kusenko and Konstantin Matchev and Daniel McKinsey and Tom Rizzo and William Shepherd and Tim M. P. Tait and Alexander M. Wijangco and Matthew Wood – arXiv:1305.1605

“four complementary approaches: direct detection, indirect detection, collider experiments, and astrophysical probes”

“diversity of possible dark matter candidates requires a balanced program drawing from all four approaches”

“complementarity of the different dark matter searches”

✓ Possible Couplings of Dark Matter (2018)

Kevin J. Ludwick – arXiv:1809.09971

“non-minimal coupling between a scalar field and the Ricci scalar, which is necessary for renormalization of the scalar field in curved space-time”

“dark matter is uniquely connected to the inflaton”

“observational astrophysical constraints to specify an upper bound on the dark matter mass”