

1 Regime shifts and alternative states in the *Sarracenia*  
2 microecosystem

3 Aaron M. Ellison,<sup>1,\*</sup> Benjamin Baiser,<sup>1,2</sup> Matthew K. Lau,<sup>1</sup> and Nicholas J. Gotelli<sup>3</sup>

4 January 4, 2016

5 <sup>1</sup>Harvard Forest, Harvard University, 324 North Main Street, Petersham, Massachusetts 01366;

6 <sup>2</sup>Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida 32611;

7 <sup>3</sup>Department of Biology, University of Vermont, Burlington, Vermont 05405

8 *Keywords:* alternative states, model system, organic-matter loading, regime shifts, *Sarracenia purpurea*

9 Article type: **Article**

# **Abstract**

text - 200 words, 1 paragraph, Using sensitivity analyses we identify model parameters that most strongly control the dynamics of the pitcher plant ecosystem and show that the system can eventually overcome its hysteresis and return to an oligotrophic state once organic-matter input is stopped.

14 •

15 •

16 •

# Introduction

Regime shifts in ecological systems are defined as rapid changes in the spatial or temporal dynamics of a more-or-less stable system caused by slow, directional changes in one or more underlying state variables (??). Ecological systems in which the occurrence of regime shifts is unambiguous are uncommon, primarily because long time series of observations are required to identify both stability of each state and a breakpoint between them (?). Detailed modeling and decades of observations and experiments have led to a thorough understanding of a canonical example of an ecological regime shift: the rapid shift from oligotrophic to eutrophic states in lakes (??).

Oxygen dynamics in lakes can be described using a simple model that yields both alternative states and hysteresis in the shift between them (?):

$$\frac{dx}{dt} = a - bx + rf(x) \quad (1)$$

In this model, the observed variable  $x$  (e.g., nutrient concentration in a lake) is positively correlated with state variable  $a$  (e.g., rate of nutrient loading) and negatively correlated with state variable  $b$  (e.g., rate of nutrient removal). A positive feedback loop,  $rf(x)$  (in the lake model, this is the rate of nutrient recycling between the lake sediments and water column). If  $r > 0$  and  $\{rf(x)\} > b$ , there will be more than one equilibrium point (i.e., stable state); the function  $f(x)$  determines the shape of the switch between the states and the degree of hysteresis (?).

Models of lake ecosystems and their food webs, and associated empirical data also have revealed that returning lakes to the more “desired” oligotrophic state can be very slow—on the order of decades to centuries—(ref: Contamin and Ellison; others from Carpenter group) and depends not only on slowing or reversing the directional changes in underlying state variables but also on the internal feedback dynamics of the system (i.e., the dynamics of  $f(x)$  in Equation 1). Other systems, including fisheries (ref: Carpenter PNAS) and ... (ref; XXX et al. PLoS One) have provided some support for these model results (ref: XXX et al. PLoS One), both in terms of dynamics and duration.

In the last several years, many researchers have suggested that a wide range of ecological systems are poised to “tip” into new regimes (refs...), or even that we are approaching a planetary tipping point. (ref: XXX, but see YYY). If, as in lakes, either changes in the underlying state variables causing these regime shifts or the regime shifts themselves span decades, ecological insights into their causes and consequences will accrue relatively slowly. More rapid progress both in understanding the mechanisms driving ecological regime shifts and in determining how to reverse them requires well-understood model systems that can be

experimentally manipulated over much shorter time scales.

Recently, we have shown experimentally that organic-matter loading (i.e., prey addition) can cause a shift from oligotrophic to eutrophic conditions in naturally-occurring microecosystems: the water and bacteria-filled modified leaves of the northern (or purple) pitcher plant, *Sarracenia purpurea* (ref: Sirota). Because bacteria that reproduce rapidly drive the nutrient cycling dynamics of the *Sarracenia* microecosystem (ref: Butler et al. 2008), the system states change in days rather than years or decades. In addition to introducing a new experimental system for regime shifts and alternative states, Sirota et al. (2013) sketched a model that, in extreme cases, could lead to a rapid (regime) shift from oligotrophic to eutrophic conditions in pitcher-plant leaves. Here, we develop fully the model of the *Sarracenia* microecosystem, introduce more realism into the underlying environmental drivers of the model, use sensitivity analysis to identify the parameters that most strongly control the dynamics of the system, and show that the system eventually can overcome its hysteresis and return to an oligotrophic state once organic-matter input is stopped.

## The *Sarracenia* microecosystem

Sketch of the biology of the system

## The Model

The detritus-based pitcher-plant food web forms in the small pools of water that accumulate in living and photosynthesizing modified leaves (“pitchers”) of *Sarracenia purpurea*. Thus, our model of this food web includes feedbacks between the food web and the plant itself (?). The full model is a pair of coupled differential equations:

$$x_{t+1} = \underbrace{a_t A_t}_{\text{Photosynthesis}} - \underbrace{\left\{ m + a_t \left[ \frac{w_{t-1}}{K_w + w_{t-1}} \right] \right\}}_{f(w,x,t)} + \underbrace{D_t(x_t)}_{g(x,t)} \quad (2a)$$

$$a_{t+1} = a_t \times \left\{ \frac{a'_{max} - a'_{min}}{1 + \exp(-s \cdot n_t - d)} + a'_{min} \right\} \quad (2b)$$

In this model, the pitcher-plant fluid is oxygenated by photosynthesis ( $A_t$ ), with  $A_{max}$  augmented by prey mineralization ( $a_t$ ), as well as some diffusion from the atmosphere ( $g(x,t)$ ). Oxygen is lost from the fluid as

Table 1: Terms, units, and interpretation for the model of oxygen dynamics in pitcher-plant fluid

Term	Units	Interpretation
$a_t$	mg/L	Maximum amount of oxygen infused from the plant to the fluid at mid-day
$a'_{min}, a'_{max}$	mg/L	minimum or maximum possible augmentation of photosynthesis as a result of nutrient fertilization from organic-matter mineralization by the food web
$d$	mg	inflection point of the sigmoidal curve relating nutrient mineralization to photosynthetic augmentation
$D_t$	mg/L	Diffusion rate of oxygen from atmosphere into the pitcher fluid
$f$	1/ $t$	constant adjusting sine wave of diurnal PAR for frequency of measurements
$K_w$	mg/min	half-saturation constant for prey consumption
$m$	mg/L	amount of oxygen used for basal metabolism (respiration) of bacteria
$n_t$	mg/L	quantity of nutrients mineralized by decomposition; a function of $w_t$ and $x_t$
$s$	dimensionless	steppens of the sigmoidal curve relating nutrient mineralization to photosynthetic augmentation
$t$	minutes	time (frequency of model iteration)
$w_t$	mg	mass of prey remaining at time $t$
$x_t$	mg/L	[O <sub>2</sub> ]: concentration of oxygen in the pitcher fluid

prey is decomposed by the food web ( $f(w, x, t)$ ). A full list of model parameters is given in Table 1.

## Photosynthesis

In the absence of the foodweb, baseline oxygen concentration [O<sub>2</sub>] in the pitcher water, is determined by diurnal photosynthesis: pitchers take up CO<sub>2</sub> from the air or water and release O<sub>2</sub> back into it (???). Photosynthesis is first modeled as  $A = A_{max}[1 - \exp(-A_{qe}[\text{PAR} - \text{LCP}])]$  (after ?), where PAR is photosynthetically active radiation, which normally ranges from 0 - 2500  $\mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  and which is modeled as a truncated diurnal sine-wave (PAR = 0 at night, a sine-wave during the day: **1**);  $A$  is photosynthetic rate (in  $\mu\text{mol CO}_2 \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ );  $A_{max}$  is the maximum photosynthetic rate of an pitcher plant that has not captured or processed any prey ( $\bar{x} \approx 4 \mu\text{mol CO}_2 \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; ??);  $A_{qe}$  is the quantum yield ( $\approx 0.3$ : the initial slope, at low light, of the  $A$  vs. PAR curve); and LCP is the light compensation point: the x-intercept when  $A = 0$ ,  $\approx 20 \text{ mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ .

Since we are interested in oxygen dynamics in volumes of water, we convert  $A$  from  $\mu\text{mol CO}_2 \cdot \text{m}^{-2} \cdot \text{s}^{-1}$  to mg O<sub>2</sub> / L by noting that there is a 1:1 relationship between  $\mu\text{mol CO}_2$  and  $\mu\text{mol O}_2$  in photosynthesis; that 1  $\mu\text{mol O}_2 = 32 \mu\text{g O}_2 \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ; and that the specific leaf area of *S. purpurea*  $\approx 70 \text{ cm}^2/\text{g}$  (?). Thus, O<sub>2</sub> production  $\approx 0.21 \mu\text{g O}_2 \cdot \text{g leaf}^{-1} \cdot \text{s}^{-1}$ . The average 5-ml leaf has a surface area of 40 cm<sup>2</sup> and weighs

Figure 1: Four-part figure. (a) truncated sine wave of PAR; (b) observed PAR; (c)  $A$  as a function of PAR w/o food web; (d)  $A$  as a function of PAR with mineralization augmentation

0.5 g, and a 1-L leaf would weigh 10g. Thus,  $O_2$  production from photosynthesis  $\approx 2.1 \text{ mg } O_2 \cdot L^{-1} \cdot s^{-1}$ .

It follows that our functions of PAR and  $A$ , taking into account LCP, are:

$$PAR = c \sin(2\pi ft) \quad (3a)$$

$$A = \begin{cases} a \cdot A_{max} [1 - \exp(-0.3[PAR - 20])] & : PAR \geq 20 \\ 0 & : PAR < 20 \end{cases} \quad (3b)$$

In Eq. 3a, the amplitude  $c$  is maximum PAR at mid-day; the constant  $f$  adjusts for the frequency of measurements. Since our model updates every minute,  $f = 1/1440$  (the reciprocal of the number of minutes per day). Without loss of generality, our model runs start at sunrise ( $t = 1$ ), day-length = 12 hours = 720 minutes, and  $PAR(t)$  is truncated = 0 for  $t \in \{721, 1440\}$ . In Eq. 3b,  $a$  is the amount that photosynthetic rate can be increased as a result of nutrients provided to the plant as prey is mineralized by the food web (1).

## Respiration

Biological oxygen demand (BOD) of bacteria as they decompose prey rapidly depletes  $[O_2]$  in the pitcher fluid. Although different  $[O_2]$  may favor different numbers of either aerobic or anaerobic bacteria in the pitcher fluid, the efficiency of organic-matter decomposition by either type of bacteria is roughly equal (?), so we assume sufficient numbers of bacteria to decompose prey at a fixed, negative exponential rate:

$$w(t) = ae^{-b[w_0t]} \quad (4)$$

Biologically, we take this to mean that easily digested parts of insect prey, such as fat bodies, are processed quickly, whereas the more recalcitrant proteins and chitins break down more slowly (?). Field observations have shown that the soft parts of a single 75- $\mu g$  wasp can be consumed completely over a 48-hour period in a pitcher with 5 ml of fluid. Thus, the baseline parameters for Equation 4 were set at  $a = 20 \text{ mg}$  and  $b = 4 \text{ mg} \cdot \text{mg}^{-1} \cdot \text{d}^{-1}$ . Because we fix the decomposition rate of given mass of prey, to maintain a 48-hour decomposition time of a single 75- $\mu g$  wasp while varying  $a$  and  $b$  in Equation 4, it is necessary to fix  $\frac{a}{b} = 5$ . Finally, because *S. purpurea* produces digestive enzymes such as proteases and chitinases only for a short period of time early in leaf growth (?), we ignore their effects here.

Oxygen used up by the bacteria in prey consumption ( $= 1 - \text{BOD}$ ) was modeled as a saturating function of the remaining prey. The proportion of the prey remaining  $= 1 -$  the amount of prey decomposed, which is a function of the maximum amount of oxygen at mid-day ( $a$  in Equation 2a, iteratively augmented by  $a'$  from Equation 2b and as detailed below in Equations 6, 7a); the mass of prey ( $w$ ); a half-saturation constant defining the prey-consumption curve ( $K_w$ ) that determines how much prey is left over each “day” of the iterated model; and the amount of oxygen needed by the bacteria for basal metabolism. Hence:

$$\text{O}_2 \text{ lost}(t) = f(w, x, t) = m + a \left[ \frac{w(t-1)}{K_w + w(t-1)} \right] \quad (5)$$

In the sensitivity analysis,  $m$  in Equation 5 was fixed  $= 1$ , whereas  $K_w$  varied.

## Diffusion and oxygenation

$[\text{O}_2]$  increases in the pitcher fluid in three ways. Some oxygen diffuses from the atmosphere into the pitcher fluid, but because this happens only at the surface of the pitcher fluid, and the “mouth” of the pitcher is at least an order of magnitude smaller than the surface area of the pitcher itself (?); we ignore this term and focus on re-oxygenation through baseline oxygen production and prey-augmented photosynthesis (Equation 3b). The latter results from a positive feedback loop in Equation 2b that links prey mineralization to the uptake of mineralized nutrients by the plant, and the subsequent usage of these nutrients to increase photosynthetic rate of the pitcher (?). Ants and wasps, the most common insect prey of pitcher plants (?), are  $\approx 50\%$  C, have a C:N ratio of 6:1, and N:P:K ratios of  $\approx 12:1.5:0.9$  (?). As the prey are mineralized, the nutrients that are released, especially  $\text{NH}_4$ (?) and P (as  $^{32}\text{P}$ : ?), are absorbed rapidly by the pitcher. Photosynthesis by *Sarracenia* is limited by both N and P (and stoichiometrically by N: (??)), and photosynthetic rates of pitcher plants significantly increase following N additions (?). We model these two processes – nutrient release and augmentation of photosynthesis – with a pair of equations.

First, nutrient release following bacterial mineralization is a function of prey mass ( $w$ ) and available oxygen ( $x$ ) used by bacteria to break down and mineralize the prey:

$$n(t) = \frac{w(t)x(t)}{c} \quad (6)$$

where  $c$  is a scaling constant (we set  $c = 100$ ).

Nutrients absorbed by pitchers and not stored for future use (?) could be used to make additional enzymes for photosynthesis (?), and we model uptake as a sigmoidal (saturating) relationship between



Table 2: Values of parameters used in the sensitivity analyses. Note that the values of  $a$  and  $b$  must change in concert to maintain a constant ratio of 5 so that prey decomposition is complete within 48 hours

Variable	values
Prey added ( $w_0$ )	0, 1, 5 mg/day
$a$	5, 10, 20, 40 mg
$b$	1, 2, 4, 8 mg · mg <sup>-1</sup> · d <sup>-1</sup>
$K_w$	0.001, 0.01, 0.1 mg/min
$d$	-1, -2, -3, -4, -5 mg

129 additional nutrients absorbed and augmentation of the peak rate of photosynthesis ( $a$  in Equation 2a):

$$a' = \frac{a'_{max} - a'_{min}}{1 + \exp(-[sn(t) - d])} + a'_{min} \quad (7a)$$

$$a(t+1) = a(t) \times a'(t) \quad (7b)$$

130 In Equation 7a,  $a'_{min}$  is the minimum possible augmentation of photosynthesis, which we set = 0;  $a'_{max}$  is  
 131 the maximum possible augmentation, which we set = 2;  $s$  is the steepness of the increase (set = 10); and  
 132  $d$  is the inflection point of the curve (= 0.5). Augmentation evolves as the day's leftover prey (i.e., that  
 133 not completely broken down in a given day) accumulates (within the  $n(t)$  term), and is mineralized on  
 134 subsequent days. We note that modeling this as discrete day-to-day carry-over is an artefact of iterations  
 135 in computer models. In real pitchers, decomposition is more continuous. Similarly, the model updates the  
 136 value of  $a'$  once each “day” using Equation 7b because conversion of nutrients to new photosynthetic enzymes  
 137 is assumed to occur slowly relatively to bacterial decomposition itself.

## 138 Sensitivity analysis

139 Five paramters were varied over a wide range of values to explore the sensitivity of the model to prey input  
 140 (mass added each day); the decomposition rate ( $a$  and  $b$  in Equation 4), the half-saturation constant for prey  
 141 consumption ( $K_w$  in Equation 5) and the inflection point  $d$  for augmentation of photosynthesis in Equation  
 142 7a. The range of values for each of these parameters is given in Table 2.

143 Models were run for each of the possible combinations of  $\{w_0, a, b, K_w, d\}$ —note that the joint set of  $a$   
 144 and  $b$  includes only 4 combinations:  $\{(5,1), (10,2), (20,4), (40,8)\}$ —for a total of 180 model runs.

# Results

## Model dynamics

Illustrative examples (?spark charts or some subset of them) illustrating the range of possible outcomes - controls, intermediate dynamics, collapse, recovery.

This should include time-series plots as well as phase-space plots. Examples in Sirota et al. PNAS supplement, but I think we can do better/differently here.

- Different state = change in O<sub>2</sub> production where the non-transient maximum values are different from control

## Sensitivity analysis

Which parameters drive the dynamics? Multivariate plots, additional analyses, ...

- Transient = returning to the maximum
- Prey: increases crashes
- D: transient when  $D = -1$ , stable or crashing otherwise
- K: increases the buffering of the system, increases return rates, increases the number of stable states
- ab: alters transient return rates, alters stable point

# Discussion

1. Why we need model systems for regime shifts and alternative states
2. Similarities and differences between pitcher plants, lakes, and other ecological systems in which regime shifts have been identified and explored
3. Insights from pitcher plant system and model that could be applied to other systems

# Literature Cited

## References

- Baiser, B., R. Ardesiri, and A. M. Ellison. 2011. Species richness and trophic diversity increase decomposition in a co-evolved food web. *PLoS One* 6:e20672.
- Bestelmeyer, B. T., A. M. Ellison, W. R. Fraser, K. B. Gorman, S. J. Holbrook, C. M. Laney, M. D. Ohman, D. P. C. Peters, F. C. Pilsbury, A. Rassweiler, R. J. Schmitt, and S. Sharma. 2011. Detecting and managing abrupt transitions in ecological systems. *Ecosphere* 2:129.
- Bradshaw, W., and R. Creelman. 1984. Mutualism between the carnivorous purple pitcher plant *Sarracenia purpurea* and its inhabitants. *American Midland Naturalist* 112:294–304.
- Butler, J., and A. Ellison. 2007. Nitrogen cycling dynamics in the carnivorous pitcher plant, *Sarracenia purpurea*. *Functional Ecology* 21:835–843.
- Cameron, C., G. Donald, and C. Paterson. 1977. Oxygen - fauna relationships in the pitcher plant *Sarracenia purpurea* l. with reference to the chironomid *Metriocnemus knabi* coq. *Canadian Journal of Zoology* 55:2018–2023.
- Carpenter, S., and W. Brock. 2006. Rising variance: a leading indicator of ecological transition. *Ecology Letters* 9:311–318.
- Carpenter, S., J. Cole, M. Pace, R. Batt, W. Brock, T. Cline, J. Coloso, J. Hodgson, J. Kitchell, D. Seekell, L. Smith, and B. Weidel. 2011. Early warnings of regime shifts: a whole-ecosystem experiment. *Science* 332:1079–1082.
- Ellison, A. 2006. Nutrient limitation and stoichiometry of carnivorous plants. *Plant Biology* 8:740–747.
- Ellison, A., and N. Gotelli. 2002. Nitrogen availability alters the expression of carnivory in the northern pitcher plant *Sarracenia purpurea*. *Proceedings of the National Academy of Sciences, USA* 99:4409–4412.
- . 2009. Energetics and the evolution of carnivorous plants - darwin's "most wonderful plants in the world". *Journal of Experimental Botany* 60:19–42.
- Farnsworth, E., and A. Ellison. 2008. Prey availability directly affects physiology, growth, nutrient allocation and scaling relationships among leaf traits in 10 carnivorous plant species. *Journal of Ecology* 96:213–221.

192 Gallie, D., and S. Chang. 1997. Signal transduction in the carnivorous plant *Sarracenia purpurea* - regulation  
193 of secretory hydrolase expression during development and in response to resources. *Plant Physiology*  
194 115:1461–1471.

195 Givnish, T., E. Burkhardt, R. Happel, and J. Weintraub. 1984. Carnivory in the bromeliad *Brocchinia*  
196 *reducta*, with a cost/ benefit model for the general restriction of carnivorous plants to sunny, moist nutrient-  
197 poor habitats. *American Naturalist* 124:479–497.

198 Murphy, M., and S. Condon. 1984. Comparison of aerobic and anaerobic growth of *Lactobacillus plantarum*  
199 in a glucose medium. *Archives of Microbiology* 138:49–53.

200 Peek, M., E. Russek-Cohen, D. Wait, and I. Forseth. 2002. Physiological response curve analysis using  
201 nonlinear mixed models. *Oecologia (Berl.)* 132:175–180.

202 Plummer, G., and J. Kethley. 1964. Foliar absorption of amino acids, peptides, and other nutrients by the  
203 pitcher plant, *Sarracenia flava*. *Botanical Gazette* 125:245–260.

204 Scheffer, M., J. Bascompte, W. Brock, V. Brovkin, S. Carpenter, V. Dakos, H. Held, E. van Nes, M. Rietkerk,  
205 and G. Sugihara. 2009. Early-warning signals for critical transitions. *Nature* 461:53–59.

206 Scheffer, M., S. Carpenter, J. Foley, C. Folke, and B. Walker. 2001. Catastrophic shifts in ecosystems. *Nature*  
207 413:591–596.

208 Sirota, J., B. Baiser, N. J. Gotelli, and A. M. Ellison. 2013. Organic-matter loading determines regime shifts  
209 and alternative states in an aquatic ecosystem. *Proceedings of the National Academy of Sciences, USA*  
210 110:7742–7747.

211 Small, E. 1972. Photosynthetic rates in relation to nitrogen recycling as an adaptation to nutrient deficiency  
212 in peat bog plants. *Canadian Journal of Botany* 50:2227–2233.

213 Wakefield, A., N. Gotelli, S. Wittman, and A. Ellison. 2005. The effect of prey addition on nutrient stoichiometry,  
214 nutrient limitation, and morphology of the carnivorous plant *Sarracenia purpurea* (sarraceniaceae).  
215 *Ecology* 86:1737–1743.