



# Integrated Athlete Monitoring Analysis

Force Plate, GPS Workload, and  
Strength Asymmetry

Angel Huang, Anita Liu, Ashley McGowan, Aarav Desai

# Introduction

Research Question: How do neuromuscular performance, GPS-derived workload, and strength asymmetry interact over time, and can deviations from these metrics serve as indicators of readiness, fatigue, or undertraining?

Hypotheses:

- H1: Athletes who experience periods of high external load, particularly elevated acceleration load, will show short-term reductions in Jump Height and Peak Propulsive Power relative to their baseline values.
- H2: Athletes with higher MaxForce asymmetry will display greater variability in CMJ performance and may demonstrate different load-response patterns compared to their more symmetrical teammates.



# The Gap

Although many teams collect force plate, GPS, and strength data, the literature shows:

- CMJ is useful for fatigue monitoring but often analyzed alone.
- GPS workload (especially accelerations) is strongly associated with neuromuscular stress but rarely paired with force-plate data.
- Strength asymmetry is studied in isolation and not connected to longitudinal workload or CMJ decline.

There is a clear gap: Teams rarely integrate all three systems into a unified readiness or monitoring framework. Our project directly addresses this gap.



# Selected Metrics

1

## Force Plate (Hawkins)

Jump Height (m)  
Peak Propulsive Power (W)



2

## GPS Workload (Kinexon)

Total Distance  
Accumulated Acceleration Load



3

## Strength (Vald)

MaxForce Left  
MaxForce Right  
Computes inter-limb asymmetry

# Methods

## Data Cleaning

- Pulled from MySQL table `research\_experiment\_refactor\_test`
- Removed rows with missing player/team/metric/value
- Standardized timestamps
- Retained original team labels for authenticity
- Long-format structure preserved
- Reflects real-world testing inconsistencies

## Analytical Approach


- Literature-based flagging thresholds:
  - Inactivity: >30 days without testing
  - Performance Decline: >10% below 3-test rolling mean
  - Team Deviations:  $\pm 2$  SD from team mean
  - Asymmetry: >10% left-right difference
- Exported flagged results to `part4_flagged_athletes.csv`




# Key Findings: Inactivity Flags



- 1,286 athletes flagged
- Most athletes had last data points from 2023 or earlier
- Consistent with Part 2 findings
- Inactivity helps detect:
  - Neuromuscular fatigue
  - workload accumulation
  - Strength asymmetry



Flag Types by Metric



# Comparison to Literature

- CMJ fatigue trends only appear with regular testing
  - (Gathercole et al., 2015; McMahon et al., 2018)
- GPS workloads require consistent session by session data to interpret meaningful data
  - (Bangsbo et al., 2006; Buchheit & Simpson, 2017)
- Strength asymmetry > 10% is repeatedly linked to higher injury risk
  - (Croisier et al., 2008)





# Practical Applications (Flag System)

- 1 Inactivity flags identify training gaps
- 2 Asymmetry flags support injury-risk decisions
- 3 Decline/outlier flags guide load adjustments
- 4 Supports individualized training plans



# Recommendation for Coaches

## Monitor Key Metrics

This helps by showing specific datasets

## Use Trend Watching

See improvements over time instead of one day

## Flag Meaningful Declines

Helps catch and fix an issue before it becomes a problem

## Compare Athletes to Team Norms

Use metrics to see where the athlete matches up against the while team

## Use Testing Frequency to Improve Metrics

Ensure metrics are tested ever 2-3 weeks to avoid sudden problems

## Connect Patterns to Actions

If jump declines week to week, decrease plyometric load

# How Findings Fill In Gaps

## Real World Data

Most Data is taken in labs. This is real world game data. That means that data shows natural fatigue and variations across multiple sports

## Compare Metrics Across Different Sports

Much research isolates certain sports. Jump Height in Volleyball, Speed in track and field, etc. By putting multiple sports together, it shows how the same metrics manifest in different sports

## Establish Team and Athlete Baselines

Scientific papers report group averages but coaches need to know about THEIR team. The finding fill in the gap by building individual baselines, team averages, percent differences, etc.

# Limitations and Future Research

## Limitations

1. Inconsistent Testing Frequencies
  - a. Some athletes measures weekly, others a couple times
2. Missing or Sparse Data
  - a. Null or Zero values limit statistical strength
3. Lack of Contextual Variables
  - a. Doesn't show why metrics changed. Injury? Training Load?

## Future Research

1. Standardize and Increase Testing Frequency
  - a. Similar data collection times would help increase accuracy
2. Integrate contextual data
  - a. Explains why metrics changed
3. Expand Beyond one Institution
  - a. Comparison against other institutions would reveal universal vs contextual specific patterns

**Any  
Questions?**

# References

- Akenhead, R., Hayes, P. R., Thompson, K. G., & French, D. (2013). Diminutions of acceleration and deceleration output during professional football match play. *Journal of Science and Medicine in Sport*, 16(6), 556–561. <https://doi.org/10.1016/j.jsams.2012.12.005>
- Bangsbo, J., Mohr, M., & Krstrup, P. (2006). Physical and metabolic demands of training and match-play in the elite football player. *Journal of Sports Sciences*, 24(7), 665–674. <https://doi.org/10.1080/02640410500482529>
- Bishop, C., Read, P., McCubbine, J., & Turner, A. (2018). Inter-limb asymmetries: Understanding how to calculate the variance between limbs. *Strength and Conditioning Journal*, 40(6), 1–6. <https://doi.org/10.1519/SSC.0000000000000371>
- Buchheit, M., & Simpson, B. M. (2017). Player-tracking technology: Half-full or half-empty glass?. *International Journal of Sports Physiology and Performance*, 12(Suppl 2), S235–S241. <https://doi.org/10.1123/ijsp.2016-0499>
- Cormack, S. J., Newton, R. U., McGuigan, M. R., & Doyle, T. L. A. (2008). Reliability of measures obtained during single and repeated countermovement jumps. *International Journal of Sports Physiology and Performance*, 3(2), 131–144. <https://doi.org/10.1123/ijsp.3.2.131>
- Croisier, J.-L., Ganteaume, S., Binet, J., Genty, M., & Ferret, J.-M. (2008). Strength imbalances and prevention of hamstring injury in professional soccer players: A prospective study. *American Journal of Sports Medicine*, 36(8), 1469–1475. <https://doi.org/10.1177/0363546508316764>
- Cummins, C., Orr, R., O'Connor, H., & West, C. (2013). Global positioning systems (GPS) and microtechnology sensors in team sports: A systematic review. *Sports Medicine (Auckland, N.Z.)*, 43(10), 1025–1042. <https://doi.org/10.1007/s40279-013-0069-2>
- Gabbett, T. J. (2016). The training-injury prevention paradox: Should athletes be training smarter and harder? *British Journal of Sports Medicine*, 50(5), 273–280. <https://doi.org/10.1136/bjsports-2015-095788>



# References

- Gathercole, R. J., Sporer, B. C., Stellingwerff, T., & Sleivert, G. (2015). Alternative countermovement-jump analysis to quantify acute neuromuscular fatigue. *International Journal of Sports Physiology and Performance*, 10(1), 84–92. <https://doi.org/10.1123/ijsp.2013-0413>
- Harper, D. J., Carling, C., & Kiely, J. (2019). High-intensity acceleration and deceleration demands in elite team sports competitive match play: A systematic review and meta-analysis of observational studies. *Sports Medicine (Auckland, N.Z.)*, 49(12), 1923–1947. <https://doi.org/10.1007/s40279-019-01170-1>
- Impellizzeri, F. M., Rampinini, E., & Marcora, S. M. (2008). Physiological assessment of aerobic training in soccer. *Journal of Sports Sciences*, 23(6), 583–592. <https://doi.org/10.1080/02640410400021278>
- Loturco, I., Pereira, L. A., Cal Abad, C. C., D'Angelo, R. A., Fernandes, V., Kitamura, K., Kobal, R., & Nakamura, F. Y. (2015). Vertical and horizontal jump tests are strongly associated with competitive performance in 100-m dash events. *Journal of Strength and Conditioning Research*, 29(7), 1966–1971. <https://doi.org/10.1519/JSC.0000000000000849>
- Markovic, G., & Mikulic, P. (2010). Neuro-musculoskeletal and performance adaptations to lower-limb plyometric training. *Sports Medicine*, 40(10), 859–895. <https://doi.org/10.2165/11318370-000000000-00000>
- McMahon, J. J., Suchomel, T. J., Lake, J. P., & Comfort, P. (2018). Understanding the key phases of the countermovement jump force-time curve. *Strength & Conditioning Journal*, 40(4), 96–106. <https://doi.org/10.1519/SSC.0000000000000375>
- Silva, J. R., Nassis, G. P., & Rebelo, A. (2015). Strength training in soccer with a specific focus on highly trained players. *Sports Medicine*, 1(1), 17. <https://doi.org/10.1186/s40798-015-0006-z>
- Suchomel, T. J., Nimphius, S., & Stone, M. H. (2016). The importance of muscular strength in athletic performance. *Sports Medicine*, 46(10), 1419–1449. <https://doi.org/10.1007/s40279-016-0486-0>

