



AGENCY FOR HEALTHCARE RESEARCH AND QUALITY



MEPS-HC Prescribed Medicines (PMED) File

Rebecca Ahrnsbrak, MPS

Household PMED Data Collection



- Respondents encouraged to use memory aids
 - ▶ Pill aids bottles/containers
 - ▶ Pharmacy receipts, patient portal, or other records
- Respondents first prompted about any medicines prescribed during reported medical visits
 - ▶ For hospital stays, respondents are asked to report only drugs prescribed on discharge
 - ▶ Respondents are asked to report only prescribed medicines that they filled

Household PMED Data Collection



- **After asking about medicines prescribed during medical events, a stand-alone prescribed medicines section asks respondents about any other prescribed medicines obtained at any pharmacy:**
 - ▶ **Any new prescription medicines or refills**
 - ▶ **Any prescribed medicines taken only as needed**
 - ▶ **Any diabetic equipment or supplies**

Pharmacy Data Collection

- **Requires written authorization from sample member to request their pharmacy records**
- **Asks pharmacies for “patient profile” or information on patient’s drug fills**
- **Used primarily for expenditure information and detailed drug characteristics (form, strength, quantity, etc.)**
- **Used as imputation donor pool for those without PC data and those with incomplete PC data**

Elements Collected: HC and PC

Household Component

- **Drug name, strength, and form**
- **Number of fills in round**
- **Associated condition(s)**
- **When first used**
- **Pharmacy information**
- **Authorization to contact pharmacies**

Pharmacy Component

- **Drug name**
- **Detailed drug characteristics (strength, form, days supplied, etc.)**
- **National Drug Code (NDC)**
- **Payers**
- **Amount paid by each payer (including out of pocket)**

Editing HC and PC Data

- **The HC and PC data do not share a common identifier**
 - ▶ **A Generic Product Indicator (GPI) is assigned to both HC and PC records to facilitate matching**
- **Before HC and PC data are combined, editing and imputation are done separately on each file**
 - ▶ **Review outliers – flag and impute when implausible**
 - ▶ **Price benchmarking for PC data**
 - ▶ **Impute missing data elements in each file**

Combining HC and PC Data

- Iterative and progressively less restrictive matching process
 - ▶ Matching is done at the person-round-drug level
 - ▶ For those with PC data, attempts to match within the person's own PC data are attempted first
 - ▶ For those with no or incomplete PC data, a donor PC record that matches as closely as possible is imputed
 - ▶ Inexact matches are permitted when exact match attempts fail
- Utilization (fill records) are based only on household-reported data
 - ▶ MEPS does not create new PMED records based on PC data
- Post-match editing is performed on the combined data

PMED Public Use File (PUF)



- **The PMED PUF is unfolded to the acquisition (fill or refill) level**
 - ▶ Each record represents a unique fill or refill of a prescribed medicine for an individual in the data year
 - ▶ Not everyone on the FYC will have PMED records
 - ▶ If people have multiple drug fills, they will have multiple PMED records
- **Rarely used drugs are masked (to -15) to protect respondent confidentiality**
 - ▶ In these cases, Multum therapeutic class information replaces the medication name in RXDRGNAM
- **Can be linked to:**
 - ▶ FYC (person-level) PUF by DUPERSID
 - ▶ Medical Conditions PUF by CLNK PUF

PMED PUF Structure

- **DUPERSID** – Unique person identifier
- **RXRECIDX** – Unique identifier for each record (fill) on the file
- **DRUGIDX** – Unique drug identifier for a given person
- **LINKIDX** – Unique person-round-drug identifier for linking to other files

| DUPERSID | DRUGIDX | RXRECIDX | PURC HRD | LINKIDX | RXDRGNAM |
|------------|---------------|---------------------|-------------|------------------|--------------|
| 2510001101 | 2510001101001 | 2510001101001103001 | 1 | 2510001101001103 | ATORVASTATIN |
| 2510001101 | 2510001101001 | 2510001101001103002 | 1 | 2510001101001103 | ATORVASTATIN |
| 2510001101 | 2510001101001 | 2510001101001203003 | 2 | 2510001101001203 | ATORVASTATIN |
| 2510001101 | 2510001101002 | 2510001101002203004 | 2 | 2510001101002203 | AMOXICILLIN |
| 1 person | 2 drugs | 4 fills | 3 LINKIDXs | | 2 drugs |


Additional PMED File Contents

- Medication name
 - ▶ RXDRGNAM: standardized generic drug name
 - ▶ RXNAME: pharmacy-reported drug name
- National Drug Code (NDC)
- Total payment
- Amount paid by payer type
- Pharmacy types
- Quantity dispensed
- Days supplied
- Strength
- Form
- Purchase round
- Flag for diabetic equipment or supplies
- Month and year person started taking drug
- Whether person has PC data
- Imputation information
- Multum therapeutic classes and pregnancy category
- Person-level weight and design variables

Multum Therapeutic Classes

- **Classifies drugs into therapeutic classes, subclasses, and sub-sub-classes**
 - ▶ **TCn** - assigns a drug to one or more therapeutic/chemical categories
 - Can have up to three therapeutic classes per drug
 - ▶ **TCnSn** - assigns one or more therapeutic subclasses to a given therapeutic class
 - ▶ **TCnSn_n** - assigns one or more sub-sub-classes to a given therapeutic subclass
- **Classification scheme can change over time**
 - ▶ Check documentation for each year
- **For 1996-2013, use the Multum Lexicon Addendum Files**

Example: Multum Therapeutic Classes



Finding PMED and Linking Files

[:: What's New](#)
[:: Mailing List](#)
[:: Discussion Forum](#)
[:: Participants' Corner](#)

☐ [Household Component Full-Year files](#) ⓘ
Expenditure and utilization data for the calendar year from several rounds of data collection.
☐ [Full-Year Consolidated Data files](#) **FYC (person-level) Files**
☐ [Full-Year Population Characteristics files](#)
☐ [Medical Conditions files](#)
☐ [Risk Adjustment Scores files](#)
☐ [Employment Variables file](#)
☐ [Jobs files](#)
☐ [Person Round Plan files](#)
☐ [Longitudinal Data files](#)
☐ [Supplemental Variables files \(1996-2000\)](#)
☐ [Health Insurance Plan Abstraction file \(1996\)](#)
☐ [Long Term Care file \(1998\)](#)
☐ [Household Component Event files](#) ⓘ
Data for the calendar year on unique household-reported medical events.
☐ [Prescribed Medicines files](#) **PMED Files**
☐ [Dental Visits files](#)
☐ [Other Medical Expenses files](#)
☐ [Hospital Inpatient Stays files](#)
☐ [Emergency Room Visits files](#)
☐ [Outpatient Visits files](#)
☐ [Office-Based Medical Provider Visits files](#)
☐ [Home Health files](#)
☐ [Appendix to MEPS Event files](#) **CLNK File**

Linking PMEDs to Medical Conditions

PMED File

| DUPERSID | DRUGIDX | PURCHRD | LINKIDX | RXRECIDX |
|------------|---------------|---------|------------------|---------------------|
| 2510001101 | 2510001101001 | 1 | 2510001101001103 | 2510001101001103001 |
| 2510001101 | 2510001101001 | 1 | 2510001101001103 | 2510001101001103002 |
| 2510001101 | 2510001101001 | 2 | 2510001101001203 | 2510001101001203003 |

CLNK File

| DUPERSID | CONDIDX | EVNTIDX | EVENT YPE |
|------------|---------------|------------------|--------------|
| 2510001101 | 2510001101003 | 2510001101001103 | 8 |
| 2510001101 | 2510001101003 | 2510001101001203 | 8 |

Medical Conditions File

| DUPERSID | CONDIDX | ICD10CDX |
|------------|---------------|----------|
| 2510001101 | 2510001101003 | Z13 |

- **EVENTYPE = 8 on CLNK file indicates PMED event**
- **Merge LINKIDX from the PMED file to EVNTIDX on the CLNK file**

Linking: Examples

- **Analyze expenditures for prescribed medicines used to treat hypertension**
 - ▶ Use CLNK PUF to link PMED PUF to conditions PUF
- **Analyze antidepressant use by gender**
 - ▶ Link FYC and PMED PUFs by DUPERSID

Caveats and Limitations

- **Potential underreporting of drugs**
- **MEPS only measures prescribed medicines obtained in an outpatient setting (retail and mail-order pharmacies)**
 - ▶ **It does not measure drugs administered in a medical office, clinic, or inpatient setting**
- **MEPS does not measure if drugs were actually taken (just those that were filled)**
- **Fills are not standardized for days supplied**

Caveats and Limitations


- **MEPS is not well suited to studying prescribers**
- **PMED expenditures do not include rebates between manufacturers and pharmacy benefit managers (PBMs) or government programs**
- **Pharmacy type variables are those reportedly used by the person in the purchase round and any prior rounds**
 - ▶ **Not unique to a specific drug or fill**
 - ▶ **Definitions of pharmacy types are fuzzy to respondents**

Interpreting Trends

- **Be cautious interpreting year-over-year changes**
 - ▶ **Policy changes**
 - ▶ **Changes in PMED data processing procedures (especially 2007-2009 and 2017)**
 - ▶ **MEPS design changes (especially 2013 and 2018)**
- **Read the PMED documentation for each year being analyzed**
- **Consider techniques to stabilize or smooth trends**

Example of the Trends

Total expenditures (\$) in millions by prescribed drug, United States, 2008 to 2021



Resources



- **Annual public use file documentation**
- Outpatient Prescription Drugs: Data Collection and Editing in the 2021 Medical Expenditure Panel Survey
- MEPS-HC online data tools
- MEPS GitHub (includes example code in R, SAS, and Stata)
- The Impacts of the COVID-19 Pandemic on the Medical Expenditure Panel Survey
- Pre-COVID-19 Retail Use and Expenditures for Drugs That Were Subsequently Used to Treat COVID-19

Thank you!



Rebecca.Ahrnsbrak@ahrq.hhs.gov