

# Homework 4

April 29, 2021

$$[B : T(u_1) \dots T(u_n)]$$

$$[B : L(u_1) L(u_2) L(u_3)]$$

1. Let  $E = \{u_1, u_2, u_3\}$  and  $F = \{b_1, b_2\}$ , where

$$\begin{aligned} D(u) &= 0 = [1 \ x \ x^2] \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \\ D(x) &= 1 = [1 \ x \ x^2] \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \\ D(x^2) &= 2x = [1 \ x \ x^2] \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} \end{aligned}$$

$V:$

$$u_1 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, u_2 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, u_3 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$$

and

$W:$

$$b_1 = (1, -1)^T, \quad b_2 = (2, -1)^T$$

For each of the following linear transformations  $L$  from  $\mathbb{R}^3$  into  $\mathbb{R}^2$ , find the matrix representing  $L$  with respect to the ordered bases  $E$  and  $F$ :

(i)  $L(x) = \begin{pmatrix} x_3 \\ x_1 \end{pmatrix}$

$$\begin{bmatrix} 1 & 2 & -1 & 1 & 1 \\ -1 & -1 & 1 & 1 & -1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -1 & 1 & 1 \\ 0 & 1 & 0 & 2 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -1 & -3 & 1 \\ 0 & 1 & 0 & 2 & 0 \end{bmatrix}$$

(ii)  $L(x) = \begin{pmatrix} x_1 + x_2 \\ x_1 - x_3 \end{pmatrix}$

$$\begin{bmatrix} -5 & -3 & 4 \\ 3 & 3 & -2 \end{bmatrix}$$

(iii)  $L(x) = \begin{pmatrix} 2x_2 \\ -x_1 \end{pmatrix}$

$$\begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 3 \end{bmatrix}$$

2. Let  $D$  be the differentiation operator on  $P_2(\mathbb{R})$ . Find the matrix  $B$  representing  $D$  with respect to  $[1, x, x^2]$ , the matrix  $A$  representing  $D$  with respect to  $[1, 2x, 4x^2 - 2]$ , and the nonsingular matrix  $S$  such that  $B = S^{-1}AS$ .

$$A = S^{-1}BS$$

$$B = SAS^{-1} \quad A = SBS^{-1}$$

3. Suppose  $V$  and  $W$  are finite-dimensional and  $T \in \mathcal{L}(V, W)$ . Prove that  $\dim T(V) = 1$  if and only if there exist a basis of  $V$  and a basis of  $W$  such that with respect to these bases, all entries of the matrix representation  $M(T)$  equal 1.

4. Suppose  $T \in \mathcal{L}(U, V)$  and  $S \in \mathcal{L}(V, W)$  are both invertible linear transformation. Prove that  $ST \in \mathcal{L}(U, W)$  is invertible and that  $(ST)^{-1} = T^{-1}S^{-1}$ .

$$S^{-1}$$

$$[1, 2x, 4x^2 - 2] = [1, x, x^2] S^{-1}$$

$$= [1, x, x^2] \begin{bmatrix} 1 & 0 & -2 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$


5. (a) Suppose  $V$  is finite-dimensional and  $T \in \mathcal{L}(V)$ . Prove that the following statements are equivalent:

- (i)  $T$  is invertible;
- (ii)  $T$  is injective;
- (iii)  $T$  is surjective.

(b) Suppose  $V$  is finite-dimensional,  $U$  is a subspace of  $V$ , and  $S \in \mathcal{L}(U, V)$ . Prove there exists an invertible operator  $T \in \mathcal{L}(V)$  such that  $Tu = Su$  for every  $u \in U$  if and only if  $S$  is injective.

(c) Suppose  $W$  is finite-dimensional and  $T_1, T_2 \in \mathcal{L}(V, W)$ . Prove that  $\ker(T_1) = \ker(T_2)$  if and only if there exists an invertible operator  $S \in \mathcal{L}(W)$  such that  $T_1 = ST_2$ .  $T_1(v) = S T_2(v)$

(d) Suppose  $V$  is finite-dimensional and  $T_1, T_2 \in \mathcal{L}(V, W)$ . Prove that  $T_1(V) = T_2(V)$  if and only if there exists an invertible operator  $S \in \mathcal{L}(W)$  such that  $T_1 = T_2 S$ .



$\{v_1, \dots, v_r\}$  is a basis of  $\ker(T_1)$  and  $\ker(T_2)$   
 $\{v_1, \dots, v_r, v_{r+1}, \dots, v_n\}$  is a basis of  $V$ .

