

Homework 4

April 29, 2021

1. Let $E = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ and $F = \{\mathbf{b}_1, \mathbf{b}_2\}$, where

$$\mathbf{u}_1 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \mathbf{u}_2 = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}, \mathbf{u}_3 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$$

and

$$\mathbf{b}_1 = (1, -1)^T, \quad \mathbf{b}_2 = (2, -1)^T$$

For each of the following linear transformations L from \mathbb{R}^3 into \mathbb{R}^2 , find the matrix representing L with respect to the ordered bases E and F :

- (i) $L(\mathbf{x}) = \begin{pmatrix} x_3 \\ x_1 \end{pmatrix}$
 - (ii) $L(\mathbf{x}) = \begin{pmatrix} x_1 + x_2 \\ x_1 - x_3 \end{pmatrix}$
 - (iii) $L(\mathbf{x}) = \begin{pmatrix} 2x_2 \\ -x_1 \end{pmatrix}$
2. Let D be the differentiation operator on $P_2(\mathbb{R})$. Find the matrix B representing D with respect to $[1, x, x^2]$, the matrix A representing D with respect to $[1, 2x, 4x^2 - 2]$, and the nonsingular matrix S such that $B = S^{-1}AS$.
3. Suppose V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$. Prove that $\dim T(V) = 1$ if and only if there exist a basis of V and a basis of W such that with respect to these bases, all entries of the matrix representation $\mathcal{M}(T)$ equal 1.
4. Suppose $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$ are both invertible linear transformations. Prove that $ST \in \mathcal{L}(U, W)$ is invertible and that $(ST)^{-1} = T^{-1}S^{-1}$.

5. (a) Suppose V is finite-dimensional and $T \in \mathcal{L}(V)$. Prove that the following statements are equivalent:
- (i) T is invertible;
 - (ii) T is injective;
 - (iii) T is surjective.
- (b) Suppose V is finite-dimensional, U is a subspace of V , and $S \in \mathcal{L}(U, V)$. Prove there exists an invertible operator $T \in \mathcal{L}(V)$ such that $Tu = Su$ for every $u \in U$ if and only if S is injective.
- (c) Suppose W is finite-dimensional and $T_1, T_2 \in \mathcal{L}(V, W)$. Prove that $\ker(T_1) = \ker(T_2)$ if and only if there exists an invertible operator $S \in \mathcal{L}(W)$ such that $T_1 = ST_2$.
- (d) Suppose V is finite-dimensional and $T_1, T_2 \in \mathcal{L}(V, W)$. Prove that $T_1(V) = T_2(V)$ if and only if there exists an invertible operator $S \in \mathcal{L}(V)$ such that $T_1 = T_2S$.