

Teacher: Yanjie Li

Assignment Number: 3

Course: Linear Algebra in Control Theory Disclosure date: May 30, 2025

Problem 1

In \mathbb{R}^4 , let

$$U = \text{span}((1, 1, 0, 0), (1, 1, 1, 2)).$$

Find $u \in U$ such that ||u - (1, 2, 3, 4)|| is as small as possible.

Problem 2

Find $p \in \mathcal{P}_3(\mathbf{R})$ such that p(0) = 0, p'(0) = 0, and

$$\int_0^1 |2 + 3x - p(x)|^2 dx$$

is as small as possible.

Problem 3

Suppose $T \in \mathcal{L}(V)$ and U is a subspace of V.

- (a) Prove that if $U \subset \text{null } T$, then U is invariant under T.
- (b) Prove that if range $T \subset U$, then U is invariant under T.

Problem 4

Suppose $S, T \in \mathcal{L}(V)$ are such that ST = TS. Prove that range S is invariant under T.

Problem 5

Suppose $S, T \in \mathcal{L}(V)$ are such that ST = TS. Prove that null S is invariant under T.

Problem 6

Define $T \in \mathcal{L}(\mathbf{F}^3)$ by

$$T(z_1, z_2, z_3) = (2z_2, 0, 5z_3)$$

Find all eigenvalues and eigenvectors of T.

Problem 7

Define $T: \mathcal{P}(\mathbf{R}) \to \mathcal{P}(\mathbf{R})$ by Tp = p'. Find all eigenvalues and eigenvectors of T.

Problem 8

Suppose $T \in \mathcal{L}(V)$. Suppose $S \in \mathcal{L}(V)$ is invertible. (a) Prove that T and $S^{-1}TS$ have the same eigenvalues. (b) What is the relationship between the eigenvectors of T and the eigenvectors of $S^{-1}TS$?

Problem 9

Find all eigenvalues and eigenvectors of the backward shift operator $T \in \mathcal{L}(\mathbf{F}^{\infty})$ defined by

$$T(z_1, z_2, z_3, \ldots) = (z_2, z_3, \ldots).$$

Problem 10

If A is a matrix with $m \times n$ dimension, please show that A^TA and AA^T have the same nonzero eigenvalues.

Pay Attention

- a) Mark your class number, student number and name on the homework.
- b) Try to write your homework on A4 size paper.
- c) Please hand in your homework to your TA before class next Sunday (June 8). If you really cannot hand in your homework by the time mentioned above, please bring it to office C202 by yourself.