

Thesis Progress Meeting

J. Raimbault^{1,2}

¹Géographie-cités (UMR 8504 CNRS)

²LVMT (UMR-T 9403 IFSTTAR)

January 26th 2016

Achieved Work (by projects)

- Biblio/Meetings/Organisation [0.7w]
- Conference [0.7w]
- Reading Records (*Synergetics*) [0.2w]
- Monitorat [1,3w]
- Cybergeog Project [1w]
- Correlated Synthetic data [3w]
- Theory construction [0.2w]
- BP Case Study / Spatial Econometrics [0,3w]

Context

Def. : *Synthetic Data* are output of generative models (and possibly inputs of models using them).

Methodology used in various fields, e.g. therapeutic evaluation [Abadie et al., 2003], territorial systems analysis [Moeckel et al., 2003, Pritchard and Miller, 2009], machine learning [Bolón-Canedo et al., 2013] or bio-informatics [Van den Bulcke et al., 2013].

Few examples at the second order : specific examples as [Ye, 2011] for discrete choices ; methods that can be interpreted this way : generation of complex networks [Newman, 2003].

Generic Method

\vec{X}_I multidimensional stochastic process, $\mathbf{X} = (X_{i,j})$ realizations.

Aim : Generate a statistical population $\tilde{\mathbf{X}} = \tilde{X}_{i,j}$ such that:

- 1 proximity to data : given a precision ε and an indicator f ,
 $\|f(\mathbf{X}) - f(\tilde{\mathbf{X}})\| < \varepsilon$
- 2 control of the estimated correlation structure : $\hat{\text{Var}} \left[(\tilde{X}_i) \right] = \Sigma R$
with R fixed.

Geographical data : Context and Objective

- In geography, generation of synthetic populations for agent-based models [Pritchard and Miller, 2009].
- Generation of spatial synthetic configuration not used (Geo. Weighted Regression [Brunsdon et al., 1998] can be interpreted this way) ; however crucial for abstract models [Schmitt, 2014]
- [Cottineau et al., 2015] recently proposed to estimate the sensitivity of spatial models of simulation to initial configuration (application to Schelling model).
- Case study : city-transportation interactions, complex to understood quantitatively [Offner, 1993, Bretagnolle, 2009] → simple model of population density and transportation network morphogenesis.

Modle

Simple coupling between

- Iterative generation of a density grid by preferential attachment/diffusion [Raimbault, 2016] calibrated on morphological objectives on european density grid.
- Heuristic network generation conditional to density :
 - Distribution of a fixed number of centers preferentially following density
 - Deterministic percolation between closest neighbors
 - Breaking of interaction potentials

$$V_{ij}(d) = \left[(1 - k_h) + k_h \cdot \left(\frac{P_i P_j}{P^2} \right)^\gamma \right] \cdot \exp \left(- \frac{d}{r_g(1 + d/d_0)} \right)$$

for a fixed number of couples N_L such that $V_{ij}(d_N)/V_{ij}(d_{ij})$ is minimal among $K \cdot N_L$ strongest euclidian potentials ($K = 5$ fixed)

- Planarization












Indicators : morphology [Le Néchet, 2015] (Moran, mean distance, entropy, hierarchy) and network (centrality, mean width, speed, diameter).

Implementation and Exploration



→ Formal and Operational coupling : modular implementation (scala/NetLogo) encapsulated by OpenMole [Reuillon et al., 2013]

→ Exploration by intensive computation on grid via OpenMole : calibration of density model alone ($\sim 1.5 \cdot 10^6$ runs) ; brutal exploration by LHS sampling for feasible correlations ($\sim 5 \cdot 10^4$ runs)



Results : Density Model alone




Results : examples of configurations



Results : examples of configurations





Results : cross-correlations




Results : feasible correlations

Mean matrices in a principal plan




Results : exemples of correlations



$$\rho[\bar{d}, \bar{c}] \simeq 0.34$$

→ gravity hierarchy more important in (1) $\gamma = 3.9, k_h = 0.7$ against
 $\gamma = 1.07, k_h = 0.25$ for (2)



$$\rho[\bar{d}, \bar{c}] \simeq -0.41$$

Applications

- 1 Calibration of the coupled model, street network data(edge effects !)
→ generation of correlated synthetic data corresponding to a given urban system → intrinsic correlations to be compared to estimated correlations between different states : non-ergodicity of urban systems [Pumain, 2012]).
- 2 Dynamical correlations in a strongly coupled model / spatio-temporal correlations in a strong spatial coupling.

Context

On Accessibility

Statistical Analysis

P. Bourguine framework for Complex Adaptive Systems

Next steps (until February 15th 2016)

- Theory exemplification, paper finalization [1w]
- Spatial Econometrics / Case study [0.5w]
- Cybergeog [0.5w]
- Wrap everything within a 1-year Memoire [1w]

References I



Abadie, A., Diamond, A., and Hainmueller, J. (2010).

Synthetic control methods for comparative case studies: Estimating the effect of california's tobacco control program.

Journal of the American Statistical Association, 105(490).



Bolón-Canedo, V., Sánchez-Marroño, N., and Alonso-Betanzos, A. (2013).

A review of feature selection methods on synthetic data.

Knowledge and information systems, 34(3):483–519.



Bretagnolle, A. (2009).

Villes et réseaux de transport : des interactions dans la longue durée, France, Europe, États-Unis.

Hdr, Université Panthéon-Sorbonne - Paris I.

References II



Brunsdon, C., Fotheringham, S., and Charlton, M. (1998).
Geographically weighted regression.
Journal of the Royal Statistical Society: Series D (The Statistician),
47(3):431–443.



Cottineau, C., Le Néchet, F., Le Texier, M., and Reuillon, R.
(2015).
Revisiting some geography classics with spatial simulation.
*In Plurimondi. An International Forum for Research and Debate on
Human Settlements*, volume 7.



Le Néchet, F. (2015).
De la forme urbaine à la structure métropolitaine: une typologie de
la configuration interne des densités pour les principales métropoles
européennes de l'audit urbain.
Cybergeo: European Journal of Geography.

References III



Moeckel, R., Spiekermann, K., and Wegener, M. (2003).

Creating a synthetic population.

In Proceedings of the 8th International Conference on Computers in Urban Planning and Urban Management (CUPUM).



Newman, M. E. (2003).

The structure and function of complex networks.

SIAM review, 45(2):167–256.



Offner, J.-M. (1993).

Les "effets structurants" du transport: mythe politique, mystification scientifique.

Espace géographique, 22(3):233–242.

References IV



Pritchard, D. R. and Miller, E. J. (2009).

Advances in agent population synthesis and application in an integrated land use and transportation model.

In *Transportation Research Board 88th Annual Meeting*, number 09-1686.



Pumain, D. (2012).

Urban systems dynamics, urban growth and scaling laws: The question of ergodicity.

In *Complexity Theories of Cities Have Come of Age*, pages 91–103. Springer.



Raimbault, J. (2016).

Calibration of a spatialized urban growth model.

Working Paper, draft at

<https://github.com/JusteRaimbault/CityNetwork/tree/master/Docs/Papers>

References V



Reuillon, R., Leclaire, M., and Rey-Coyrehourcq, S. (2013).
Openmole, a workflow engine specifically tailored for the distributed
exploration of simulation models.
Future Generation Computer Systems, 29(8):1981–1990.



Schmitt, C. (2014).
*Modélisation de la dynamique des systèmes de peuplement: de
SimpopLocal à SimpopNet*.
PhD thesis, Paris 1.



Van den Bulcke, T., Van Leemput, K., Naudts, B., van Remortel,
P., Ma, H., Verschoren, A., De Moor, B., and Marchal, K. (2006).
Syntren: a generator of synthetic gene expression data for design
and analysis of structure learning algorithms.
BMC bioinformatics, 7(1):43.

References VI



Ye, X. (2011).

Investigation of underlying distributional assumption in nested logit model using copula-based simulation and numerical approximation.

Transportation Research Record: Journal of the Transportation Research Board, (2254):36–43.