### Multivariate

### Hao

N / - - - I - I

OVB

Mod

DGF

The Least Co...

Moscure the

Goodness of Fit

Multi-collinearit

Control Variab

CI

Hypothesis Single

CI Single Application

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1

Confidence Sets for Multiple

References

# Multivariate Regression <sup>1</sup>

Jasmine(Yu) Hao

Faculty of Business and Economics Hong Kong University

September 9, 2021

<sup>&</sup>lt;sup>1</sup>This section is based on Stock and Watson (2020), €hapter 6-7. ■ ▶

Model

OVE

DO

OLS Estimat

Assumption

Measure the Goodness of F

Goodness of F

Control Varia

CI

Hypothesis Singl

Testing Hypothese

on Two or More Coefficients

Restrictions
Involving Multiple
Coefficients

Confidence Sets for Multiple

Reference

- ▶ Although school districts with lower student-teacher ratios tend to have higher test scores in the California data set,
- other advantages that help them perform well on standardized tests.
- Could this have produced a misleading estimate of the causal effect of class size on test scores, and, if so, what can be done?

Mode

OVB

DGP

OLS Estimate

Assumption

Measure the

Goodness of F

Control Varia

Hypothesis an

Munotheric Single

er er

Application
Testing Hynothes

on Two or More Coefficients

Restrictions
Involving Multiple
Coefficients

Extension to q ¿ 1 Confidence Sets fo Multiple

Reference

# **Omitted Variable Bias**

- By focusing only on the student-teacher ratio, ignored some potentially important determinants of test scores.
  - school characteristics, such as teacher quality and computer usage,
  - student characteristics, such as family background.
- Considering an omitted student characteristic that is particularly relevant in California because of its large immigrant population: the prevalence in the school district of students who are still learning English.

Assumption

Measure the Goodness of Fi

Multi-collinear

Control V

CÍ

Hypothesis Sing

CI Single

Testing Hypothes

Coefficients
Testing Single

Restrictions
Involving Multiple
Coefficients

Confidence Sets for Multiple

Reference:

# Control for percentage of English learners

- ▷ By ignoring the percentage of English learners in the district, the OLS estimator could be biased:
  - not equal the true causal effect.
- Students who are still learning English might perform worse on standardized tests than native English speakers.
- Direction of the bias?
  - If positive correlation(districts with large classes also have many students still learning English).

The Least Sq

Measure the

Multi-collinear

Control Va

Hypothesis a

Hypothesis Sing

CI C:--I-

Application

Testing Hypothes

on Two or More Coefficients

Restrictions
Involving Multiple
Coefficients

Extension to q ¿ 1

Confidence Sets for Multiple

References

# Control for percentage of English learners

- ▶ Wrong Implication: Accordingly, based on the analysis of bivariate regression, the superintendent might hire enough new teachers to reduce the student-teacher ratio by 2, but her hoped-for improvement in test scores will fail to materialize if the true coefficient is small or zero.
- ▶ The correlation between the student-teacher ratio and the percentage of English learners (students who are not native English speakers and who have not yet mastered English) in the district is 0.19.
- ▶ If the student-teacher ratio were unrelated to the percentage of English learners, safe to ignore.

Model

OVB

IVIOC

OLS Estimat

The Least Co.

Measure the

Goodness of Fi

Mulai -- Ili----i

Control Variable

CI

Hypothesis Sing

CI Single

CI Single

Testing Hypothes on Two or More

Coefficients
Testing Single

Restrictions Involving Multiple Coefficients

Confidence Sets f

Multiple Coefficients

References

# Control for percentage of English learners

| TABLE 6.1 Differences in Test Scores for California School Districts with Low and High Student-Teacher Ratios, by the Percentage of English Learners in the District |                               |     |                               |     |  |             |
|--|-------------------------------|-----|-------------------------------|-----|--|-------------|
|  | Student-Teacher<br>Ratio < 20 |     | Student-Teacher<br>Ratio ≥ 20 |     | Difference in Test Scores,<br>Low vs. High Student-<br>Teacher Ratio |             |
|  | Average<br>Test Score         | n   | Average<br>Test Score         | n   | Difference   | t-statistic |
| All districts  | 657.4                         | 238 | 650.0                         | 182 | 7.4  | 4.04        |
| Percentage of English learners   |                               |     |                               |     |  |             |
| < 1.9%   | 664.5                         | 76  | 665.4                         | 27  | -0.9   | -0.30       |
| 1.9-8.8%   | 665.2                         | 64  | 661.8                         | 44  | 3.3  | 1.13        |
| 8.8-23.0%  | 654.9                         | 54  | 649.7                         | 50  | 5.2  | 1.72        |
| > 23.0%  | 636.7                         | 44  | 634.8                         | 61  | 1.9  | 0.68        |

**Figure** 

OLC Estima

The Least Squ

Measure the Goodness of F

Multi collingari

Control Varia

Hypothesis ar

Hypothesis Singl

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Coefficients

Extension to q / 1

Confidence Sets fo Multiple

Reference

# Control for percentage of English learners IV

What can you do about omitted variable bias?

- ▷ select a subset of districts that have the same fraction of English learners but have different class sizes:
- ▶ class size cannot be picking up the English learner effect because the fraction of English learners is held constant.

# Mathematical Expression for OVB

$$\triangleright$$
 **DGP**:  $Y_i = X_i\beta + u_i$ ,

$$\triangleright$$
  $cor(u_i, x_i) = \rho_{ux} \neq 0.$ 

$$\hat{\beta}_1 \rightarrow_{p} \beta_1 + \frac{\sigma_{u}}{\sigma_{x}} \rho_{ux}$$

- 1. OVB cause problem for the consistency of  $\hat{\beta}$ .
- 2. The magnitude of the bias depends on  $|\rho_{ux}|$ .
- 3. The direction of the bias depends on the sign of  $\rho_{ux}$ .

OLS Estimate

Measure the

Goodness of F

Control Var

Hypothosis a

Hypothesis Single

CI Single

Application
Testing Hypothe

Coefficients
Testing Single

Involving Multiple Coefficients Extension to q / 1

Confidence Sets fo Multiple

References

# Multiple Regression Model

- ▶ Multivariable OLS extends the single variable OLS, include additional variables as regressors.
- ▷ Causal inference, permits estimating the effect on  $Y_i$  of changing one variable  $X_{1i}$  while holding the other regressors ( $X_{2i}$  and  $X_{3i}$ , and so forth) constant.
- $\triangleright$  e.g. **isolate the effect** on test scores  $Y_i$  of the student–teacher ratio  $X_{1i}$  while **holding constant** the percentage of students in the district who are English learners  $X_{2i}$ .
- ▶ the multiple regression model can improve predictions by using multiple variables as predictors.

Goodness of Fi

Control Variable

Hypothesis ar

Hypothesis Sing

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1
Confidence Sets for Multiple

Reference

# The Population Regression Line

- $\triangleright$  Two independent variables,  $X_{1i}$  and  $X_{2i}$ .
- ▶ Interest in average relationship between these two independent variables and the dependent variable:

$$E[Y_i|X_{1i} = x_1, X_{2i} = x_2] = \beta_0 + \beta_1 x_1 + \beta_2 x_2,$$

where  $E[Y_i|X_{i1},X_{i2}]$  is the conditional expectation of  $Y_i$  given X.

- ▶ The equation is referred as the **population regression line**.
- ▶ The interpretation of the coefficient  $\beta_1$  is the predicted change in Y between two observations with a unit difference in  $X_1$  controlling for  $X_2$ :

$$\beta_1 = \frac{\Delta Y}{\Delta X_1}.$$

## Distribution of Frror

- $\triangleright$  The error term  $u_i$  in the multiple regression model is homoskedastic if the variance of the conditional distribution of  $u_i$  given  $X_{1i}, \ldots, X_{ki}$  is constant for  $i = 1, \ldots, n$ ,
- $\triangleright$  and thus does not depend on the values of  $X_{1i}, \ldots, X_{ki}$ . Otherwise, the error term is **heteroskedastic**.

Mode

OVE

DO

### OLS Estimator

The Least Squa

Measure the

Goodness of F

Control Var

.....

CI

CI Single

Application Testing Hypothes

on Two or More Coefficients Testing Single

Restrictions Involving Multiple Coefficients

Extension to q ¿ 1 Confidence Sets for

Multiple Coefficients

Reference

- $\triangleright$  We estimate the unknown population coefficients  $\beta_0, \beta_1, \dots, \beta_k$  using a sample of data.
- ightharpoonup The estimators of the coefficients  $\hat{eta}_0,\hat{eta}_1,\ldots,\hat{eta}_k$  that minimize the sum of squared mistakes are called the **ordinary least** squares (OLS) estimators of  $eta_0,eta_1,\ldots,eta_k$ .

# The Least Square Assumption

Parameters of interest: estimate the coefficients  $\beta_1, \ldots, \beta_k$ .

There are **four least squares assumptions** for causal inference in the multiple regression model.

The first three are those of Section 4.3 for the single-regressor model extended to allow for multiple regressors.

- 1. Assumption 1: The conditional distribution of  $u_i$  given  $X_1, \ldots, X_k$  has mean of 0.
- 2. Assumption 2:  $(X_{1i}, \ldots, X_{ki}, Y_i)$ ,  $i = 1, \ldots, n$  are i.i.d
- 3. Assumption 3: Large outliers are unlikely.
- 4. Assumption 4: No perfect multicollinearity. other regressors.

Mode

Mo

OLS Ect

The Least Squa

### Measure the Goodness of Fit

Goodness of Fit

Control Varial

Hypothesis a

Hypothesis Single

er er

Testing Hypothes

Coefficients
Testing Single

Restrictions Involving Multiple Coefficients

Confidence Sets for

Multiple Coefficients

Reference

- 1. The Standard Error of the Regression (SER).
  - ⋄ The standard error of the regression (SER) estimates the standard deviation of the error term u<sub>i</sub>.
  - $\diamond$  and where SER is the sum of squared residuals,  $SER = s_{\hat{u}} = \sqrt{s_{\hat{u}}^2}$  where  $s_{\hat{u}}^2 = \frac{1}{n-k-1} \sum_{i=1}^n \hat{u}_i^2$ .
  - $\diamond$  compare SER with the divisor is n k 1 rather than n-2.

Multi-collinearit

Control Variable

CI .....

er er

Application
Testing Hypothe

Coefficients
Testing Single

Restrictions Involving Multiple Coefficients

Extension to q ¿ 1 Confidence Sets fo Multiple

References

2. The  $R^2$ . the fraction of the sample variance of Y explained by the regressors.

$$R^2 = \frac{ESS}{TSS} = 1 - \frac{SSR}{TSS},$$

where the **explained sum of squares** is  $ESS = \sum_{i=1}^{n} (\hat{Y} - \bar{Y})^2$  and the **total sum of squares** is  $TSS = \sum_{i=1}^{n} (Y_i - \bar{Y})^2$ .

- 3. The Adjusted  $R^2$ , or  $\bar{R}^2$ .
  - $\diamond$  More variables increase  $R^2$ .
  - $\diamond$  Use adjusted  $ar{R}^2$ ,  $ar{R}^2=1-rac{n-1}{n-k-1}rac{SSR}{TSS}=1-rac{s_{\hat{\mu}}^2}{s_{Y}^2}.$

OVI

DGF

The Least Say

Assumption

Goodness of

Multi-collinearity

CI

Hypothesis Sing

CI Single

Testing Hypothes

Coefficients
Testing Single

Involving Multipl Coefficients

Confidence Sets for

Confidence Sets for Multiple Coefficients

Reference

Easy to attempt to calculate a regression with linearly dependent regressors.

- 1. Including the same regressor twice.
- 2. Including regressors which are a linear combination of one another, such as education, experience

Mode

...

DGF

The Least Co.

Assumption

Goodness

Multi-collinearity

WillierColline

Hypothesis ar

Hypothesis Single

CI Single

Testing Hypothese

Testing Single
Restrictions

Coefficients
Extension to a / 1

Confidence Sets for Multiple

Reference:

# Some Examples:

- 1. Including a dummy variable and its square.
- 2. Estimating a regression on a sub-sample for which a dummy variable is either all zeros or all ones.
- 3. Including a dummy variable interaction which yields all zeros.
- 4. Including more regressors than observations.

Goodness of Fi

Control Variable

### Hypothesis an

CI

CLC: 1

Application

Testing Hypotheses on Two or More Coefficients

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1 Confidence Sets for Multiple

Deference

Consider the multiple regression model with control variables

$$Y_i = X_i^{\top} \beta_{(1)} + W_i^{\top} \beta_{(2)},$$

- $\triangleright$  k variables of interest, denoted by X,
- $\triangleright$  and r control variables, denoted by W where  $\beta_{(1)}$  is the causal effect,  $W_i$  is the control variable,

Control Variable

# Assumptions:

- 1. Assumption 1:(conditional independence)  $E(u_i|X_i, W_i) = E(u_i|W_i) = 0.$
- 2. Assumption 2:  $(X_i, W_i, Y_i)$  are i.i.d.
- 3. Assumption 3:  $X_i$  and  $W_i$  have nonzero finite fourth moment.
- 4. Assumption 4: No perfect collinearity.

OV

- -

OLS Estimate

Assumption
Measure the

Goodness of F

### Control Variable

CI

Hypothesis Sing

CI Single

Testing Hypothese

Testing Single Restrictions Involving Multiple

Coefficients
Extension to q ¿ 1

Confidence Sets for Multiple

References

# Conditional mean independence

**Conditional mean independence** requires that the conditional expectation of  $u_i$  given the variable of interest and the control variables does not depend on the variable of interest, although it can depend on control variables.

- $\triangleright$  once you control for the W's the X's can be treated as if they were randomly assigned.
- X's uncorrelated with the error term.
- $\triangleright$  OLS can estimate the causal effects on Y of a change in each of the X's.

Measure the

Goodness of F

Control Variable

### Hypothesis an

CI Hypothesis Single

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1 Confidence Sets for Multiple

References

# Example of Control Variable I

$$\widehat{TestScore} = 686.0 - 1.10 \times STR - 0.65 \times PctEL, \qquad (6.12)$$

$$\widehat{TestScore} = 700.2 - 1.00 \times STR - 0.122 \times PctEL - 0.547 \times LchPct. \quad (6.16)$$

- Consider the potential OVB arising from outside learning opportunities.
  - difficult to measure,
  - correlated with the students' economic background, (can be measured).

OVB

IVIO

OLS Estimate

Assumption

Goodness of Fit

Multi-collinearity

Control Variable

Hypothesis a

Li...aharia Ciral

or or a

Application
Testing Hypothes

Testing Single Restrictions

Extension to q ¿ 1 Confidence Sets fo

Reference

# Example of Control Variable II

- ▶ regression of test scores on STR and PctEL
- ▶ with the percentage of students receiving a free or subsidized school lunch(*LchPct*).
- Students are eligible if their family income less than 150% of the poverty line
- b the coefficient on the STR is the effect of the student−teacher ratio on test scores,
  - $\diamond$  Including the control variable *LchPct* does not substantially change any conclusions about the class size effect: The coefficient on STR changes only slightly from its value of -1.10 in Equation (6.12) to -1.00 in Equation (6.16)

OV

IVIC

OLS Estimate

Assumption

Goodness of Fit

Multi-collinearity

Hypothesis a

CI

Hypothesis Sing

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1

Multiple Coefficients

Reference

## Choose Control Variables I

Multiple regression makes it possible to control for factors that could lead to omitted variable bias in the estimate of the effect of interest.

- ▶ how does one determine the "right" set of control variables?
- ▶ the conditional mean independence condition.
- $\triangleright$  to eliminate omitted variables bias, a set of control variables must satisfy  $E[u_i|X_i,W_i]=E[u_i|W_i]$ ,
  - where X<sub>i</sub> denotes the variable or variables of interest and W<sub>i</sub> denotes one or more control variables.

### . . . .

OVB Mode

OLS Estimator

The Least Squar

Measure the Goodness of Fit

Goodness of Fi

Control Variable

### CI

Hypothesis Singl

Application

on Two or More Coefficients

Restrictions
Involving Multiple
Coefficients

Extension to q ¿ 1 Confidence Sets for Multiple

Reference

# Choose Control Variables II

- ▶ Use judgement to determining which control variables to include.
  - ⋄ For example, economic conditions vary across school districts with the same percentage of English learners.
  - more affluent districts would be expected to have lower class sizes,
  - Affluent families tend to have more access to outside learning opportunities.
  - $\diamond~$  OVB even after controlling for the percentage of English learners.
- ▶ Logic leads to including one or more additional control variables in the test score regressions, where the additional control variables measure economic conditions of the district.
- Our approach to the challenge of choosing control variables is twofold.
  - First, a core or base set of regressors should be chosen using a combination of expert judgment, economic theory, and knowledge of how the data were collected; the regression using this base set of regressors is sometimes referred to as a base specification.

OV

IVIOC

OLS Estimato

Assumption

Measure the

Goodness of Fi

Control Variable

cí

Hypothesis Sing

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Coefficients

Confidence Sets fo Multiple

Reference

## Choose Control Variables III

- This base specification should contain the variables of primary interest and the control variables suggested by expert judgment and economic theory.
- develop a list of candidate alternative specifications—that is, alternative sets of regressors.
- ▶ If the estimates of the coefficients of interest change substantially across specifications—OVB.

Goodness of Fi

Control Var

Hypothesis and

Hypothesis Single

### riypotiicaia biiig

Application
Testing Hypothes

Coefficients
Testing Single

Restrictions
Involving Multiple
Coefficients

Extension to q ¿ 1
Confidence Sets fo

Reference

# Hypothesis test for single coefficient I

- ▶ Want to test the hypothesis that a change in *STR* has no effect on test scores, holding constant *PctEl*.
- $\vdash H_0: \beta_1 = 0.$
- $\triangleright$  Moregenerally: can test  $H_0: \beta_j = \beta_{j,0}$ .
- ▶ If the alternative hypothesis is two-sided, then the two hypotheses can be written mathematically as

$$H_0: \beta_j = \beta_{j,0} \ v.s. \ H_1: \beta_j \neq \beta_{j,0}.$$

### Mode

OV

D

OLS Estimat

Assumption

Goodness of Fit

ividiti-collinea

Hypothesis and CI

### Hypothesis Single

### CI Single

Testing Hypothes

Testing Single Restrictions Involving Multiple

Coefficients

Extension to a + 1

Confidence Sets for Multiple

Reference

# Hypothesis test for single coefficient II

- $\triangleright$  For example, if the first regressor is STR, then the null hypothesis that changing the student–teacher ratio has no effect on test scores corresponds to the null hypothesis that  $\beta_1=0$  (so  $\beta_{1,0}=0$ ).
- $\triangleright$  Task: test the null hypothesis  $H_0$  against the alternative  $H_1$  using a sample of data.
- ▶ Steps:
  - 1. compute the standard error
  - 2. compute the t-statistics
  - 3. compute the p-value

Measure the

Goodness of Fi

Multi-collinear

CI

Hypothesis Singl

CI Single

Testing Hypothe on Two or More

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1

Confidence Sets for Multiple

References

# Confidence Intervals for a Single Coefficient I

- ▶ Same as in the single-regressor model.
- ▶ Note: only work in large samples.
- ▶ When the sample size is large, the 95% confidence interval is

$$CI_{0.95}(\beta_j) = [\hat{\beta}_j - 1.96se(\hat{\beta}_j), \hat{\beta}_j + 1.96se(\hat{\beta}_j)].$$

Measure the

Goodness of F

Control Va

ar ar a

Hypothesis Sing

CI Single

Application

on Two or More Coefficients

Restrictions
Involving Multiple
Coefficients

Extension to q ¿ 1 Confidence Sets for

References

- be test the null hypothesis that a change in the student-teacher ratio has no effect on test scores, control for the percentage of English learners in the district?
- ▶ The regression of test scores against STR and PctEL,

$$\widehat{TestScore} = 686.0 - 1.10 * STR - 0.650 * PctEL(7.5)$$

- ▷ Construct the t-statistic t = (-1.10 0)/0.43 = -2.54.
- ▶ Associated p-value is  $2\Phi(-2.54) = 0.011$ .
- $\triangleright$  Reject at  $H_0$ .

### Application

# Application II

- ▶ A 95% confidence interval for the population coefficient on STR is  $-1.10 \pm 1.96 * 0.43 = (-1.95, -0.26)$ .
- ▶ If the superintendent's interest in decreasing the student-teacher ratio by 2, the 95% confidence interval for the effect on test scores of this reduction is (-0.26\*-2,-1.95\*-2) = (0.52,3.902).

. . . .

OVE

Mod

OLS Estimate

The Least Squa Assumption

Measure the Goodness of Fi

C------ V--:--

Hypothocic

CI

rrypotnesis sing

CI Single Application

Testing Hypothe

Testing Single Restrictions

Extension to q ¿ 1
Confidence Sets for

References

# Example with expenditures per student I

Adding expenditures per pupil to the equation.

▶ What is the effect on test scores of reducing the student-teacher ratio, holding expenditures per student constst?

 $\triangleright$ 

$$\widehat{\textit{TestScore}} = \underset{(15.5)}{649.6} - \underset{(0.48)}{0.29} * STR + \underset{(1.59)}{3.87} * \textit{Expn} - \underset{(0.032)}{0.656} * \textit{PctEL}, (7.6)$$

- coefficient on STR is -1.10, but after adding Expn as a regressor in Equation (7 .6), it is only -0.29.
- t = (-0.29 0)/0.48 = -0.60,
- $\diamond$  Cannot reject  $H_0$  at the 10% significance level.
- Interpretation of (7.6) is that: in these California data, school administrators allocate their budgets efficiently.
- Counterfactual, the coefficient on STR in (7.6) were negative and large.
  - school districts could raise their test scores simply by decreasing funding for other purposes (textbooks, technology, sports, and so on) and using those funds to hire more teachers.
  - this transfer would have little effect on test scores.

### Multivariate

### Hao

OVE

...

OLS Estima

The Least Squa

Measure the

Goodness of Fi

Control Variab

CÍ

CI Single

Application

Testing Hypothesion Two or More

Testing Single Restrictions Involving Multiple

Coefficients

nfidence Sets for ultiple

References

# Example with expenditures per student II

 Remark: the standard error on STR increased when Expn was added, from 0.43 to 0.48. Imperfect multicollinearity, that correlation between regressors (the correlation between STR and Expn is -0.62) can make the OLS estimators less precise.

OVE

DO

The Least Sq

Measure the

Goodness of Fi Multi-collineari

CI

Hypothesis Sing

Application

### Testing Hypotheses on Two or More Coefficients

Testing Single Restrictions Involving Multiple Coefficients

Extension to q ¿ 1
Confidence Sets fo

References

# Testing Hypotheses on Two or More Coefficients I

- Consider the model
- Consider the test

$$H0: \beta_1 = 0$$
 and  $\beta_2 = 0$  vs  $H1: \beta_1 \neq 0$  and  $/$  or  $\beta_2 \neq 0$ .

- ▶ Example of joint hypothesis on the coefficients in the multiple regression model.
- ▶ The null hypotheses is  $H_0$ :  $\beta_1 = 0$  and  $\beta_2 = 0$ .

A joint hypothesis is a hypothesis that imposes two or more restrictions on the regression coefficients.

Consider the alternative form:  $H_0: \beta_j = \beta_{j,0}, \dots \beta_m = \beta_{m,0}$ , for a total of q restrictions, vs.  $H_1:$  one or more of the q restrictions under H0 does not hold.

Testing Hypotheses on Two or More Coefficients

# Why can't I just test the individual coefficients one at a time?

- Seemingly possible to test a joint hypothesis by using t-statistics to test the restrictions one at a time, the following calculation shows that this approach is unreliable.
- Specifically, suppose you are interested in testing the joint null hypothesis that  $\beta_1 = 0$  and  $\beta_2 = 0$ .
- $\triangleright$  Let  $t_1$  and  $t_2$  be the t-statistic for testing  $\beta_1=0$  and  $\beta_2=0$
- ▶ What happens when you use the "one-at-a-time" testing procedure: Reject the joint null hypothesis if either  $t_1$  or  $t_2$ exceeds 1.96 in absolute value?

We derive the joint distribution of  $t_1$  and  $t_2$ .

The Least So

Measure the

Goodness of Fi

CI Hypothesis Single

CI Single

Testing Hypotheses on Two or More Coefficients

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1 Confidence Sets fo

Multiple Coefficients

Reference

- ▶ The homoskedasticity-only F-statistic is computed using a simple formula based on the sum of squared residuals from two regressions.
- ▶ **Restricted regression**: the null hypothesis is forced to be true.
- ▶ **Unrestricted regression**: the alternative hypothesis is allowed to be true.
- ▶ The homoskedasticity-only F-statistic is given by the formula

$$F = \frac{(SSR_{restricted} - SSR_{unrestricted})/q}{SSR_{unrestricted}/(n - k_{unrestricted} - 1)},$$

where *SSR* is the sum of squared residuals.

Goodness of F

Control Varial

CI

Hypothesis Sing

Testing Hypotheses on Two or More

Coefficients
Testing Single
Restrictions

Extension to q ¿ 1 Confidence Sets for Multiple

Reference

We can write the F-test as

$$F = \frac{(R_{unrestricted}^2 - R_{restricted}^2)/q}{(1 - R_{unrestricted}^2)/(n - k_{unrestricted} - 1)}.$$

- ▶ If the errors are homoskedastic, then the difference between the homoskedasticityonly F-statistic and the heteroskedasticityrobust F-statistic vanishes as the sample size n increases.
- ightharpoonup If the errors are homoskedastic, the sampling distribution of the homoskedasticity-only F-statistic under the null hypothesis is, in large samples,  $F_{q,\infty}$ .
- Advantages: easy to compute, clear to interpret. Disadvantage: only apply to homoskedastic errors.
- ▶ In practice the homoskedasticity-only F-statistic is not a satisfactory substitute for the heteroskedasticityrobust F-statistic.

OVE

IVIC

OLS Estimate

Assumption
Measure the

Goodness of Fi

Control Varia

Hypothesis ar

Hypothesis Singl

CI Single

Testing Hypotheses on Two or More Coefficients

Testing Single Restrictions Involving Multiple

Extension to q ¿ 1 Confidence Sets fo Multiple

References

## Using the homoskedasticity-only F-statistic when n is small

- ightharpoonup If the errors are i.i.d., homoskedastic, and normally distributed, then the homoskedasticity-only F-statistic defined has an  $F_{q,n-k_{unrestricted}-1}$  distribution under the null hypothesis.
- ▷ Critical values depend on both q and  $k_{unrestricted}$ , check SW Appendix Table 5 or use qf(p, df1, df2) in R.
- $\triangleright F_{q,n-k_{unrestricted}-1} \rightarrow_d F_{q,\infty}$  as n increases;
- ▶ For small samples, however, the two sets of critical values differ. However, it is easy to read more into them than they deserve.

Measure the Goodness of F

Multi-collineari

Control vari

CI

Hypothesis Sing

Testing Hypotheses on Two or More

On I wo or More Coefficients

Restrictions
Involving Multiple
Coefficients

Confidence Sets f

Confidence Sets fo Multiple Coefficients

References

# Application to test scores and the student-teacher ratio I

- $\triangleright$   $H_0$ : coefficients on STR and and Expn are 0, controlling for PctEL.
- ▶ The unrestricted regression has the regressors STR, Expn, and PctEL:

$$\widehat{TestScore} = 649.6 - 0.29 * STR + 3.87 * Expn - 0.656 * PctEL,$$
  
 $(0.032)$   $R^2 = 0.4366$ 

▶ The restricted model:

$$\widehat{TestScore} = 664.7 - 0.671 * PctEL,$$

$$(1.0) \qquad (0.032)$$

$$R^2 = 0.4149$$

### OV

Mo

OLS Estimate

Assumption
Massure the

Goodness of F

Control Va

Hypothesis a

Ulunothoric Single

CLC: I

Testing Hypotheses

### on Two or More Coefficients

Testing Single Restrictions Involving Multiple Coefficients

Confidence Sets for

Reference

# Application to test scores and the student-teacher ratio II

- $\triangleright$  The number of restrictions is q = 2,
- $\triangleright$  The number of observations is n = 420,
- $\triangleright$  The number of regressors in the unrestricted regression is k=3.
- ▶ The homoskedasticity-only F-statistic,

$$F = \frac{(0.4366 - 0.4149)/2}{(1 - 0.4366)/(420 - 3 - 1)} = 8.01.$$

- $\triangleright$  For  $F_{2,\infty}$ , 1% critical value of 4.61
- $\triangleright$  Rejected at the 1% level using the homoskedasticity-only test.
- ▶ Remark: The heteroskedasticity-robust F-statistic testing this joint hypothesis is 5.43, quite different from the less reliable homoskedasticityonly value of 8.01.

Goodness of F

Control Va

Hypothesis ar

Hypothesis Singl

CIC: I

Testing Hypothes

on Two or More Coefficients Testing Single

Restrictions
Involving Multiple
Coefficients

Confidence Sets 1

Multiple Coefficients

References

# Testing Single Restrictions Involving Multiple Coefficients I

Economic theory may suggest a single restriction that involves two or more regression coefficients. For example, theory might suggest a null hypothesis of the form  $\beta_1 = \beta_2$ ;

D

$$H_0: \beta_1 = \beta_2, \text{ v.s. } H_1: \beta_1 \neq \beta_2.$$

 $\triangleright$  single restriction, q=1, involves multiple coefficients ( $\beta_1$  and  $\beta_2$ ).

We need to modify the methods presented so far to test this hypothesis. There are two approaches; which is easier depends on your software

OVE

Mo

OLS Estimate

The Least Squa

Measure the Goodness of Fi

Mula: --II:----

Control Variab

Hypothesis an

Hypothesis Singl

Cl Single

Testing Hypothes

on Two or More Coefficients

Testing Single Restrictions Involving Multiple Coefficients

Confidence Sets for Multiple

D-f----

# Testing Single Restrictions Involving Multiple Coefficients II

- 1. Approach 1: Test the restriction directly.
  - $\diamond$  Test with an F-statistic that, because q=1, has an  $F_1$ -distribution
  - $\diamond$  Under the null hypothesis, the 95% percentile of the  $F_1$ -distribution is  $1.96^2 = 3.84$ .

. . .

OVE

...

OLS Estimat

Assumption

Goodness of Fi

Multi-collineari

Control Variable

Hypothesis an

Hypothesis Singl

01.01

CI Single

Testing Hypothesi on Two or More

Testing Single Restrictions Involving Multiple Coefficients

Extension to q ¿ 1
Confidence Sets fo

Multiple

References

# Testing Single Restrictions Involving Multiple Coefficients III

- 2. Approach 2: Transform the regression.
  - Rewrite the equation and turn the restriction on on a single regression coefficient.

OLS Estimat

Moscure the

Goodness of F

Control Vari

and the second

CI

CI Single

Application Testing Hypothese

Coefficients
Testing Single

Coefficients

Extension to q / 1

Confidence Sets fo

Deference

## Extension to q > 1 l

- ▶ In general, *q* restrictions under the null hypothesis in which some or all of these restrictions involve multiple coefficients.
- ▶ User the *F*-statistic to test this type of joint hypothesis.
- ightharpoonup The *F*-statistic can be computed by either of the two methods just discussed for q=1.

0)//

Mo

OLS Estima

The Least Squ. Assumption

Measure the Goodness of Fi

Goodness of Fi

Control V

CI

Hypothesis Sing

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Coefficients

Confidence Sets for Multiple Coefficients

Reference

## Confidence Sets for Multiple Coefficients I

- ▶ The method is conceptually similar to that of a single coefficient using the *t*-statistic except that the confidence set for multiple coefficients is based on the *F*-statistic.
- A 95% confidence set for two or more coefficients is a set that contains the true population values of these coefficients in 95% of randomly drawn samples.
- ▶ Recall that a 95% confidence interval is computed by finding the set of values of the coefficients that are not rejected using a t-statistic at the 5% significance level.

Goodness of F

C--+--| \/--i-h

Hypothesis ar

Hypothesis Singl

CI Single

Testing Hypothes

Testing Single Restrictions


Coefficients
Extension to q ¿

Confidence Sets for Multiple Coefficients

Reference

# Confidence Sets for Multiple Coefficients

This approach can be extended to the case of multiple coefficients.



 $\triangleright$  This ellipse does not include the point (0,0).

Measure the

Goodness of Fi

Control Varia

. . . . . . .

CI

CLC: 1

Application

on Two or More Coefficients

Testing Single Restrictions Involving Multiple Coefficients

Extension to q ¿

Confidence Sets for Multiple Coefficients

Reference:

# Confidence Sets for Multiple Coefficients

- This means that the null hypothesis that these two coefficients are both 0 is rejected using the F-statistic at the 5% significance level.
- ▷ The confidence ellipse is oriented in the lower-left/upper-right direction.
- ▶ Estimated correlation.

OLS Estimat

Assumption

Measure the

Mark: --II:----

Control V

CI

Hypothesis Singl

CI Single

Testing Hypothese on Two or More

Testing Single
Restrictions
Involving Multiple
Coefficients

Confidence Sets for Multiple

Coefficients References

## Some Remarks I

- 1. An increase in the  $R^2$  or  $R^2$  does not necessarily mean that an added variable is statistically significant.
- 2. A high  $R^2$  or  $R^2$  does not mean that the regressors are a true cause of the dependent variable.
- 3. A high  $R^2$  or  $R^2$  does not mean that there is no omitted variable bias.
- 4. A high  $R^2$  or  $R^2$  does not necessarily mean that you have the most appropriate set of regressors, nor does a low  $R^2$  or  $R^2$  necessarily mean that you have an inappropriate set of regressors.

Goodness of Fi

Control Variable

Hypothesis ar

Hypothesis Single

CI Single

Testing Hypothese on Two or More

Testing Single Restrictions Involving Multiple

Extension to q ¿ .

**Summary Statistics** 

SER

 $\overline{R}^2$ 

Confidence Sets for Multiple Coefficients

Reference

## Tabulate Results I

TABLE 7.1 Results of Regressions of Test Scores on the Student-Teacher Ratio and Student
Characteristic Control Variables Using California Elementary School Districts

Dependent variable, average test score in the district

| Regressor                                     | (1)            | (2)            | (3)            | (4)            | (5)            |
|---|----------------|----------------|----------------|----------------|----------------|
| Student-teacher ratio (X <sub>1</sub> )       | -2.28          | -1.10          | -1.00          | -1.31          | -1.01          |
|   | (0.52)         | (0.43)         | (0.27)         | (0.34)         | (0.27)         |
|   | [-3.30, -1.26] | [-1.95, -0.25] | [-1.53, -0.47] | [-1.97, -0.64] | [-1.54, -0.49] |
| Control variables                             |                |                |                |                |                |
| Percentage English learners (X <sub>2</sub> ) |                | -0.650         | -0.122         | -0.488         | -0.130         |
|   |                | (0.031)        | (0.033)        | (0.030)        | (0.036)        |
| Percentage eligible for subsidized            |                |                | -0.547         |                | -0.529         |
| lunch (X <sub>3</sub> )                       |                |                | (0.024)        |                | (0.038)        |
| Percentage qualifying for income              |                |                |                | -0.790         | 0.048          |
| assistance $(X_4)$                            |                |                |                | (0.068)        | (0.059)        |
| Intercept                                     | 698.9          | 686.0          | 700.2          | 698.0          | 700.4          |
|   | (10.4)         | (8.7)          | (5.6)          | (6.9)          | (5.5)          |

| n  | 420                 | 420                  | 420 | 420 | 420 |
|--|---------------------|----------------------|-----|-----|-----|
| These regressions were estimated using the da<br>robust standard errors are given in parenthes<br>confidence interval is given in brackets below | es under coefficien | ts. For the variable |     |     |     |

14.46

420

0.424

9.08

0.773

420

18.58

0.049

420

11.65

420

0.626

9.08

0.773

420

14100

Mo

OLS Estimato

Assumption

Measure the Goodness of Fi

Multi-collinear

CONLION VAIN

CI

rrypotnesis singi

Testing Hypothes

Coefficients
Testing Single

Involving Multiple Coefficients

Confidence Sets for Multiple

Coefficients

## Tabulate Results II

- Way compare several possible sets of regressors, tabulate the results.
- ▶ Each column presents a separate regression.
  - same dependent variable: test score.
  - statistics
  - standard error in brackets is a 95% confidence interval for the population coefficient
- ▶ The final three rows contain summary statistics:
  - SER: the standard error of the regression, SER
  - $\diamond R^2$
  - the sample size (same for all of the regressions, 420 observations).
- ▶ Use the results to test single regressor.
- ▶ Although the table does not report t-statistics, they can be computed from the information provided.

Confidence Sets for

Multiple Coefficients

## Tabulate Results III

## These results suggest three conclusions:

- 1. Controlling for these student characteristics cuts the estimated effect of the student-teacher ratio on test scores approximately in half.
  - This estimated effect is not very sensitive to which specific control variables are included in the regression.
  - In all cases, the hypothesis that the coefficient on the student teacher ratio is 0 can be rejected at the 5% level. In the four specifications with control variables, regressions (2) through (5), reducing the student-teacher ratio by one student per teacher is estimated to increase average test scores by approximately 1 point, holding constant student characteristics.

OVE

Mod

OLS Estimator

The Least Squa Assumption

Goodness of Fi

Control Va

Hypothesis

Librarah asia Cian

CLC: I

Testing Hypothes

on Two or More Coefficients

Restrictions Involving Multiple Coefficients

Confidence Sets for Multiple

Coefficients References

## Tabulate Results IV

- 2. The student characteristic variables are potent predictors of test scores. The student–teacher ratio alone explains only a small fraction of the variation in test scores: The  $R^2$  in column (1) is 0.049. The  $R^2$  jumps when the student characteristic variables are added. For example, the R2 in the base specification, regression (3), is 0.773. Districts with many English learners and districts with many poor children have lower test scores.
- 3. In contrast to the other two control variables, the percentage qualifying for income assistance appears to be redundant. As reported in regression (5), adding it to regression (3) has a negligible effect on the estimated coefficient on the student-teacher ratio or its standard error.

Model

OVE

DC

OLS Estimat

The Least Squa

Measure the

Goodness of Fi

maici comican

Control Variable

Hypothesis Sing

CI Single

Application
Testing Hypothe

on Two or More Coefficients

Testing Single Restrictions Involving Multiple Coefficients

Extension to q ¿ 1 Confidence Sets for

Multiple

References

Stock, J. H. and Watson, M. W. (2020). *Introduction to econometrics*, volume 4. Pearson New York.