

# Deep Learning

## 2.1 Quick visit to ML concepts

Dr. Konda Reddy Mopuri  
kmopuri@iittp.ac.in  
Dept. of CSE, IIT Tirupati  
Aug-Dec 2021

# Machine Learning

- ① General objective is to capture the data regularity for making predictions

# Machine Learning

- ① General objective is to capture the data regularity for making predictions
- ② In our linear regression example: we modeled the  $x$  and  $y$  as linearly related for fitting a line and predict  $y$  from  $x$

# Machine Learning

- ① General objective is to capture the data regularity for making predictions
- ② In our linear regression example: we modeled the  $x$  and  $y$  as linearly related for fitting a line and predict  $y$  from  $x$
- ③ Broadly these are types of the inferences
  - Regression (e.g. customer satisfaction, stock prediction, etc.)
  - Classification (e.g. object recognition, speech processing, disease detection etc.)
  - Density estimation (e.g. sampling/synthesize, outlier detection, etc.)

# Standard formalization

- ① Classification and Regression: considers a measure of joint probability density  $f_{X,Y}$  over the observation/value of interest and training (i.i.d.) samples  $(x_n, y_n), n = 1, \dots, N$

# Standard formalization

- ① Classification and Regression: considers a measure of joint probability density  $f_{X,Y}$  over the observation/value of interest and training (i.i.d.) samples  $(x_n, y_n), n = 1, \dots, N$
- ② Density estimation: distribution  $f_X$  and  $x_n, n = 1, \dots, N$

# Intuitive interpretation

① Classification:  $f_{X,Y}(x,y) = f_{X/Y=y}(x)P(Y=y)$

# Intuitive interpretation

- ① Classification:  $f_{X,Y}(x,y) = f_{X/Y=y}(x)P(Y=y)$
- ② Draw Y first, then given the value of Y generate X



# Intuitive interpretation

- ① Classification:  $f_{X,Y}(x,y) = f_{X/Y=y}(x)P(Y = y)$
- ② Draw  $Y$  first, then given the value of  $Y$  generate  $X$
- ③ Conditional distribution  $f_{X/Y}$  stands for the distribution of observable signal for category  $y$  (e.g. image of a dog, weight of a 30 year Indian male)

# Intuitive interpretation

① Regression:  $f_{X,Y}(x,y) = f_{Y/X=x}(y)f_X(x)$


# Intuitive interpretation

- ① Regression:  $f_{X,Y}(x, y) = f_{Y/X=x}(y)f_X(x)$
- ② First generate X, given its value generate Y

# Summary: three types of inferences

## ① Classification


- $X, Y$  random variables on  $\mathcal{L} = \mathcal{R}^D \times \{1, \dots, C\}$
- Aim is to estimate the  $\operatorname{argmax}_y P(Y = y/X = x)$



# Summary: three types of inferences


## ① Classification

- $X, Y$  random variables on  $\mathcal{L} = \mathcal{R}^D \times \{1, \dots, C\}$
- Aim is to estimate the  $\operatorname{argmax}_y P(Y = y/X = x)$



## ① Regression


- $X, Y$  random variables on  $\mathcal{L} = \mathcal{R}^D \times \mathcal{R}$
- Aim is to estimate  $\mathbb{E}(Y/X = x)$



# Summary: three types of inferences


## ① Classification

- $X, Y$  random variables on  $\mathcal{L} = \mathcal{R}^D \times \{1, \dots, C\}$
- Aim is to estimate the  $\operatorname{argmax}_y P(Y = y/X = x)$




## ① Regression

- $X, Y$  random variables on  $\mathcal{L} = \mathcal{R}^D \times \mathcal{R}$
- Aim is to estimate  $\mathbb{E}(Y/X = x)$



## ① Density estimation

- $X$  is random variable  $\mathcal{R}^D$
- Aim is to estimate the  $f_X$



# This categorization is not hard

Boundaries are vague

- ① We may perform classification via class score regression

# This categorization is not hard

Boundaries are vague

- ① We may perform classification via class score regression
- ② Density estimation can perform classification using Baye's rule



## Risk/Loss

# Loss/Risk

- ① Learning: finding a good function  $f^*$  from a set of functions  $\mathcal{F}$

# Loss/Risk

- ① Learning: finding a good function  $f^*$  from a set of functions  $\mathcal{F}$
- ② How to find the goodness of a function  $f$ ?

# Loss/Risk

- ① Learning: finding a good function  $f^*$  from a set of functions  $\mathcal{F}$
- ② How to find the goodness of a function  $f$ ?
- ③ Through a loss  $l : \mathcal{F} \times \mathcal{L} \rightarrow \mathcal{R}$

# Loss/Risk

- ① Learning: finding a good function  $f^*$  from a set of functions  $\mathcal{F}$
- ② How to find the goodness of a function  $f$ ?
- ③ Through a loss  $l : \mathcal{F} \times \mathcal{L} \rightarrow \mathcal{R}$
- ④ Such that value of  $l(f, z)$  increases with the *wrongness* of  $f$  on  $z$ :  
(measure of discrepancy between the expected and predicted)

# Loss/Risk

- ① Learning: finding a good function  $f^*$  from a set of functions  $\mathcal{F}$
- ② How to find the goodness of a function  $f$ ?
- ③ Through a loss  $l : \mathcal{F} \times \mathcal{L} \rightarrow \mathcal{R}$
- ④ Such that value of  $l(f, z)$  increases with the *wrongness* of  $f$  on  $z$ :  
(measure of discrepancy between the expected and predicted)
- ⑤
  - Regression:  $l(f, (x, y)) = (f(x) - y)^2$
  - Classification:  $l(f, (x, y)) = \mathbf{1}(f(x) \neq y)$
  - Density estimation:  $l(q, z) = -\log(q(z))$

# Loss/Risk

- ① Learning: finding a good function  $f^*$  from a set of functions  $\mathcal{F}$
- ② How to find the goodness of a function  $f$ ?
- ③ Through a loss  $l : \mathcal{F} \times \mathcal{L} \rightarrow \mathcal{R}$
- ④ Such that value of  $l(f, z)$  increases with the *wrongness* of  $f$  on  $z$ :  
(measure of discrepancy between the expected and predicted)
- ⑤
  - Regression:  $l(f, (x, y)) = (f(x) - y)^2$
  - Classification:  $l(f, (x, y)) = \mathbf{1}(f(x) \neq y)$
  - Density estimation:  $l(q, z) = -\log(q(z))$
- ⑥ Loss may have additional terms (from prior knowledge)

# Expected Risk

- ① We want  $f$  with small *expected (average) risk*  $R(f) = \mathbb{E}_z(l(f, z))$



# Expected Risk

- ① We want  $f$  with small *expected (average) risk*  $R(f) = \mathbb{E}_z(l(f, z))$
- ②  $f^* = \operatorname{argmin}_{f \in \mathcal{F}} R(f)$

# Expected Risk

- ① We want  $f$  with small *expected (average) risk*  $R(f) = \mathbb{E}_z(l(f, z))$
- ②  $f^* = \operatorname{argmin}_{f \in \mathcal{F}} R(f)$
- ③ This is unknown. However, if the training data  $\mathcal{D} = \{z_1, \dots, z_N\}$  is i.i.d. we can estimate the risk empirically (known as empirical risk),

$$\hat{R}(f; \mathcal{D}) = \hat{\mathbb{E}}_{\mathcal{D}}(l(f, z)) = \frac{1}{N} \sum_{i=1}^N l(f, z_n)$$