

**AE/ME 330 Spring 2013 Homework I, Due: Friday February 8th**

1. The non-dimensional displacement of an airplane wing tip ( $f(t)$ ) obtained from an aeroelastic flutter study is represented by

$$f(t) = 2\sin(t^2) - 3t\sin(t^2) + t^2\sin(t^2).$$

where  $t$  is the non-dimensional time.

- (a) For three different starting points  $t_0 = 2.2$ ,  $t_0 = 2.3$ , and  $t_0 = 2.4$ , find the roots of  $f(t)$  (non-dimensional times at which the displacement is equal to zero) using Newton's Method. For each case, tabulate  $t_n$ ,  $f(t_n)$ ,  $\Delta_n t = t_{n+1} - t_n$ , and  $t_{n+1}$  at each iteration level  $n$ . Use a convergence criteria of  $(\epsilon_n)_r < 10^{-8}$  where  $(\epsilon_n)_r = \frac{|t_{n+1} - t_n|}{|t_n|}$ .
  - (b) Use the method of linear interpolation (Regula-Falsi) to find the root of  $f(t) = 0$  for  $3.2 \leq t \leq 3.8$  with the same convergence criteria given in part (a). Tabulate  $t_{n+1}$ ,  $f(t_{n+1})$ , and  $(\epsilon_n)_r$  at each iteration level  $n$ .
2. The free vibrations of a cantilever beam satisfy the equation

$$\cosh(\alpha) \times \cos(\alpha) = -1$$

where the dimensionless parameter  $\alpha$  is proportional to the square root of the natural frequency. Using Secant Method, find the three smallest positive values of  $\alpha$  satisfying this equation (pick the appropriate initial guesses for finding each root). For each root: (1) use a convergence criteria of  $(\epsilon_n)_r = \frac{|\alpha_{n+1} - \alpha_n|}{|\alpha_n|} < 10^{-6}$ , (2) tabulate the values of  $\alpha_n$ ,  $f(\alpha_n)$ ,  $\Delta_n \alpha = \alpha_{n+1} - \alpha_n$ , and  $\alpha_{n+1}$  at each iteration.

3. For  $-1.0 \leq x \leq 0.0$ , find the root of the function

$$f(x) = e^{6x} + 3(\ln 2)^2 e^{2x} - e^{4x} \ln 8 - (\ln 2)^3 = 0.$$

- (a) Use Newton's Method with a starting point of  $x_0 = -1.0$ .
- (b) Use Modified Newton's Method with a starting point of  $x_0 = -1.0$ .

For each method: (1) approximate the root with a convergence criteria of  $(\epsilon_n)_r = \frac{|x_{n+1}-x_n|}{|x_n|} < 10^{-5}$ , (2) tabulate the values of  $x_n$ ,  $f(x_n)$ ,  $\Delta_n x = x_{n+1} - x_n$ , and  $x_{n+1}$  at each iteration, and (3) plot  $|f(x_n)|/|f(x_0)|$  vs. iteration number on a semi-log plot. Comment on the convergence of each method.

4. (Modified from problem 8.35 in the textbook) Mechanical and Aerospace engineers use thermodynamics extensively in their work. The following polynomial (curve-fit) can be used to relate the specific heat at constant pressure of dry air,  $c_p$  ( $kJ/kgK$ ) to temperature  $T$  for a range of values:

$$c_p = 0.99403 + 1.671 \times 10^{-4}T + 9.7215 \times 10^{-8}T^2 \\ - 9.5838 \times 10^{-11}T^3 + 1.9520 \times 10^{-14}T^4$$

Use a fixed point iteration scheme to determine the temperature that corresponds to specific heat of  $1.2$  ( $kJ/kgK$ ) with a convergence criteria of  $(\epsilon_n)_r = \frac{|T_{n+1}-T_n|}{|T_n|} < 10^{-5}$ . Show the derivation of the fixed-point function  $g(T)$  used in your scheme. Solve the problem with two different initial guesses,  $T_0 = 500$   $K$  and  $T_0 = 2000$   $K$ . For each case, tabulate the values of  $T_n$ ,  $T_{n+1}$ , and  $(\epsilon_n)_r$  at each iteration level  $n$  and limit the maximum number of iterations to  $N_{max} = 100$ .