

# THESIS INTRODUCTION

TIM LUCAS. JANUARY 14, 2016

## 1. ABSTRACT

**ABSTRACT.** Pathogens make up a huge proportion of global diversity and their role in disease strongly affects human disease, economics and development as well as having an important ecological role. However, the factors that control the number of pathogen species are poorly understood.

The patterns of contacts between individuals — in both human and animal populations — are nonrandom and depend on the density of individuals. Population structure and density have important epidemiological consequences, but their role in the control of pathogen richness is unknown.

It is still unknown whether population structure and density controls pathogen richness in bats. Furthermore, the specific mechanisms by which structure might influence pathogen richness have not been studied.

In this thesis I have studied how population structure and density control pathogen richness using bats as a case study. I have used epidemiological simulations and comparative analyses of data from wild bat populations to show that population structure does not have a strong affect on pathogen richness. Using further simulations I have shown that the interaction between population density and population structure is an important consideration. Finally, I have created a model for estimating bat densities — previously an incredibly challenging task — using acoustic data. Together, these studies clarify the relative roles of population density and structure and facilitate further study of population density in bats as a driver of pathogen richness.

While theory previously predicted that population structure should increase pathogen richness, the expectation in the ecological literature was that population structure would decrease pathogen richness. My studies support neither of these views, instead suggesting that population structure does not have a strong affect in either direction.

## 2. INTRODUCTION

The diversity of pathogens is huge and largely unknown (Poulin 2014). Recent large studies have found tens (Anthony et al. 2013) or even hundreds of virus species in a single host species (Anthony et al. 2015). This suggests the number of mammalian viruses globally is of the order of hundreds of thousands of virus species (Anthony et al. 2013) while only three thousand species of virus, across all host groups, are currently described (King et al. 2011). Recent large databases include nearly 2,000 pathogens from approximately 400 wild animal hosts (Wardeh et al. 2015). Given that there are nearly 4,000 named mammal species (Wilson & Reeder 2005), the undiscovered diversity of pathogens is likely huge.

This diversity of pathogens presents a risk to human health. 60% of newly emerged disease are zoonotic (acquired from animals) with wild animals being the predominant source (Jones et al. 2008, Taylor et al. 2001, Woolhouse & Gowtage-Sequeria 2006). Zoonotic diseases can be extremely virulent: viruses such as Nipah (Luby et al. 2009), Ebola (Lefebvre et al. 2014) having case fatality rates over 50%. Furthermore these pathogens can have large economic costs (e.g. SARS is estimated to have cost \$40 billion (Knobler et al. 2004)). In particular these impacts can have huge affects on developing economies. For example the 2014 ebola epidemic caused both Liberia and Guinea to fall from positive to negative per capita growth rates (World Bank n.d., World Bank 2014). In general we can expect competition between pathogens. Competition between

pathogens can occur by different mechanisms: immunological mechanisms such as cross-immunity or shared immune response (Fenton & Perkins 2010) and ecological mechanisms such as removal of susceptible hosts by death (Rohani et al. 2003) or competition for internal host resources (Griffiths et al. 2014). Like ecological systems this competition leads us to expect competitive exclusions (Ackleh & Allen 2003, Ackleh & Salceanu 2014, Bremermann & Thieme 1989, Martcheva & Li 2013, Turner & Garnett 2002) so the diversity of parasites needs an explanation.

Surveillance of zoonotic diseases is crucial. In particular we want to categorise diseases before they spillover into humans (SARS was not identified until months into the pandemic for example (Drosten et al. 2003)) and we want to anticipate outbreaks. If we know there is an increased number of infections in host species, increased numbers of a species that is a known reservoir of a high risk zoonotic disease, or increased contacts between humans and a pathogen reservoir, we can prepare for a potential outbreak in that area. However, funds for zoonotic surveillance are limited and so efforts must be optimised. Knowing which species are likely to have many pathogens allows us to sample and identify potentially zoonotic viruses efficiently. Many factors have been suggested in having a role in controlling pathogen richness including body mass (Arneberg 2002, Kamiya et al. 2014, Poulin 1995), animal density (Arneberg 2002, Kamiya et al. 2014, Nunn et al. 2003) and species range size (Kamiya et al. 2014, Nunn et al. 2003). This understanding provides a basis for predicting which species will have high pathogen richness and should be prioritised for sampling and surveillance.

In recent decades bats have been implicated in a number of high profile zoonotic outbreaks including Nipah (Field et al. 2001), Ebola (Leroy et al. 2005), SARS (Li et al. 2005) and Hendra (Field et al. 2001). This has led to much research on whether bats are a particular source of zoonotic disease (Luis et al. 2013, Olival et al. 2015, Wang et al. 2011) and examinations of factors, such as flight, social living and longevity, that might predispose them to being reservoirs of zoonotic viruses (Calisher et al. 2006, Dobson 2005, O'Shea et al. 2014). Given that bats are the second largest order of mammals (Wilson & Reeder 2005), we may expect them to be the source of many viruses simply through weight of numbers (Luis et al. 2013). The broad conclusions are that while bats do host more zoonotic viruses than other groups (Luis et al. 2013) they do not host more virus species in general (Olival et al. 2015). Questions remain as to why bat viruses have a tendency to have such high zoonotic potential.

Define. Simple count of pathogens across whole species. Ignores heterogeneity in prevalence, across space and time.

#### REFERENCES

- Ackleh, AS & LJ Allen (2003). "Competitive exclusion and coexistence for pathogens in an epidemic model with variable population size". *Journal of mathematical biology* 47.2, pp. 153–168. DOI: 10.1007/s00285-003-0207-9.
- Ackleh, AS & PL Salceanu (2014). "Robust uniform persistence and competitive exclusion in a nonautonomous multi-strain SIR epidemic model with disease-induced mortality". *Journal of mathematical biology* 1-2, pp. 453–475. DOI: 10.1007/s00285-012-0636-4.
- Anthony, SJ, JH Epstein, KA Murray, I Navarrete-Macias, CM Zambrana-Torrel, A Solovyov, R Ojeda-Flores, NC Arrigo, A Islam, SA Khan, P Hosseini, TL Bogich, KJ Olival, MD Sanchez-Leon, WB Karesh, T Goldstein, SP Luby, SS Morse, JAK Mazet, P Daszak, & WI Lipkin (2013). "A strategy to estimate unknown viral diversity in mammals". *MBio* 4.5, e00598–13.
- Anthony, SJ, A Islam, C Johnson, I Navarrete-Macias, E Liang, K Jain, PL Hitchens, X Che, A Solovyov, AL Hicks, R Ojeda-Flores, C Zambrana-Torrel, W Ulrich, MK Rostal, A Petrosov, J Garcia, N Haider, N Wolfe, T Goldstein, SS Morse, M Rahman, JH Epstein,

- JK Mazet, P Daszak, & WI Lipkin (2015). "Non-random patterns in viral diversity". *Nature communications* 6. DOI: 10.1038/ncomms9147.
- Arneberg, P (2002). "Host population density and body mass as determinants of species richness in parasite communities: comparative analyses of directly transmitted nematodes of mammals". *Ecography* 1, pp. 88–94. DOI: 10.1034/j.1600-0587.2002.250110.x.
- Bremermann, HJ & H Thieme (1989). "A competitive exclusion principle for pathogen virulence". *Journal of mathematical biology* 27, pp. 179–190. DOI: 10.1007/BF00276102.
- Calisher, C, J Childs, H Field, K Holmes, & T Schountz (2006). "Bats: important reservoir hosts of emerging viruses". *Clinical Microbiology Reviews* 19.3, pp. 531–545. DOI: 10.1128/CMR.00017-06.
- Dobson, AP (2005). "What links bats to emerging infectious diseases?" *Science* 310.5748, pp. 628–629. DOI: 10.1126/science.1120872.
- Drosten, C, S Günther, W Preiser, S Van Der Werf, HR Brodt, S Becker, H Rabenau, M Panning, L Kolesnikova, RA Fouchier, et al. (2003). "Identification of a novel coronavirus in patients with severe acute respiratory syndrome". *New England Journal of Medicine* 348.20, pp. 1967–1976. DOI: 10.1056/NEJMoa030747.
- Fenton, A & SE Perkins (2010). "Applying predator-prey theory to modelling immune-mediated, within-host interspecific parasite interactions". *Parasitology* 137.06, pp. 1027–1038. DOI: 10.1017/s0031182009991788.
- Field, H, P Young, J Yob, J Mills, L Hall, & J Mackenzie (2001). "The natural history of Hendra and Nipah viruses". *Microbes and infection* 3.4, pp. 307–314. DOI: 10.1016/S1286-4579(01)01384-3.
- Griffiths, EC, AB Pedersen, A Fenton, & OL Petchey (2014). "Analysis of a summary network of co-infection in humans reveals that parasites interact most via shared resources". *Proceedings of the Royal Society of London B: Biological Sciences* 281.1782, p. 20132286. DOI: 10.1098/rspb.2013.2286.
- Jones, KE, NG Patel, MA Levy, A Storeygard, D Balk, JL Gittleman, & P Daszak (2008). "Global trends in emerging infectious diseases". *Nature* 451.7181, pp. 990–993. DOI: 10.3201/eid0903.020336.
- Kamiya, T, K O'Dwyer, S Nakagawa, & R Poulin (2014). "What determines species richness of parasitic organisms? A meta-analysis across animal, plant and fungal hosts". *Biological Reviews* 89.1, pp. 123–134. DOI: 10.1111/brv.12046.
- King, AM, MJ Adams, & EJ Lefkowitz (2011). *Virus taxonomy: classification and nomenclature of viruses: Ninth Report of the International Committee on Taxonomy of Viruses*. Vol. 9. San Diego, CA: Elsevier. ISBN: 978-0123846846.
- Knobler, S, S Lemon, A Mack, L Sivitz, & K Oberholtzer (2004). *Learning from SARS: Preparing for the Next Disease Outbreak: Workshop Summary*. National Academies Press Washington, DC, USA: DOI: 10.1016/j.socscimed.2006.08.004.
- Lefebvre, A, C Fiet, C Belpois-Duchamp, M Tiv, K Astruc, & LA Glélé (2014). "Case fatality rates of Ebola virus diseases: a meta-analysis of World Health Organization data". *Médecine et maladies infectieuses* 44.9, pp. 412–416. DOI: 10.1016/j.medmal.2014.08.005.
- Leroy, EM, B Kumulungui, X Pourrut, P Rouquet, A Hassanin, P Yaba, A Délicat, JT Paweska, JP Gonzalez, & R Swanepoel (2005). "Fruit bats as reservoirs of Ebola virus". *Nature* 438.7068, pp. 575–576.
- Li, W, Z Shi, M Yu, W Ren, C Smith, JH Epstein, H Wang, G Crameri, Z Hu, H Zhang, J Zhang, J McEachern, H Field, P Daszak, BT Eaton, S Zhang, & LF Wang (2005). "Bats are natural reservoirs of SARS-like coronaviruses". *Science* 310.5748, pp. 676–679. DOI: 10.1126/science.1118391.
- Luby, SP, MJ Hossain, ES Gurley, BN Ahmed, S Banu, SU Khan, N Homaira, PA Rota, PE Rollin, JA Comer, et al. (2009). "Recurrent zoonotic transmission of Nipah virus

- into humans, Bangladesh, 2001–2007”. *Emerging infectious diseases* 15.8, p. 1229. DOI: 10.3201/eid1508.081237.
- Luis, AD, DTS Hayman, TJ O’Shea, PM Cryan, AT Gilbert, JRC Pulliam, JN Mills, ME Timonin, CKR Willis, AA Cunningham, AR Fooks, CE Rupprecht, JLN Wood, & CT Webb (2013). “A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special?” *Proceedings of the Royal Society B: Biological Sciences* 280.1756, p. 20122753. DOI: 10.1098/rspb.2012.2753.
- Martcheva, M & XZ Li (2013). “Competitive exclusion in an infection-age structured model with environmental transmission”. *Journal of Mathematical Analysis and Applications* 408.1, pp. 225–246. DOI: 10.1016/j.jmaa.2013.05.064.
- Nunn, CL, S Altizer, KE Jones, & W Sechrest (2003). “Comparative tests of parasite species richness in primates”. *The American Naturalist* 162.5, pp. 597–614. DOI: 10.1086/378721.
- Olival, KJ, CC Weekley, & P Daszak (2015). “Are bats really “special” as viral reservoirs? What we know and need to know”. In: *Bats and Viruses: A New Frontier of Emerging Infectious Diseases*. Ed. by LF Wang & C Cowled. New Jersey: Wiley, pp. 281–294. DOI: 10.1002/9781118818824.ch11.
- O’Shea, TJ, PM Cryan, AA Cunningham, AR Fooks, DT Hayman, AD Luis, AJ Peel, RK Plowright, & JL Wood (2014). “Bat flight and zoonotic viruses”. *Emerging infectious diseases* 20.5, p. 741. DOI: 10.3201/eid2005.130539.
- Poulin, R (1995). “Phylogeny, ecology, and the richness of parasite communities in vertebrates”. *Ecological Monographs*, pp. 283–302. DOI: 10.2307/2937061.
- Poulin, R (2014). “Parasite biodiversity revisited: frontiers and constraints”. *International journal for parasitology* 44.9, pp. 581–589. DOI: 10.1016/j.ijpara.2014.02.003.
- Rohani, P, C Green, N Mantilla-Beniers, & B Grenfell (2003). “Ecological interference between fatal diseases”. *Nature* 422.6934, pp. 885–888. DOI: 10.1038/nature01542.
- Taylor, LH, SM Latham, & E Mark (2001). “Risk factors for human disease emergence”. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 356.1411, pp. 983–989. DOI: 10.1093/acprof:oso/9780198525738.003.0035.
- Turner, K & G Garnett (2002). “The impact of the phase of an epidemic of sexually transmitted infection on the evolution of the organism”. *Sexually transmitted infections* 78, pp. i20–i30. DOI: 10.1136/sti.78.suppl\_1.i20.
- Wang, L, P Walker, & L Poon (2011). “Mass extinctions, biodiversity and mitochondrial function: are bats ‘special’ as reservoirs for emerging viruses?” *Current Opinion in Virology*. DOI: 10.1016/j.coviro.2011.10.013.
- Wardeh, M, C Risley, MK McIntyre, C Setzkorn, & M Baylis (2015). “Database of host-pathogen and related species interactions, and their global distribution”. *Scientific data* 2. DOI: 10.1038/sdata.2015.49.
- Wilson, DE & DM Reeder (2005). *Mammal species of the world: a taxonomic and geographic reference*. Vol. 12. JHU Press. ISBN: 9780801882210.
- Woolhouse, ME & S Gowtage-Sequeria (2006). “Host range and emerging and reemerging pathogens”. In: *Ending the War Metaphor:: The Changing Agenda for Unraveling the Host-Microbe Relationship-Workshop Summary*. Vol. 192. National Academies Press.
- World Bank. “World Development Indicators: Size of the economy”. URL: <http://wdi.worldbank.org/table/1.1>.
- World Bank (2014). *Update on the economic impact of the 2014 Ebola epidemic on Liberia, Sierra Leone, and Guinea*. Washington, DC. DOI: 10.1211/pj.2014.20066832. URL: <http://documents.worldbank.org/curated/en/2014/12/20454884/update-economic-impact-2014-ebola-epidemic-liberia-sierra-leone-guinea>.