

# Reminder of Random Variables I

Dr Ahmad Khonsari

ECE Dept.

The University of Tehran

## 4. Random variables (is nor random nor variable)

### Basic notes:

- events: sets of outcomes of the experiment;
- in many experiments we are interested in some number associated with the experiment:
- **random variable**: function which associates a number with experiment.

### Examples:

- number of voice calls  $N$  that exists at the switch at time  $t$ :
  - random variable which takes on integer values in  $(0, 1, \dots, \infty)$ .
- service time  $t_s$  of voice call at the switch:
  - random variable which takes on any real value  $(0, \infty)$ .

### Classification based on the nature of RV:

- continuous:  $R \in (-\infty, \infty)$
- discrete:  $N \in \{0, 1, \dots\}$ ,  $Z \in \{\dots, -1, 0, 1, \dots\}$ .

## 4.1. Definitions (measure theoretic)

**Definition:** a real valued RV  $X$  is a mapping from  $\Omega$  to  $\mathbb{R}$  such that:

$$\{\omega \in \Omega : X(\omega) \leq x\} \in \mathcal{F} \quad (45) \quad \text{for all } x \in \mathbb{R};$$

- This means that once we know the (random) value  $X(\omega)$  we know which of the events in  $\mathcal{F}$  have happened.
  - $\mathcal{F} = \{\emptyset, \Omega\}$ : only constant functions are measurable
  - $\mathcal{F} = 2^\Omega$ : all functions are measurable

**Definition:** an integer valued RV  $X$  is a mapping from  $\Omega$  to  $\mathbb{Z}$  such that:

$$\{\omega \in \Omega : X(\omega) \leq x\} \in \mathcal{F} \quad (46)$$

- for all  $x \in \mathbb{Z}$ ;

**Note!** in teletraffic and queuing theories:

- most RVs are time intervals, number of channels, packets etc.
- continuous:  $(0, \infty)$ , discrete:  $0, 1, \dots$

## 4.1. Definitions Random Variable (classic)

- We are often more interested in a some number associated with the experiment rather than the outcome itself.
- Example 1. The number of heads in tossing coin rather than the sequence of heads/tails

A real-valued random variable  $X$  is a mapping

$$X : \mathcal{S} \rightarrow \mathcal{R}$$

which associates the real number  $X(e)$  to each outcome  $e \in \mathcal{S}$ .

- **The image of a random variable  $X$**
- $\mathcal{S}_X = \{x \in \mathcal{R} \mid X(e) = x, e \in \mathcal{S}\}$  (complete set of values  $X$  can take)
- may be finite or countably infinite: discrete random variable :  $0, 1, \dots$
- uncountably infinite: continuous random variable :  $(0, \infty)$

## 4.1. Definitions Random Variable (classic)

- **Example 2:** The number of heads in three consecutive tossings of a coin (head = **h**, tail=**t** (tail)) .

$e$	$X(e)$
hhh	3
hht	2
hth	2
htt	1
thh	2
tht	1
tth	1
ttt	0

- The values of  $X$  are “drawn” by “drawing”  $e$
- $e$  represents a “lottery ticket”, on which the value of  $X$  is written

- **Note!**
- in teletraffic and queuing theories: most RVs are time intervals, number of channels, packets etc.

## 4.2. Full descriptors(PDF, pdf, pmf)

**Definition:** the probability that a random variable  $X$  is not greater than  $x$ :

$\Pr\{X \leq x\}$  = probability of the **Event**  $\{X \leq x\}$   
=function of  $x = F_X(x)$  with  $(-\infty \leq x \leq \infty)$



is called probability (cumulative) distribution function (PDF, CDF) of  $X$ .

**Definition:** complementary (cumulative) probability distribution function (CDF, CCDF)

- $F^C(x) = \Pr\{X > x\} = 1 - F(x) = G(x) \quad (48)$

# Cumulative Distribution Function

## -Example-



## 4.3. Properties of PDF

**For PDF the following properties holds:**

- PDF  $F(x)$  is monotone and non-decreasing with:

$$F(-\infty) = 0, F(\infty) = 1, 0 \leq F(x) \leq 1 \quad (51)$$

- for any  $a < b$ :

$$\Pr\{a < X \leq b\} = F(b) - F(a) \quad (52)$$

- right continuity: if  $F(x)$  is **discontinuous** at  $x = a$ , then:

$$F(a) = F(a - 0) + \Pr\{X = a\} \quad (53)$$

- If  $X$  is continuous:  $F(x) = \int_{-\infty}^x f(y)dy$

**Definition:** if  $X$  is a continuous RV, and  $F(x)$  is differentiable, then:


$$f(x) = \frac{dF(x)}{dx} = \lim_{dx \rightarrow 0} \frac{\Pr\{x < X \leq x + dx\}}{dx}$$

is called **probability density function** (pdf).





- $X$  is discrete:  $F(x) = \sum_{j \leq x} \Pr\{X = j\} \quad (54)$

**Note:** if  $X$  is discrete RV it is often preferable to deal with pmf (probability mass function) instead of PDF.





# Perf Eval of Comp Systems



# Perf Eval of Comp Systems




Fig.2-1

(a) The probability distribution and  
(b) The distribution function of a discrete RV.

## 4.4. Discrete RVs

- **Definition:** Let the values that can be assumed by  $X$  be  $x_k$ ,  $k = 0, 1, 2, \dots$
- The distribution function will have the staircase
- The steps occur at each  $x_k$  and have size  $P(X = x_k)$ .




Fig. A discrete distribution function has a finite number of discontinuities. The random variable has a nonzero probability only at the points of discontinuity.

## 4.4. Discrete RVs

CDF and pdf of discrete case

$$\begin{aligned} F_X(x) &= \Pr\{X \leq x\} \\ &= \sum_{j \leq x} \Pr\{X = j\} \\ &= \sum_{j=1}^N \Pr\{X = x_j\} u(x - x_j) \\ &= \sum_{j=1}^N p(x_j) u(x - x_j) \end{aligned}$$

, where  $p(x_j)$  is a shorthand for  $\Pr\{X = x_j\}$

**Note: accumulates  
up to  $x_j$ , and not to  $N$**





Fig. Discrete distribution and density functions

#### 4.4. Discrete RVs (pdf) !

$$\begin{aligned} f_X(x) &= \frac{F_X(x)}{dx} \\ &= \sum_{j=1}^N \Pr\{X = x_j\} \frac{du(x-x_j)}{dx} \\ &= \sum_{j=1}^N \Pr\{X = x_j\} \delta(x - x_j) \\ &= \sum_{j=1}^N p(x_j) \delta(x - x_j) \\ &= p(x_j) \text{ for } j=1, \dots, N \end{aligned}$$

Q: what is pmf of a discrete RV:



## 4.5. More Properties of pdf (continuous RV)

- pdf  $f(x)$  non-negative:

$$f(x) \geq 0, \quad x \in (-\infty, \infty) \quad (55)$$

- if  $f(x)$  is integrable then for any  $x_1 < x_2$ :

$$\Pr\{x_1 < X \leq x_2\} = F(x_2) - F(x_1)$$



$$= \int_{x_1}^{x_2} f(x) dx$$

- $F_X(x_0) = \int_{-\infty}^{x_0} f_X(x) dx$

- integration to 1:  $\int_{-\infty}^{\infty} f(x) dx = 1 \quad (57)$

**Note:** all these properties hold for pmf (you have to replace integral by sum).

**Q:** what does  $f(x)$  mean?



Note: **Not** All Continuous Random Variables Have PDFs , e.g. *Cantor set*

- <https://blogs.ubc.ca/math105/continuous-random-variables/the-pdf/>


## 4.6. mixed RVs

**Definition:**  $X$  is a continuous RV, and  $F(x)$  is differentiable, and with discontinuities at some discrete points:

The first term r.h.s are impulse components and the second is non-impulse component

$$f_X(x) = \sum_{j=1}^n p_j \delta(x - x_j) + \mathcal{C}_X(x)$$

$$\int_{-\infty}^{\infty} f_X(x) dx = \sum_{j=1}^n p_j U(x - x_j) + \int_{-\infty}^{\infty} \mathcal{C}(x) dx = 1$$





## 4.7. notes on Full descriptors cntd.

In what follows we assume integer values for discrete RVs i.e. :

$$p_j = \Pr\{X = j\} \quad (50)$$

Which is also called probability function (PF) or probability mass function (pmf).

- Q: X is a continuous RV with no jump, then  $P(x=x_0)=0$  or
- If we are ignorant:  $p(x \approx x_0) = f_X(x_0)|\Delta x|$  since

$$P\{x_0 < X(\xi) \leq x_0 + \Delta x\} = \int_{x_0}^{x_0 + \Delta x} f_X(u) du \approx f_X(x_0) \cdot \Delta x$$

- jumps in the CDF correspond to points x for which  $P(X=x)>0$

## 4.8. Parameters of RV

### **Basic notes:**

### **Full descriptors (i.e.)**

- continuous RV: PDF and pdf give all information regarding properties of RV;
- discrete RV: PDF and pdf(pmf) give all information regarding properties of RV.

### **Why we need something else:**

- problem 1: PDF, pdf and pmf are sometimes not easy to deal with;
- problem 2: sometimes it is hard to estimate from data;
- solution: use parameters (summaries) of RV.

### **What parameters (summaries):**

- mean, median;
- variance;
- skewness;
- excess (also known as excess kurtosis or simply kurtosis).

## 4.9-a: Mean

**Definition:** the mean of RV X is given by:

$$E[X] = \sum_{\forall i} x_i p_i, \quad E[x] = \int_{-\infty}^{\infty} x f(x) dx \quad (58)$$

- mean  $E[X]$  of RV X is between max and min value of non-complex RV:

$$\min_k x_k \leq E[x] \leq \max_k x_k \quad (59)$$

- mean of the constant is constant:

$$E[c] = c \quad (60)$$

- mean of RV multiplied by constant value is constant value multiplied by the mean:

$$E[cX] = cE[X] \quad (61)$$

- mean of constant and RV X is the mean of X and constant value:

$$E[c + X] = c + E[X] \quad (62)$$

- Linearity of Expectation:

$$E[X_1 + \cdots + X_n] = E[X_1] + \cdots + E[X_n]$$

## 4.9-a. Conditional Expectation

The expectation of the random variable  $X$  given that another random variable  $Y$  takes the value  $Y = y$  is

$$E[X|Y = y] = \int_{-\infty}^{\infty} x f_{X|Y}(x, y) dx$$

obtained by using the conditional distribution of  $X$ .

$E[X|Y = y]$  is a function of  $y$ .

By applying this function on the value of the random variable  $Y$  one obtains a random variable  $E[X/Y]$  (a function of the random variable  $Y$ ).

*Properties of conditional expectation*

$$E[X/Y] = E[X]$$

if  $X$  and  $Y$  are independent

$$E[cX/Y] = c E[X/Y]$$

$c$  is constant

$$E[X + Y/Z] = E[X/Z] + E[Y/Z]$$

$$E[g(Y)/Y] = g(Y)$$

$$E[g(Y)X/Y] = g(Y)E[X/Y]$$

## 4.9-b: median

- **Definition:** The *median* of  $X$  is defined to be any value  $m$  such that

$$\Pr(X \leq m) \geq 1/2 \text{ and } \Pr(X \geq m) \geq 1/2.$$

- **Theorem 3.9-mitzen:** For any random variable  $X$  with finite expectation  $\mathbf{E}[X]$  and finite median  $m$ ,

*1. the expectation  $\mathbf{E}[X]$  is the value of  $c$  that minimizes the expression*

$$\mathbf{E}[(X - c)^2], \text{ and}$$

*2. the median  $m$  is a value of  $c$  that minimizes the expression*

$$[|X - c|].$$

- **Theorem 3.10-mitzen:** If  $X$  is a random variable with finite standard deviation  $\sigma$ , expectation  $\mu$ , and median  $m$ , then

$$|\mu - m| \leq \sigma.$$

For a random variable  $X$ , consider the function

$$g(c) = E[(X - c)^2] \quad (3.57)$$

Remember, the quantity  $E[(X - c)^2]$  is a number, so  $g(c)$  really is a function, mapping a real number  $c$  to some real output.

We can ask the question, What value of  $c$  minimizes  $g(c)$ ? To answer that question, write:

$$g(c) = E[(X - c)^2] = E(X^2 - 2cX + c^2) = E(X^2) - 2cEX + c^2 \quad (3.58)$$

where we have used the various properties of expected value derived in recent sections.

Now differentiate with respect to  $c$ , and set the result to 0. Remembering that  $E(X^2)$  and  $EX$  are constants, we have

$$0 = -2EX + 2c \quad (3.59)$$

so the minimizing  $c$  is  $c = EX$ !

In other words, the minimum value of  $E[(X - c)^2]$  occurs at  $c = EX$ .

## 4.10. Variance and standard deviation

**Definition:** the mean of the square of difference between RV  $X$  and its mean  $E[X]$ :

$$V[X] = E[(X - E[X])^2] \quad (63)$$

**How to compute variance:**

- assume that  $X$  is discrete, compute variance as:

$$V[X] = \sum_{\forall n} (X - E[X])^2 p_n \quad (64)$$

- assume that  $X$  is continuous, compute variance as:

$$V[X] = \int_{-\infty}^{\infty} (x - E[X])^2 f(x) dx \quad (65)$$

- the another approach to compute variance:

$$V[X] = E[X^2] - (E[X])^2 \quad (66)$$

## 4.10 cntd. Properties of the variance:

- the variance of the constant value is 0:

$$V[c] = E[(X - E[X])^2] = E[(c - c)^2] = E[0] = 0 \quad (67)$$

- variance of RV multiplied by constant value:

$$V[cX] = E[(cX - cE[X])^2] = E[c^2(X - E[X])^2] = c^2V[X] \quad (68)$$

- variance of the constant value and RV X:

$$\begin{aligned} V[c + X] &= E[((c + X) - E(c + E[X]))^2] = E[(c + X - (c + E[X]))^2] \\ &= E[(X - E[X])^2] = V[X] \end{aligned} \quad (69)$$

**Definition:** the standard deviation of RV X is given by:

$$\sigma[X] = \sqrt{V[X]} \quad (70)$$

**Note:** standard deviation is dimensionless parameter.



# Perf Eval of Comp Systems

## 4.10 cntd. Properties of variance (summary):

- $V[X_1 + \cdots + X_n] = V[X_1] + \cdots + V[X_n]$  only when the  $X_i$  are independent
- $V[X_1 + \cdots + X_n] = \sum_{i,j=1}^n \text{Cov}[X_i, X_j]$  always

Proof:

$$\begin{aligned} \bullet \quad V[X_1 + \cdots + X_n] &= E\{(\sum_{j=1}^n (X_j - E(X_j)))^2\} \\ &= E\{\sum_{j=1}^n (X_j - E(X_j)) \sum_{k=1}^n (X_k - E(X_k))\} \\ &= \sum_{j=1}^n \sum_{k=1}^n E\{(X_j - E(X_j))(X_k - E(X_k))\} \\ &= \sum_{j,k=1}^n \text{Cov}[X_j, X_k] = \sum_{k=1}^n V(X_k) \\ &\quad + \sum_{j=1}^n \sum_{k=1}^n \text{Cov}(X_j, X_k) \end{aligned}$$

Properties of covariance

- $\text{Cov}[X, Y] = \text{Cov}[Y, X]$
- $\text{Cov}[X + Y, Z] = \text{Cov}[X, Z] + \text{Cov}[Y, Z]$

# Perf Eval of Comp Systems

## 4.10 cntd. Conditional variance

### Conditional variance

$V[X|Y] = E[(X - E[X|Y])^2|Y]$       Deviation with respect to the conditional expectation

### Conditional covariance

$COV[X, Y|Z] = E[(X - E[X|Z])(Y - E[Y|Z])|Z]$

### Conditioning rules

$E[X] = E[E[X|Y]]$  (inner conditional expectation is a function of  $Y$ )

$V[X] = E[V[X|Y]] + V[E[X|Y]]$       **Law of Total Variance**

$COV[X, Y] = E[COV[X, Y|Z]] + COV[E[X|Z], E[Y|Z]]$

$$E[E[Y|X]] = E[Y]$$

$$E[E[h(Y)|X]] = E[h(Y)]$$

$$E[E[Y^k|X]] = E[Y^k] \quad : \quad \text{على التخصيص } h(Y) = Y^k$$

ثانيًا:  $E[Y] = \int_{-\infty}^{\infty} E[Y|x] f_X(x) dx$

$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_Y(y|x) dy f_X(x) dx$$

$$= \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} \frac{f_{X,Y}(x,y)}{f_Y(y)} dx dy = \int_{-\infty}^{\infty} y f_Y(y) dy = E[Y]$$

269 مسألة 6°

$$E[h(X)g(X)] = \int_{-\infty}^{\infty} \frac{h(u)g(u)f_X(u)du}{\text{توزیع } X}$$

سنگین ص ۱۷۵

اگر  $g(x) = E[Y|X=x]$

$$= \int_{-\infty}^{\infty} h(u) E[Y|X=u] f_X(u) du$$

$$= \int_{-\infty}^{\infty} h(u) \int_{-\infty}^{\infty} y f_{Y|X}(y|u) f_X(u) dy du$$

$$= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(u) y f_{XY}(u, y) dy du = E[h(X)Y]$$

$$E[Y] = E_X[E_Y[Y|X=x]]$$

توزیع  $X$  ← هر مقدار از  $h(u)=1$  به همان وزن صرف می‌کند

در این شرط:

$$\text{Var}(X|Y=y) = E[(X - E(X|Y=y))^2 | Y=y]$$

نیز در این شکل:

$$\text{Var}(X) = E(\text{Var}(X|Y)) + \text{Var}(E(X|Y))$$

اثبات: می‌توانیم از این روش استفاده کنیم:

$$X - E(X) = (X - E(X|Y)) + (E(X|Y) - E(X))$$

حالا دو طرف را در هم ضرب می‌کنیم و امید می‌گیریم. به این ترتیب داریم:

$$\begin{aligned} \text{Var}(X) = E(X - E(X))^2 &= E((X - E(X|Y))^2) + E((E(X|Y) - E(X))^2) \\ &\quad + 2E((X - E(X|Y))(E(X|Y) - E(X))) \end{aligned}$$

طبق قانون امید می‌گیریم، جداگانه:

$$E(E((X - E(X|Y))^2 | Y))$$

در این شرط است،  $\text{Var}(X|Y)$

همچنین هم بداند  $\text{Var}(E(X|Y))$  است. چون  $E(X)$  یک عدد ثابت است.

$$E((E(X|Y) - E(X))^2) = \text{Var}(E(X|Y))$$

$$\text{Var}(X) = E[(X - E(X))^2]$$

حالا می‌توانیم بنویسیم که این دو طرف را در هم ضرب می‌کنیم:

در اینجا از فرمول  $h(Y) = 2(E(X|Y) - E(X))$  داریم. (نکته)

$$\begin{aligned} E[(X - E(X|Y))h(Y)] &= E(Xh(Y)) - E(E(X|Y)h(Y)) \\ &= E(Xh(Y)) - E(E(Xh(Y)|Y)) \\ &= E(Xh(Y)) - E(Xh(Y)) = 0 \end{aligned}$$

## 4.11. Other parameters: moments

**Let us assume the following:**

- $X$  be RV (discrete or continuous);
- $k \in 1, 2, \dots$  be the natural number;
- $Y = X^k, k = 1, 2, \dots$ , be the set of random variables.

**Definition:** the mean of RVs  $Y$  can be computed as follows:

- assume  $X$  is a discrete RV:

$$E[Y] = \sum_{\forall i} x_i^k p_i \quad (71)$$

- assume  $X$  is a continuous one.

$$E[Y] = \int_{-\infty}^{\infty} x^k f_X(x) dx \quad (72)$$

**Note:** for example, mean is obtained by setting  $k = 1$ .

**Definition:** (**raw**) **moment** of order  $k$  of RV  $X$  is the mean of RV  $X$  in power of  $k$ :

$$\alpha_k = E[X^k] \quad (73)$$

**Definition:** **central** moment (moment around the mean) of order  $k$  of RV  $X$  is given by:

$$\mu_k = E[(X - E[X])^k] \quad (74)$$


One can note that:

$$E[X] = \alpha_1, \quad V[X] = \sigma[X] = \mu_2 = \alpha_2 - \alpha_1^2 \quad (75)$$


measures of shape:

**Definition:** skewness (the degree of symmetry in the variable distribution) of RV is given by:


$$S_X = \frac{\mu_3}{(\sigma[X])^3} \quad (76)$$



Negatively skewed distribution  
or Skewed to the left  
Skewness < 0




Normal distribution  
Symmetrical  
Skewness = 0



Positively skewed distribution  
or Skewed to the right  
Skewness > 0

for **unimodal** (one peak), **skewed** to one side (i.e. not **symmetric**), If the bulk of the data is at the left and the right tail is longer, we say that the distribution is **skewed right or positively skewed**; and vice versa.

**Application:** three bandit (robbing your money) with the above distributions; the left distribution is the best Machine in terms of maximizing your net profit



Beta( $\alpha=4.5$ ,  
 $\beta=2$ )  
skewness =  
+0.5370



Skewness gives us the Shape of the data. It is the 'Lack of Symmetry'

### Positively Skewed


- Right Tail is longer
  - Mass of the distribution is concentrated on the left
- $\text{Mode} < \text{Median} < \text{Mean}$



### Symmetric


- Both tails are equal
  - Mass of the distribution is equally distributed
- $\text{Mean} = \text{Median} = \text{Mode}$

Normal Distribution is symmetric distribution




### Negatively Skewed

- Left Tail is longer
  - Mass of the distribution is concentrated on the right
- $\text{Mean} < \text{Median} < \text{Mode}$



Uniform( $\min=-\sqrt{3}$ ,  $\max=\sqrt{3}$ )  
kurtosis = 1.8, excess = -1.2



## measures of shape:

**Definition:** kurtosis (excess of kurtosis )

of RV is given by:


$$e_X = \frac{\mu_4}{(\sigma[X])^4} \quad (77)$$

the degree of tailedness in the variable distribution (Westfall 2014).


increasing kurtosis is associated with the “movement of probability mass from the shoulders of a distribution into its center and tails.”



Platykurtic  
distribution  
Thinner tails  
Kurtosis < 0





Normal  
distribution  
Mesokurtic  
distribution  
Kurtosis = 0



Leptokurtic  
distribution  
Fatter tails  
Kurtosis > 0

Kurtosis is defined as a measure of 'peakedness'. It is generally measured relative to Normal distribution. (Which means 'excess of kurtosis' is measured)



a distribution with kurtosis approximately equal to 3 or excess of kurtosis=0 is called mesokurtic. A value of kurtosis less than 3 indicates a platykurtic distribution and a value greater than 3 indicates a leptokurtic distribution. A normal distribution is a mesokurtic distribution.

## 4.12. Meaning of moments





### Parameters meanings:

- measures of central tendency:
  - mean:  $E[X] = \sum_{\forall i} x_i p_i$
  - mode: value corresponding to the highest probability;
  - median: value that equally separates weights of the distribution.
- measures of variability:
  - variance:  $V[X] = E[(X - E[X])^2]$
  - standard deviation:  $\sqrt{V[X]}$
  - squared coefficient of variation(squared COV):  $k_X^2 = \frac{V[X]}{E[X]^2}$
- other measures:
  - skewness of distribution: skewness;
  - excess of the mode: excess.

**Note:** not all parameters exist for a given distribution!

Pareto distribution has no mean when  $\alpha \leq 1$

Pareto distribution has no variance when  $\alpha \in (1, 2]$



تمرین ۱: اثبات کنید امید یک متغیر تصادفی ~~متغیر تصادفی~~ متغیر تصادفی را در صورتی که  $\text{tail}$   $\leq$  ثابت است:

$$E(X) = \int_{-\infty}^{\infty} x f_X(x) dx = - \underbrace{x \int_x^{\infty} f_X(u) du}_{\substack{\text{حد } x=0 \text{ برابر } 0 \\ \text{حد } x=\infty \text{ برابر } 0}} \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} dx \underbrace{\int_x^{\infty} f_X(u) du}_{\substack{\downarrow \\ L(F_X)}} \quad \int u dv = uv - \int u dv$$

$n = u \Rightarrow dn = du$   
 $\int_x^{\infty} f_X(x) dx = \int_x^{\infty} f_X(u) du$

$$= \int_{-\infty}^{\infty} (1 - F_X(x)) dx$$

$$E(X) = \int_{-\infty}^{\infty} x f_X(x) dx = x \int_{-\infty}^x f_X(u) du \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} dx \int_{-\infty}^x f_X(u) du$$

$$= \int_{-\infty}^{\infty} F_X(x) dx$$

$$E(x) = \sum_{k=-\infty}^{\infty} k P_r(x=k) = \sum_{k=-\infty}^{-1} k P_r(x=k) + \sum_{k=0}^{\infty} k P(x=k) \quad \therefore \text{نصفه}$$

$$= \sum_{k=-\infty}^{-1} k \{P(X \leq k) - P(X \leq k-1)\} + \sum_{k=0}^{\infty} k \{P(X \geq k) - P(X \geq k+1)\}$$

$$= \sum_{k=-\infty}^{-1} k P(X \leq k) - \sum_{k=-\infty}^{-2} (k+1) P(X \leq k) + \sum_{k=1}^{\infty} k P(X \geq k) - \sum_{k=1}^{\infty} (k-1) P(X \geq k)$$

$$= \sum_{k=-\infty}^{-1} k p(x \leq k) - \sum_{k=-\infty}^{-2} k p(x \leq k) - \sum_{k=-\infty}^{-2} p(x \leq k) + \sum_{k=1}^{\infty} k p(x \geq k) - \sum_{k=1}^{\infty} k p(x \geq k) + \sum_{k=1}^{\infty} p(x \geq k)$$

$$= -p(x \leq -1) - \sum_{k=-2}^{\infty} p(x \leq k) + \sum_{k=1}^{\infty} p(x \geq k) = 0$$

Handwritten mathematical diagrams illustrating the derivation of the Fourier transform of a function  $p(x, z)$ . The diagrams show the complex plane with various contours and poles. Key labels include  $p(x, z)$ ,  $p(-x, z)$ ,  $\sum$ ,  $R = -\infty$ ,  $R = \infty$ , and  $R = 1$ . The final result is shown as a sum of residues:  $\textcircled{1} + \textcircled{2} + \textcircled{3} = \sum p(x, z)$ .

$$= \sum_{k=1}^{\infty} p(x_7, k) - \sum_{R=-\infty}^{-1} p(x_7, k)$$





\* اگر یک RV متناهی عددی غیر منفی را بگیریم پس از  $\{0, 1, 2, \dots\}$  و براندازی قرار است

$$E(X) = \sum_{i=1}^{\infty} p(X \geq i)$$

$$\begin{aligned} &= \sum_{i=1}^{\infty} p(X \geq i) = \sum_{i=1}^{\infty} \sum_{j=i}^{\infty} p(X=j) = \sum_{j=1}^{\infty} \sum_{i=1}^j p(X=j) \\ &= \sum_{j=1}^{\infty} j p(X=j) = E(X) \end{aligned}$$

$$= E(X)$$

$$\begin{aligned} & p(X=1) + p(X=2) + p(X=3) + \dots \\ & + p(X=2) + p(X=3) + \dots \\ & + p(X=3) + \dots \end{aligned}$$

$$p(X=1) + 2p(X=2) + 3p(X=3) + \dots$$

مثبت و پیوسته، اگر تغییر تعداد را بر حسب  $x$  مقادیر غیر منفی را بنویسیم (بگیریم) در آن صورت داریم:

(و مشتق تابع چگالی  $f_x(t)$  را به دست می آوریم)

$$E(x) = \int_0^{\infty} p(x, n) dn$$

$$= \int_0^{\infty} p(x, n) dn = \int_0^{\infty} \int_0^{\infty} \underbrace{f_x(t)}_{p(x, t)} dt dn = \int_0^{\infty} \int_0^{\infty} f_x(t) dx dt$$

$$\int_0^{\infty} f_x(t) dx = f_x(t)$$

$$= \int_0^{\infty} \underbrace{f_x(t)}_{(2)} dt = E(x)$$

متساوی

اگر چگالی  $f_x(t)$  را در نظر بگیریم:

$$E(x) = \int_0^{\infty} \int_0^{\infty} dt dF(t) =$$

$$\int_0^{\infty} \int_t^{\infty} dF(t) dt = \int_0^{\infty} (1 - F(t)) dt$$

**Theorem 4.1** (Continuous Tail Sum Formula). *Let  $X$  be a non-negative random variable. Then*

$$E(X) = \int_0^{\infty} (1 - F_X(x)) \, dx \quad (4.16)$$

*Proof.*

$$\begin{aligned} E(X) &= \int_0^{\infty} x f_X(x) \, dx \\ &= \int_0^{\infty} \int_0^x f_X(x) \, dt \, dx \\ &= \int_0^{\infty} \int_t^{\infty} f_X(x) \, dx \, dt \\ &= \int_0^{\infty} \Pr(X > t) \, dt \\ &= \int_0^{\infty} (1 - F_X(t)) \, dt \end{aligned}$$

The proof is quite similar to the discrete case. Interchanging the bounds of integration in line 3 is justified by Fubini's Theorem from multivariable calculus.  $\square$

**Theorem 2.2.5** *Let  $X$  be a non-negative continuous random variable with its distribution function  $F(x)$ . Suppose that  $\lim_{x \rightarrow \infty} x\{1 - F(x)\} = 0$ . Then, we have:*

$$E(X) = \sum_{x=0}^{\infty} \{1 - F(x)\}.$$

**Proof** We have assumed that  $X \geq 0$  w.p.1 and thus

$$\begin{aligned} E(X) &= \int_0^{\infty} x f(x) dx \\ &= \int_0^{\infty} x dF(x), \because dF(x)/dx = f(x) \text{ from (1.6.10)} \\ &= - \int_0^{\infty} x d\{1 - F(x)\} = \\ &\quad - \{[x\{1 - F(x)\}]_{x=0}^{x=\infty} - \int_0^{\infty} \{1 - F(x)\} dx\}, \\ &\quad \text{using integration by parts from (1.6.28)} \\ &= \int_0^{\infty} \{1 - F(x)\} dx \text{ since } \lim_{x \rightarrow \infty} x\{1 - F(x)\} \\ &\quad \text{is assumed to be zero.} \end{aligned}$$

The proof is now complete. ■

۲۰ در مورد استواریت شرطی - Conditioning rule: استواریت که در زیر آورده شده است

\* قانون استواریت \* (یا قانون استواریت تکرار یا تکرار برج یا قضیه هموار کننده)

اگر  $X$  متغیر تصادفی استوار پذیر باشد (یعنی  $E|X| < \infty$ ) و  $Y$  یک متغیر تصادفی که نزدیک استوار پذیر نیست و محدود و در حد تغییر احوال باشد (همچون داده (تکرار) آنها یکسان باشد)

داریم:

$$E(X) = E(E(X|Y)) \quad (1)$$

$$E(E(X|Y)) = \sum_y (E(X|Y=y) \cdot P(Y=y))$$

المطلوب (1):  $E(X|Y)$  هو متوسط  $E(X|Y)$  على  $Y$   
 بر  $E$ ،  $Y$  متغير عشوائي

$$= \sum_y \left( \sum_x x (P(X=x|Y=y)) \right) \cdot P(Y=y)$$

$$= \sum_y \sum_x x P(X=x|Y=y) \cdot P(Y=y)$$

$$= \sum_y \sum_x x P(X=x, Y=y)$$

$$= \sum_x \sum_y x P(Y=y|X=x) P(X=x)$$

$$= \sum_x x P(X=x) \left( \sum_y P(Y=y|X=x) \right)$$

$$= \sum_x x P(X=x)$$

$$= E(X)$$

$$E(X) = \sum_x x \sum_y P(X=x, Y=y) =$$

$$\sum_x x P(X=x) = E(X)$$

$$\sum_y P(Y=y|X=x) = \frac{\sum_y P(X=x, Y=y)}{P(X=x)} = 1$$

$$P(Y=y|X=x) = \frac{P(Y=y, X=x)}{P(X=x)}$$

موزع مشترك

$$P(X=x) = \sum_y P(X=x, Y=y)$$

ناتج



تاریک‌داری ۴:

$E(X|Y)$  عددی که متغیر تصادفی است چون مقدار آن بستگی به  $Y$  دارد، چون:

نویافته که مقدار این شرطی  $X$  به شرط  $Y=y$  تناسب از  $y$  است

• پس برای  $Y$  های مختلف،  $E(X|Y)$  مقدار تصادفی است (در این اصل متغیر تصادفی است)

• اگر بنویسیم  $E(X|Y=y)=g(y)$ ، پس متغیر تصادفی،  $E(X|Y)$  را می‌توان نوشت

$$E(X|Y) = \sum_x x \cdot P(X=x|Y=y)$$

$g(y)$  بیان کرد.