

# Gaussian Processes for Regression

COMP9418 — Advanced Topics in Statistical Machine Learning

**Edwin V. Bonilla**

School of Computer Science and Engineering  
UNSW Sydney




**UNSW**  
SYDNEY

September 20th, 2017

(Last Update: Tuesday 19<sup>th</sup> September, 2017 at 18:15)

# Acknowledgements




Carl Edward Rasmussen and Christopher K. I. Williams

All chapters available online along with software and datasets:  
<http://www.gaussianprocess.org/gpml>


This lecture will allow you to understand Gaussian processes as priors over functions and apply them to regression problems. Following it you should be to:

- Understand and apply Bayesian approaches to linear regression.
- Understand and apply Bayesian linear-in-the-parameters models to non-linear regression problems.
- Understand the connection between Bayesian regression with non-linear feature spaces and Gaussian process regression.
- Derive and apply the function-space view of Gaussian process regression.
- Carry out model selection in Gaussian process regression models.


## Some Applications of Gaussian Process (GP) Models (1)




## Spatio-temporal modelling



## Classification




## Robot inverse dynamics



## Data fusion / multi-task learning


# Some Applications of GP Models (2)




Style-based inverse kinematics



Preference learning



Bayesian optimisation




Bayesian quadrature

# How can We 'Solve' All these Problems with the Humble Gaussian Distribution?

## Key components of GP models:

- Non-parametric prior
- Bayesian
- Kernels (covariance functions)




# How can We 'Solve' All these Problems with the Humble Gaussian Distribution?

## Key components of GP models:

- Non-parametric prior
- Bayesian
- Kernels (covariance functions)

## Tasks

- Prediction (posterior inference)
- Hyperparameter learning



# How can We 'Solve' All these Problems with the Humble Gaussian Distribution?

## Key components of GP models:


- Non-parametric prior
- Bayesian
- Kernels (covariance functions)

## Tasks

- Prediction (posterior inference)
- Hyperparameter learning

## Challenges


- Intractability for non-Gaussian likelihoods
  - ▶ E.g. a sigmoid likelihood for classification
- High computational cost (in time and memory) with  $\#$  datapoints






# The Prediction Problem

Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



# The Prediction Problem


Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



- What parameterization?

# The Prediction Problem


Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j x_j$

# The Prediction Problem


Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$

# The Prediction Problem


Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$
- Flexibility v generalization

# The Prediction Problem


Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$
- Flexibility v generalization
- What basis functions?  
How many?

# The Prediction Problem

Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .




- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$
- Flexibility v generalization
- What basis functions?  
How many?

- What about Neural nets?

# The Prediction Problem

Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .




- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$
- Flexibility v generalization
- What basis functions?  
How many?

- What about Neural nets?
- How to avoid overfitting? (cf regularization)



# The Prediction Problem

Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .




- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$
- Flexibility v generalization
- What basis functions?  
How many?

- What about Neural nets?
- How to avoid overfitting? (cf regularization)
- Confidence on our predictions?

# The Prediction Problem


Learn mapping  $\mathbf{x} \rightarrow f(\mathbf{x})$  from observations  $\{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ .



- What parameterization?
- $f(\mathbf{x}) = \sum_j w_j \phi_j(\mathbf{x})$
- Flexibility v generalization
- What basis functions?  
How many?

- What about Neural nets?
- How to avoid overfitting? (cf regularization)
- Confidence on our predictions?

We can address these issues in a principled way with Gaussian process models



- Smooth functions
- Closeness in input space  $\rightarrow$  closeness in output space

# Why Gaussian Processes

- Parametric models constrain the class of functions we consider
- Flexibility (no underfitting) due to **non-parametric** nature

# Why Gaussian Processes

- Parametric models constrain the class of functions we consider
- Flexibility (no underfitting) due to non-parametric nature
- Generalization (no overfitting)
- Bayesian, distribution over functions: prior, likelihood, posterior

# Why Gaussian Processes

- Parametric models constrain the class of functions we consider
- Flexibility (no underfitting) due to non-parametric nature
- Generalization (no overfitting)
- Bayesian, distribution over functions: prior, likelihood, posterior
- How can we do computations with infinite vectors?
  - ▶ “Efficient” Inference due to consistency (Gaussian distributions)

# Why Gaussian Processes

- Parametric models constrain the class of functions we consider
- Flexibility (no underfitting) due to **non-parametric** nature
- Generalization (no overfitting)
- **Bayesian**, **distribution over functions**: prior, likelihood, posterior
- How can we do computations with **infinite vectors**?
  - ▶ “Efficient” Inference due to **consistency** (Gaussian distributions)
- Characteristics of the functions can be learned from data
  - ▶ **Covariance function**: smoothness, stationarity, length-scale
  - ▶ Hyperparameter learning

# Why Gaussian Processes

- Parametric models constrain the class of functions we consider
- Flexibility (no underfitting) due to **non-parametric** nature
- Generalization (no overfitting)
- **Bayesian**, **distribution over functions**: prior, likelihood, posterior
- How can we do computations with **infinite vectors**?
  - ▶ “Efficient” Inference due to **consistency** (Gaussian distributions)
- Characteristics of the functions can be learned from data
  - ▶ **Covariance function**: smoothness, stationarity, length-scale
  - ▶ Hyperparameter learning
- Many standard regression models are special cases of GPs



# Why Gaussian Processes


- Parametric models constrain the class of functions we consider
- Flexibility (no underfitting) due to **non-parametric** nature
- Generalization (no overfitting)
- **Bayesian**, **distribution over functions**: prior, likelihood, posterior
- How can we do computations with **infinite vectors**?
  - ▶ “Efficient” Inference due to **consistency** (Gaussian distributions)
- Characteristics of the functions can be learned from data
  - ▶ **Covariance function**: smoothness, stationarity, length-scale
  - ▶ Hyperparameter learning
- Many standard regression models are special cases of GPs
- GP models also applicable to non-regression settings

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

# The Gaussian Distribution


## 1D Example




$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right)$$

# The Gaussian Distribution

## 1D Example




$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right)$$




$$F(x) = \int_{-\infty}^x \mathcal{N}(z|\mu, \sigma^2) dz$$

# The Gaussian Distribution

## 1D Example



$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right)$$




$$F(x) = \int_{-\infty}^x \mathcal{N}(z|\mu, \sigma^2) dz$$

In general:  $p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{|2\pi\boldsymbol{\Sigma}|^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right)$

# The Gaussian Distribution

## 2D Example




$$p(x_1, x_2) \sim \mathcal{N}(\mu, \Sigma)$$

Joint


# The Gaussian Distribution

## 2D Example



$$p(x_1, x_2) \sim \mathcal{N}(\mu, \Sigma)$$

Joint




$$p(x_1)$$

Marginal




# The Gaussian Distribution

## 2D Example




$$p(x_1, x_2) \sim \mathcal{N}(\mu, \Sigma)$$

Joint



$$p(x_1)$$

Marginal




$$p(x_1|x_2)$$

Conditional


# The Gaussian Distribution

## 2D Example




$$p(x_1, x_2) \sim \mathcal{N}(\mu, \Sigma)$$

Joint



$$p(x_1)$$

Marginal



$$p(x_1|x_2)$$

Conditional

The **marginal** and the **conditional** distributions are also Gaussians

# Partitioned Gaussians

For general Gaussian random vectors  $\mathbf{x}$ , we can partition:

$$\mathbf{x} \stackrel{\text{def}}{=} \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{12}^T & \boldsymbol{\Sigma}_{22} \end{bmatrix} \right),$$

# Partitioned Gaussians

For general Gaussian random vectors  $\mathbf{x}$ , we can partition:

$$\mathbf{x} \stackrel{\text{def}}{=} \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{12}^T & \boldsymbol{\Sigma}_{22} \end{bmatrix} \right),$$

hence, the **marginal**  $\mathbf{x}_1 \sim \mathcal{N}(\mathbf{x}_1 | \boldsymbol{\mu}_1, \boldsymbol{\Sigma}_{11})$

# Partitioned Gaussians

For general Gaussian random vectors  $\mathbf{x}$ , we can partition:

$$\mathbf{x} \stackrel{\text{def}}{=} \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{12}^T & \boldsymbol{\Sigma}_{22} \end{bmatrix} \right),$$

hence, the **marginal**  $\mathbf{x}_1 \sim \mathcal{N}(\mathbf{x}_1 | \boldsymbol{\mu}_1, \boldsymbol{\Sigma}_{11})$

and the **conditional**  $\mathbf{x}_1 | \mathbf{x}_2 \sim \mathcal{N}(\mathbf{x}_1 | \boldsymbol{\mu}_{1|2}, \boldsymbol{\Sigma}_{1|2})$

# Partitioned Gaussians

For general Gaussian random vectors  $\mathbf{x}$ , we can partition:

$$\mathbf{x} \stackrel{\text{def}}{=} \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{12}^T & \boldsymbol{\Sigma}_{22} \end{bmatrix} \right),$$

hence, the **marginal**  $\mathbf{x}_1 \sim \mathcal{N}(\mathbf{x}_1 | \boldsymbol{\mu}_1, \boldsymbol{\Sigma}_{11})$

and the **conditional**  $\mathbf{x}_1 | \mathbf{x}_2 \sim \mathcal{N}(\mathbf{x}_1 | \boldsymbol{\mu}_{1|2}, \boldsymbol{\Sigma}_{1|2})$

where  $\boldsymbol{\mu}_{1|2} = \boldsymbol{\mu}_1 + \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} (\mathbf{x}_2 - \boldsymbol{\mu}_2)$ ,

and  $\boldsymbol{\Sigma}_{1|2} = \boldsymbol{\Sigma}_{11} - \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} \boldsymbol{\Sigma}_{12}^T$

# The Gaussian Distribution

## Covariance and Precision Matrices

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{|2\pi\boldsymbol{\Sigma}|^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right)$$

$\boldsymbol{\Sigma}$  : is the **covariance** matrix

$\boldsymbol{\Sigma}^{-1}$  : is the **precision** matrix

# The Gaussian Distribution

## Covariance and Precision Matrices

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{|2\pi\boldsymbol{\Sigma}|^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right)$$

$\boldsymbol{\Sigma}$  : is the **covariance** matrix

$\boldsymbol{\Sigma}^{-1}$  : is the **precision** matrix

- An entry  $\boldsymbol{\Sigma}_{ij}^{-1} = 0$  indicates that the variables  $i$  and  $j$  are **conditionally independent** given all the other variables.



# The Gaussian Distribution

## Covariance and Precision Matrices

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{|2\pi\boldsymbol{\Sigma}|^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right)$$

$\boldsymbol{\Sigma}$  : is the **covariance** matrix

$\boldsymbol{\Sigma}^{-1}$  : is the **precision** matrix

- An entry  $\boldsymbol{\Sigma}_{ij}^{-1} = 0$  indicates that the variables  $i$  and  $j$  are **conditionally independent** given all the other variables.
- An entry  $\boldsymbol{\Sigma}_{ij} = 0$  indicates that the variables  $i$  and  $j$  are **marginally independent** given all the other variables.

# The Gaussian Distribution

## Covariance and Precision Matrices

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{|2\pi\boldsymbol{\Sigma}|^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right)$$

$\boldsymbol{\Sigma}$  : is the **covariance** matrix

$\boldsymbol{\Sigma}^{-1}$  : is the **precision** matrix

- An entry  $\boldsymbol{\Sigma}_{ij}^{-1} = 0$  indicates that the variables  $i$  and  $j$  are **conditionally independent** given all the other variables.
- An entry  $\boldsymbol{\Sigma}_{ij} = 0$  indicates that the variables  $i$  and  $j$  are **marginally independent** given all the other variables.
- Marginalizing out a variable leaves  $\boldsymbol{\Sigma}$  unchanged but changes  $\boldsymbol{\Sigma}^{-1}$ .
  - ▶ This is crucial when parameterizing a Gaussian process.

## Gaussian Quiz

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

# The Standard Linear Regression Model

## Notation and Settings

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** :  $\mathbf{x} \xrightarrow{f(\mathbf{x})} \mathbf{y}$

# The Standard Linear Regression Model

## Notation and Settings

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ ,  $\mathbf{x} \in \mathbb{R}^D$ ,  $y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** :  $\mathbf{x} \xrightarrow{f(\mathbf{x})} \mathbf{y}$

**Model**       $f(\mathbf{x}) = \sum_{j=1}^D w_j x_j \quad = \mathbf{w}^T \mathbf{x}$

**Noise**       $y = f(\mathbf{x}) + \eta \quad \text{with } \eta \sim \mathcal{N}(\eta|0, \sigma^2)$

**Likelihood**    $y|f(\mathbf{x}) \sim \mathcal{N}(y|f(\mathbf{x}), \sigma^2) \quad = \mathcal{N}(y|\mathbf{w}^T \mathbf{x}, \sigma^2)$

# The Standard Linear Regression Model

## Notation and Settings

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ ,  $\mathbf{x} \in \mathbb{R}^D$ ,  $y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** :  $\mathbf{x} \xrightarrow{f(\mathbf{x})} \mathbf{y}$

**Model**       $f(\mathbf{x}) = \sum_{j=1}^D w_j x_j \quad = \mathbf{w}^T \mathbf{x}$

**Noise**       $y = f(\mathbf{x}) + \eta$       with  $\eta \sim \mathcal{N}(\eta|0, \sigma^2)$

**Likelihood**    $y|f(\mathbf{x}) \sim \mathcal{N}(y|f(\mathbf{x}), \sigma^2) \quad = \mathcal{N}(y|\mathbf{w}^T \mathbf{x}, \sigma^2)$

Thus, the data-likelihood is given by:

$$\begin{aligned} p(\mathbf{y}|\mathbf{X}, \mathbf{w}) &= \prod_{n=1}^N p(y^{(n)}|\mathbf{x}^{(n)}, \mathbf{w}) = \prod_{n=1}^N \mathcal{N}(y^{(n)}|\mathbf{w}^T \mathbf{x}^{(n)}, \sigma^2) \\ &= \mathcal{N}(\mathbf{y}|\mathbf{X}^T \mathbf{w}, \sigma^2 \mathbf{I}) \end{aligned}$$

# The Standard Linear Regression Model

## Notation and Settings

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ ,  $\mathbf{x} \in \mathbb{R}^D$ ,  $y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** :  $\mathbf{x} \xrightarrow{f(\mathbf{x})} \mathbf{y}$

**Model**       $f(\mathbf{x}) = \sum_{j=1}^D w_j x_j = \mathbf{w}^T \mathbf{x}$

**Noise**       $y = f(\mathbf{x}) + \eta$       with  $\eta \sim \mathcal{N}(\eta|0, \sigma^2)$

**Likelihood**     $y|f(\mathbf{x}) \sim \mathcal{N}(y|f(\mathbf{x}), \sigma^2) = \mathcal{N}(y|\mathbf{w}^T \mathbf{x}, \sigma^2)$

Thus, the data-likelihood is given by:

$$\begin{aligned} p(\mathbf{y}|\mathbf{X}, \mathbf{w}) &= \prod_{n=1}^N p(y^{(n)}|\mathbf{x}^{(n)}, \mathbf{w}) = \prod_{n=1}^N \mathcal{N}(y^{(n)}|\mathbf{w}^T \mathbf{x}^{(n)}, \sigma^2) \\ &= \mathcal{N}(\mathbf{y}|\mathbf{X}^T \mathbf{w}, \sigma^2 \mathbf{I}) \end{aligned}$$

We need to do inference on  $\mathbf{w}$ .



# Bayesian Linear Regression

## Posterior Distribution

Consider a zero-mean Gaussian prior over the weights:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

# Bayesian Linear Regression

## Posterior Distribution

Consider a zero-mean Gaussian prior over the weights:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

Then the **posterior distribution** over the weights is given by:

$$\begin{aligned} p(\mathbf{w}|\mathbf{X}, \mathbf{y}) &= \frac{p(\mathbf{w}) p(\mathbf{y}|\mathbf{X}, \mathbf{w})}{p(\mathbf{y}|\mathbf{X})} \\ &= \mathcal{N}(\mathbf{w}|\bar{\mathbf{w}}, \mathbf{A}^{-1}) \end{aligned}$$

where  $\bar{\mathbf{w}} = \frac{1}{\sigma^2} \mathbf{A}^{-1} \mathbf{X} \mathbf{y}$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \mathbf{X} \mathbf{X}^T + \mathbf{\Sigma}_w^{-1})$ .

# Bayesian Linear Regression

## Posterior Distribution

Consider a zero-mean Gaussian prior over the weights:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

Then the **posterior distribution** over the weights is given by:

$$\begin{aligned} p(\mathbf{w}|\mathbf{X}, \mathbf{y}) &= \frac{p(\mathbf{w}) p(\mathbf{y}|\mathbf{X}, \mathbf{w})}{p(\mathbf{y}|\mathbf{X})} \\ &= \mathcal{N}(\mathbf{w}|\bar{\mathbf{w}}, \mathbf{A}^{-1}) \end{aligned}$$

where  $\bar{\mathbf{w}} = \frac{1}{\sigma^2} \mathbf{A}^{-1} \mathbf{X} \mathbf{y}$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \mathbf{X} \mathbf{X}^T + \mathbf{\Sigma}_w^{-1})$ .

- Mean of posterior is equal to its mode

# Bayesian Linear Regression

## Posterior Distribution

Consider a zero-mean Gaussian prior over the weights:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

Then the **posterior distribution** over the weights is given by:

$$\begin{aligned} p(\mathbf{w}|\mathbf{X}, \mathbf{y}) &= \frac{p(\mathbf{w}) p(\mathbf{y}|\mathbf{X}, \mathbf{w})}{p(\mathbf{y}|\mathbf{X})} \\ &= \mathcal{N}(\mathbf{w}|\bar{\mathbf{w}}, \mathbf{A}^{-1}) \end{aligned}$$

where  $\bar{\mathbf{w}} = \frac{1}{\sigma^2} \mathbf{A}^{-1} \mathbf{X} \mathbf{y}$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \mathbf{X} \mathbf{X}^T + \mathbf{\Sigma}_w^{-1})$ .

- Mean of posterior is equal to its mode
- MAP solution (non-Bayesian): negative log prior as penalty term

# Bayesian Linear Regression

## Posterior Distribution

Consider a zero-mean Gaussian prior over the weights:

$$\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

Then the **posterior distribution** over the weights is given by:

$$\begin{aligned} p(\mathbf{w}|\mathbf{X}, \mathbf{y}) &= \frac{p(\mathbf{w}) p(\mathbf{y}|\mathbf{X}, \mathbf{w})}{p(\mathbf{y}|\mathbf{X})} \\ &= \mathcal{N}(\mathbf{w}|\bar{\mathbf{w}}, \mathbf{A}^{-1}) \end{aligned}$$

where  $\bar{\mathbf{w}} = \frac{1}{\sigma^2} \mathbf{A}^{-1} \mathbf{X} \mathbf{y}$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \mathbf{X} \mathbf{X}^T + \mathbf{\Sigma}_w^{-1})$ .

- Mean of posterior is equal to its mode
- MAP solution (non-Bayesian): negative log prior as penalty term
- This penalized maximum likelihood is known as **ridge regression**
  - ▶ Consider  $\mathbf{\Sigma}_w = \lambda \mathbf{I}$  Then :

$$\bar{\mathbf{w}} = (\mathbf{X} \mathbf{X}^T + \frac{1}{\lambda} \sigma^2 \mathbf{I})^{-1} \mathbf{X} \mathbf{y}$$

# Bayesian Linear Regression

## Predictive Distribution

We are interested in making predictions at a new test point  $\mathbf{x}_*$

- In fact we obtain the **predictive distribution** by averaging over all possible parameter values (weighted by their posterior probabilities):

$$p(f_*|\mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \int p(f_*|\mathbf{x}_*, \mathbf{w})p(\mathbf{w}|\mathbf{X}, \mathbf{y}) d\mathbf{w} = \mathcal{N}(f_*|\mathbf{x}_*^T \bar{\mathbf{w}}, \mathbf{x}_*^T \mathbf{A}^{-1} \mathbf{x}_*)$$

- ▶ Predictive mean: linear combination of weights' posterior mean
- ▶ Predictive variance: grows with the magnitude of the test point

# Bayesian Linear Regression

## Predictive Distribution

We are interested in making predictions at a new test point  $\mathbf{x}_*$

- In fact we obtain the **predictive distribution** by averaging over all possible parameter values (weighted by their posterior probabilities):


$$p(f_*|\mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \int p(f_*|\mathbf{x}_*, \mathbf{w})p(\mathbf{w}|\mathbf{X}, \mathbf{y}) d\mathbf{w} = \mathcal{N}(f_*|\mathbf{x}_*^T \bar{\mathbf{w}}, \mathbf{x}_*^T \mathbf{A}^{-1} \mathbf{x}_*)$$

- ▶ Predictive mean: linear combination of weights' posterior mean
- ▶ Predictive variance: grows with the magnitude of the test point
- **Point predictions**: Need to consider the expected loss (or **risk**):

$$y_{\text{opt}} = \underset{y_{\text{pred}}}{\operatorname{argmin}} \int \mathcal{L}(f_*, y_{\text{pred}}) p(f_*|\mathbf{x}_*, \mathbf{X}, \mathbf{y}) df_*$$

- ▶ e.g. Square loss  $\mathcal{L} = (y_{\text{pred}} - f_*)^2$
- ▶ c.f. Empirical risk minimization (ERM)


# Bayesian Linear Regression Example




Prior Weights



# Bayesian Linear Regression Example




Prior Weights




Observed Data


# Bayesian Linear Regression Example



Prior Weights




Observed Data




Likelihood


# Bayesian Linear Regression Example




Prior Weights



Observed Data




Likelihood




Posterior Weights


# Bayesian Linear Regression Example




Prior Weights



Predictive Distribution



Likelihood



Posterior Weights

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

# Non-linear Feature Spaces

- Consider the model  $f(\mathbf{x}) = \sum_{i=1}^{D'} w_i \phi_i(\mathbf{x}) = \mathbf{w}^T \boldsymbol{\phi}(\mathbf{x})$ 
  - ▶ Each  $\phi_i(\mathbf{x})$  is a (non-linear) feature on  $\mathbf{x}$ , e.g.  $x_1, x_2, x_1^2, x_2^2, x_1 x_2 \dots$
  - ▶ We have a non-linear mapping but a **linear-in-the-parameters** model
  - ▶ The number of these features can be very large, i.e.  $D' \gg D$

# Non-linear Feature Spaces

- Consider the model  $f(\mathbf{x}) = \sum_{i=1}^{D'} w_i \phi_i(\mathbf{x}) = \mathbf{w}^T \boldsymbol{\phi}(\mathbf{x})$ 
  - ▶ Each  $\phi_i(\mathbf{x})$  is a (non-linear) feature on  $\mathbf{x}$ , e.g.  $x_1, x_2, x_1^2, x_2^2, x_1 x_2 \dots$
  - ▶ We have a non-linear mapping but a **linear-in-the-parameters** model
  - ▶ The number of these features can be very large, i.e.  $D' \gg D$
- All the Bayesian analysis is similar to the standard linear model:

$$p(f_* | \mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \mathcal{N}(f_* | \sigma^{-2} \boldsymbol{\phi}_*^T \mathbf{A}^{-1} \boldsymbol{\Phi} \mathbf{y}, \boldsymbol{\phi}_*^T \mathbf{A}^{-1} \boldsymbol{\phi}_*)$$

where:  $\boldsymbol{\phi}_* = \boldsymbol{\phi}(\mathbf{x}_*)$ ,  $\boldsymbol{\Phi} = \boldsymbol{\Phi}(\mathbf{X})$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \boldsymbol{\Phi} \boldsymbol{\Phi}^T + \boldsymbol{\Sigma}_w^{-1})$

# Non-linear Feature Spaces

- Consider the model  $f(\mathbf{x}) = \sum_{i=1}^{D'} w_i \phi_i(\mathbf{x}) = \mathbf{w}^T \boldsymbol{\phi}(\mathbf{x})$ 
  - Each  $\phi_i(\mathbf{x})$  is a (non-linear) feature on  $\mathbf{x}$ , e.g.  $x_1, x_2, x_1^2, x_2^2, x_1 x_2 \dots$
  - We have a non-linear mapping but a **linear-in-the-parameters** model
  - The number of these features can be very large, i.e.  $D' \gg D$
- All the Bayesian analysis is similar to the standard linear model:

$$p(f_* | \mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \mathcal{N}(f_* | \sigma^{-2} \phi_*^T \mathbf{A}^{-1} \boldsymbol{\Phi} \mathbf{y}, \phi_*^T \mathbf{A}^{-1} \phi_*)$$

where:  $\phi_* = \phi(\mathbf{x}_*)$ ,  $\boldsymbol{\Phi} = \boldsymbol{\Phi}(\mathbf{X})$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \boldsymbol{\Phi} \boldsymbol{\Phi}^T + \boldsymbol{\Sigma}_w^{-1})$

- Note we need to invert  $\mathbf{A}$  of ? dimensions.



# Non-linear Feature Spaces

- Consider the model  $f(\mathbf{x}) = \sum_{i=1}^{D'} w_i \phi_i(\mathbf{x}) = \mathbf{w}^T \boldsymbol{\phi}(\mathbf{x})$ 
  - Each  $\phi_i(\mathbf{x})$  is a (non-linear) feature on  $\mathbf{x}$ , e.g.  $x_1, x_2, x_1^2, x_2^2, x_1 x_2 \dots$
  - We have a non-linear mapping but a **linear-in-the-parameters** model
  - The number of these features can be very large, i.e.  $D' \gg D$
- All the Bayesian analysis is similar to the standard linear model:

$$p(f_* | \mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \mathcal{N}(f_* | \sigma^{-2} \boldsymbol{\phi}_*^T \mathbf{A}^{-1} \boldsymbol{\Phi} \mathbf{y}, \boldsymbol{\phi}_*^T \mathbf{A}^{-1} \boldsymbol{\phi}_*)$$

where:  $\boldsymbol{\phi}_* = \boldsymbol{\phi}(\mathbf{x}_*)$ ,  $\boldsymbol{\Phi} = \boldsymbol{\Phi}(\mathbf{X})$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \boldsymbol{\Phi} \boldsymbol{\Phi}^T + \boldsymbol{\Sigma}_w^{-1})$

- Note we need to invert  $\mathbf{A}$  of ? dimensions.

- We can rewrite the predictive distribution as:

$$p(f_* | \mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \mathcal{N}(f_* | \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y}, \mathbf{k}_{**} - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*)$$

where  $\mathbf{k}_* = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_*$ ,  $\mathbf{k}_{**} = \boldsymbol{\phi}_*^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_*$ , and  $\tilde{\mathbf{K}} = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\Phi} + \sigma^2 \mathbf{I}$

# Non-linear Feature Spaces

- Consider the model  $f(\mathbf{x}) = \sum_{i=1}^{D'} w_i \phi_i(\mathbf{x}) = \mathbf{w}^T \boldsymbol{\phi}(\mathbf{x})$ 
  - Each  $\phi_i(\mathbf{x})$  is a (non-linear) feature on  $\mathbf{x}$ , e.g.  $x_1, x_2, x_1^2, x_2^2, x_1 x_2 \dots$
  - We have a non-linear mapping but a **linear-in-the-parameters** model
  - The number of these features can be very large, i.e.  $D' \gg D$
- All the Bayesian analysis is similar to the standard linear model:

$$p(f_* | \mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \mathcal{N}(f_* | \sigma^{-2} \boldsymbol{\phi}_*^T \mathbf{A}^{-1} \boldsymbol{\Phi} \mathbf{y}, \boldsymbol{\phi}_*^T \mathbf{A}^{-1} \boldsymbol{\phi}_*)$$

where:  $\boldsymbol{\phi}_* = \boldsymbol{\phi}(\mathbf{x}_*)$ ,  $\boldsymbol{\Phi} = \boldsymbol{\Phi}(\mathbf{X})$ , and  $\mathbf{A} = (\frac{1}{\sigma^2} \boldsymbol{\Phi} \boldsymbol{\Phi}^T + \boldsymbol{\Sigma}_w^{-1})$

- Note we need to invert  $\mathbf{A}$  of ? dimensions.

- We can rewrite the predictive distribution as:

$$p(f_* | \mathbf{x}_*, \mathbf{X}, \mathbf{y}) = \mathcal{N}(f_* | \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y}, \mathbf{k}_{**} - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*)$$

where  $\mathbf{k}_* = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_*$ ,  $\mathbf{k}_{**} = \boldsymbol{\phi}_*^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_*$ , and  $\tilde{\mathbf{K}} = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\Phi} + \sigma^2 \mathbf{I}$

- Now we need to invert  $\tilde{\mathbf{K}}$  of ? dimensions ▶ GP prediction

# The Kernel Trick

- Note that in:

$$\mathbf{k}_* = \Phi^T \Sigma_w \phi_*, k_{**} = \phi_*^T \Sigma_w \phi_* \text{ and } \tilde{\mathbf{K}} = \Phi^T \Sigma_w \Phi + \sigma^2 \mathbf{I}$$

the features always enter in the form  $\phi(\mathbf{x})^T \Sigma_w \phi(\mathbf{x}')$

# The Kernel Trick

- Note that in:

$$\mathbf{k}_* = \Phi^T \Sigma_w \phi_*, k_{**} = \phi_*^T \Sigma_w \phi_* \text{ and } \tilde{\mathbf{K}} = \Phi^T \Sigma_w \Phi + \sigma^2 \mathbf{I}$$

the features always enter in the form  $\phi(\mathbf{x})^T \Sigma_w \phi(\mathbf{x}')$

- This is an inner product wrt  $\Sigma_w$

# The Kernel Trick

- Note that in:

$$\mathbf{k}_* = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_*, \quad k_{**} = \boldsymbol{\phi}_*^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_* \quad \text{and} \quad \tilde{\mathbf{K}} = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\Phi} + \sigma^2 \mathbf{I}$$

the features always enter in the form  $\boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}(\mathbf{x}')$

- This is an inner product wrt  $\boldsymbol{\Sigma}_w$
- As  $\boldsymbol{\Sigma}_w$  is PD we can rewrite:

$$\begin{aligned} \boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}(\mathbf{x}') &= \boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\phi}(\mathbf{x}') \\ &= \underbrace{(\boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\phi}(\mathbf{x}))^T}_{\boldsymbol{\psi}(\mathbf{x})} \underbrace{(\boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\phi}(\mathbf{x}'))}_{\boldsymbol{\psi}(\mathbf{x}')} \\ \kappa(\mathbf{x}, \mathbf{x}') &= \boldsymbol{\psi}(\mathbf{x}) \cdot \boldsymbol{\psi}(\mathbf{x}') \end{aligned}$$

# The Kernel Trick

- Note that in:

$$\mathbf{k}_* = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_*, \quad k_{**} = \boldsymbol{\phi}_*^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}_* \quad \text{and} \quad \tilde{\mathbf{K}} = \boldsymbol{\Phi}^T \boldsymbol{\Sigma}_w \boldsymbol{\Phi} + \sigma^2 \mathbf{I}$$

the features always enter in the form  $\boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}(\mathbf{x}')$

- This is an inner product wrt  $\boldsymbol{\Sigma}_w$
- As  $\boldsymbol{\Sigma}_w$  is PD we can rewrite:

$$\begin{aligned} \boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\Sigma}_w \boldsymbol{\phi}(\mathbf{x}') &= \boldsymbol{\phi}(\mathbf{x})^T \boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\phi}(\mathbf{x}') \\ &= \underbrace{(\boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\phi}(\mathbf{x}))^T}_{\boldsymbol{\psi}(\mathbf{x})} \underbrace{(\boldsymbol{\Sigma}_w^{1/2} \boldsymbol{\phi}(\mathbf{x}'))}_{\boldsymbol{\psi}(\mathbf{x}')} \\ \kappa(\mathbf{x}, \mathbf{x}') &= \boldsymbol{\psi}(\mathbf{x}) \cdot \boldsymbol{\psi}(\mathbf{x}') \end{aligned}$$

- $\kappa(\cdot, \cdot)$  is called a kernel or covariance function

# The Kernel Trick

- Note that in:

$$\mathbf{k}_* = \Phi^T \Sigma_w \phi_*, \quad k_{**} = \phi_*^T \Sigma_w \phi_* \quad \text{and} \quad \tilde{\mathbf{K}} = \Phi^T \Sigma_w \Phi + \sigma^2 \mathbf{I}$$

the features always enter in the form  $\phi(\mathbf{x})^T \Sigma_w \phi(\mathbf{x}')$

- This is an inner product wrt  $\Sigma_w$
- As  $\Sigma_w$  is PD we can rewrite:

$$\begin{aligned} \phi(\mathbf{x})^T \Sigma_w \phi(\mathbf{x}') &= \phi(\mathbf{x})^T \Sigma_w^{1/2} \Sigma_w^{1/2} \phi(\mathbf{x}') \\ &= \underbrace{(\Sigma_w^{1/2} \phi(\mathbf{x}))^T}_{\psi(\mathbf{x})} \underbrace{(\Sigma_w^{1/2} \phi(\mathbf{x}'))}_{\psi(\mathbf{x}')} \\ \kappa(\mathbf{x}, \mathbf{x}') &= \psi(\mathbf{x}) \cdot \psi(\mathbf{x}') \end{aligned}$$

- $\kappa(\cdot, \cdot)$  is called a kernel or **covariance function**
- We can replace all occurrences of inner products by  $\kappa(\cdot, \cdot)$

# The Kernel Trick

- Note that in:

$$\mathbf{k}_* = \Phi^T \Sigma_w \phi_*, k_{**} = \phi_*^T \Sigma_w \phi_* \text{ and } \tilde{\mathbf{K}} = \Phi^T \Sigma_w \Phi + \sigma^2 \mathbf{I}$$

the features always enter in the form  $\phi(\mathbf{x})^T \Sigma_w \phi(\mathbf{x}')$

- This is an inner product wrt  $\Sigma_w$
- As  $\Sigma_w$  is PD we can rewrite:

$$\begin{aligned}\phi(\mathbf{x})^T \Sigma_w \phi(\mathbf{x}') &= \phi(\mathbf{x})^T \Sigma_w^{1/2} \Sigma_w^{1/2} \phi(\mathbf{x}') \\ &= \underbrace{(\Sigma_w^{1/2} \phi(\mathbf{x}))^T}_{\psi(\mathbf{x})} \underbrace{(\Sigma_w^{1/2} \phi(\mathbf{x}'))}_{\psi(\mathbf{x}')} \\ \kappa(\mathbf{x}, \mathbf{x}') &= \psi(\mathbf{x}) \cdot \psi(\mathbf{x}')\end{aligned}$$

- $\kappa(\cdot, \cdot)$  is called a kernel or covariance function
- We can replace all occurrences of inner products by  $\kappa(\cdot, \cdot)$
- We do not need to compute the feature vectors explicitly



# From a Prior over Weights to a Prior over Functions

- Consider the kinds of functions that can be generated from a set of basis functions with **random weights**.

# From a Prior over Weights to a Prior over Functions

- Consider the kinds of functions that can be generated from a set of basis functions with **random weights**.
- Then  **$f(\mathbf{x})$**  at a particular point is a random variable:

$$f(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x}) \text{ with } \mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

defined as a linear combination of Gaussian random variables.

# From a Prior over Weights to a Prior over Functions

- Consider the kinds of functions that can be generated from a set of basis functions with **random weights**.
- Then  $f(\mathbf{x})$  at a particular point is a random variable:

$$f(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x}) \text{ with } \mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

defined as a linear combination of Gaussian random variables.

- A collection of these random variables indexed by  $\mathbf{x}$ :  
 $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$ , define a **stochastic process** in a **consistent** way.

# From a Prior over Weights to a Prior over Functions

- Consider the kinds of functions that can be generated from a set of basis functions with **random weights**.
- Then  $f(\mathbf{x})$  at a particular point is a random variable:

$$f(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x}) \text{ with } \mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{\Sigma}_w)$$

defined as a linear combination of Gaussian random variables.

- A collection of these random variables indexed by  $\mathbf{x}$ :  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$ , define a **stochastic process** in a **consistent** way.
- The mean and the covariance function for this stochastic process is given by:

$$\begin{aligned}\mathbb{E}_{\mathbf{w}}[f(\mathbf{x})] &= 0 \\ \mathbb{E}_{\mathbf{w}}[f(\mathbf{x})f(\mathbf{x}')] &= \phi^T(\mathbf{x})\mathbf{\Sigma}_w\phi(\mathbf{x}')\end{aligned}$$

# From a Prior over Weights to a Prior over Functions

- Consider the kinds of functions that can be generated from a set of basis functions with **random weights**.
- Then  $f(\mathbf{x})$  at a particular point is a random variable:

$$f(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x}) \text{ with } \mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \Sigma_w)$$

defined as a linear combination of Gaussian random variables.

- A collection of these random variables indexed by  $\mathbf{x}$ :  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$ , define a **stochastic process** in a **consistent** way.
- The mean and the covariance function for this stochastic process is given by:

$$\mathbb{E}_{\mathbf{w}}[f(\mathbf{x})] = 0$$

$$\mathbb{E}_{\mathbf{w}}[f(\mathbf{x})f(\mathbf{x}')] = \phi^T(\mathbf{x})\Sigma_w\phi(\mathbf{x}')$$

- The Bayesian linear model is a **Gaussian process**

# From a Prior over Weights to a Prior over Functions

- Consider the kinds of functions that can be generated from a set of basis functions with **random weights**.
- Then  $f(\mathbf{x})$  at a particular point is a random variable:

$$f(\mathbf{x}) = \mathbf{w}^T \phi(\mathbf{x}) \text{ with } \mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \Sigma_w)$$

defined as a linear combination of Gaussian random variables.

- A collection of these random variables indexed by  $\mathbf{x}$ :  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$ , define a **stochastic process** in a **consistent** way.
- The mean and the covariance function for this stochastic process is given by:

$$\mathbb{E}_{\mathbf{w}}[f(\mathbf{x})] = 0$$

$$\mathbb{E}_{\mathbf{w}}[f(\mathbf{x})f(\mathbf{x}')] = \phi^T(\mathbf{x})\Sigma_w\phi(\mathbf{x}')$$


- The Bayesian linear model is a **Gaussian process**
  - The Function values corresponding to any number of inputs have a **joint Gaussian distribution**.

# Sample Functions from the Linear Model

- 1 Define  $\phi_i(x) = \exp(-\frac{1}{2}(x - \mu_i)^2)$ , for  $i = 1, 2, 3$
- 2 Construct  $\Phi(i, j) = \phi_i(x^{(j)})$ , for  $i = 1, 2, 3$
- 3 Draw  $\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{I})$
- 4 Draw  $\mathbf{f} = \Phi^T \mathbf{w}$

# Sample Functions from the Linear Model



- 1 Define  $\phi_i(x) = \exp(-\frac{1}{2}(x - \mu_i)^2)$ , for  $i = 1, 2, 3$
- 2 Construct  $\Phi(i, j) = \phi_i(x^{(j)})$ , for  $i = 1, 2, 3$
- 3 Draw  $\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{I})$
- 4 Draw  $\mathbf{f} = \Phi^T \mathbf{w}$





# Sample Functions from the Linear Model

- 1 Define  $\phi_i(x) = \exp(-\frac{1}{2}(x - \mu_i)^2)$ , for  $i = 1, 2, 3$
- 2 Construct  $\Phi(i, j) = \phi_i(x^{(j)})$ , for  $i = 1, 2, 3$
- 3 Draw  $\mathbf{w} \sim \mathcal{N}(\mathbf{w}|\mathbf{0}, \mathbf{I})$
- 4 Draw  $\mathbf{f} = \Phi^T \mathbf{w}$



- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

## Gaussian Process (GP)

*$f(\mathbf{x})$  is a Gaussian process if for any finite subset of points  $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}$ , the function values  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$  follow a Gaussian distribution.*

$$f(\mathbf{x}) \sim \mathcal{GP}(\mu(\mathbf{x}), \kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})),$$

$$\mu(\mathbf{x}) = \mathbb{E}[f(\mathbf{x})],$$

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \mathbb{E}[(f(\mathbf{x}) - \mu(\mathbf{x}))(f(\mathbf{x}') - \mu(\mathbf{x}'))],$$

$\mu(\mathbf{x})$ : mean function, consider  $\mu(\mathbf{x}) \equiv \mathbf{0}$

$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})$ : parameterized covariance function, notion of similarity

## Gaussian Process (GP)

$f(\mathbf{x})$  is a Gaussian process if for any finite subset of points  $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}$ , the function values  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$  follow a Gaussian distribution.

$$f(\mathbf{x}) \sim \mathcal{GP}(\mu(\mathbf{x}), \kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})),$$

$$\mu(\mathbf{x}) = \mathbb{E}[f(\mathbf{x})],$$

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \mathbb{E}[(f(\mathbf{x}) - \mu(\mathbf{x}))(f(\mathbf{x}') - \mu(\mathbf{x}'))],$$

$\mu(\mathbf{x})$ : mean function, consider  $\mu(\mathbf{x}) \equiv \mathbf{0}$

$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})$ : parameterized covariance function, notion of similarity

- Stochastic process: collection of random variables

## Gaussian Process (GP)

$f(\mathbf{x})$  is a Gaussian process if for any finite subset of points  $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}$ , the function values  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$  follow a Gaussian distribution.

$$f(\mathbf{x}) \sim \mathcal{GP}(\mu(\mathbf{x}), \kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})),$$

$$\mu(\mathbf{x}) = \mathbb{E}[f(\mathbf{x})],$$

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \mathbb{E}[(f(\mathbf{x}) - \mu(\mathbf{x}))(f(\mathbf{x}') - \mu(\mathbf{x}'))],$$

$\mu(\mathbf{x})$ : mean function, consider  $\mu(\mathbf{x}) \equiv \mathbf{0}$

$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})$ : parameterized covariance function, notion of similarity

- Stochastic process: collection of random variables
- These variables are the values of the function  $f(\mathbf{x})$  indexed by the set of all possible input

## Gaussian Process (GP)

$f(\mathbf{x})$  is a Gaussian process if for any finite subset of points  $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(N)}$ , the function values  $f(\mathbf{x}^{(1)}), \dots, f(\mathbf{x}^{(N)})$  follow a Gaussian distribution.

$$f(\mathbf{x}) \sim \mathcal{GP}(\mu(\mathbf{x}), \kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})),$$

$$\mu(\mathbf{x}) = \mathbb{E}[f(\mathbf{x})],$$

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \mathbb{E}[(f(\mathbf{x}) - \mu(\mathbf{x}))(f(\mathbf{x}') - \mu(\mathbf{x}'))],$$

$\mu(\mathbf{x})$ : **mean function**, consider  $\mu(\mathbf{x}) \equiv \mathbf{0}$

$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta})$ : parameterized **covariance function**, notion of similarity

- **Stochastic process**: collection of random variables
- These variables are the values of the function  $f(\mathbf{x})$  **indexed** by the set of all possible input
- **Consistency**: marginalization property  
 $(f_1, f_2) \sim \mathcal{N}(\mathbf{f}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) \rightarrow f_1 \sim \mathcal{N}(f_1|\mu_1, \Sigma_{11})$

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\mathbb{C}\text{ov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\mathbb{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs



# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\mathbb{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
- ▶ A crucial component in GPs

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\mathbb{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
- ▶ A crucial component in GPs
- ▶ Intuitively, it describes the notion of **similarity**

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\mathbb{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
- ▶ A crucial component in GPs
- ▶ Intuitively, it describes the notion of **similarity**
- ▶ It can be parametrized and we can learn its **hyperparameters** from data

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\mathbb{C}\text{ov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
  - ▶ A crucial component in GPs
  - ▶ Intuitively, it describes the notion of **similarity**
  - ▶ It can be parametrized and we can learn its **hyperparameters** from data
- The matrix **K** such that  $K_{i,j} = \kappa(\mathbf{x}^{(i)}, \mathbf{x}^{(j)})$  at all pairwise input points is known as the covariance matrix or **Gram matrix**.

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\text{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
  - ▶ A crucial component in GPs
  - ▶ Intuitively, it describes the notion of **similarity**
  - ▶ It can be parametrized and we can learn its **hyperparameters** from data
- The matrix  $\mathbf{K}$  such that  $K_{i,j} = \kappa(\mathbf{x}^{(i)}, \mathbf{x}^{(j)})$  at all pairwise input points is known as the covariance matrix or **Gram matrix**.
- it must generate a positive semidefinite (PSD) matrix at any subset of points, i.e.  $\mathbf{b}^T \mathbf{K} \mathbf{b} \geq 0, \forall \mathbf{b} \in \mathbb{R}^N$

# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\text{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
  - ▶ A crucial component in GPs
  - ▶ Intuitively, it describes the notion of **similarity**
  - ▶ It can be parametrized and we can learn its **hyperparameters** from data
- The matrix  $\mathbf{K}$  such that  $K_{i,j} = \kappa(\mathbf{x}^{(i)}, \mathbf{x}^{(j)})$  at all pairwise input points is known as the covariance matrix or **Gram matrix**.
- it must generate a positive semidefinite (PSD) matrix at any subset of points, i.e.  $\mathbf{b}^T \mathbf{K} \mathbf{b} \geq 0, \forall \mathbf{b} \in \mathbb{R}^N$
- **Stationary**:  $\varphi(\mathbf{x} - \mathbf{x}')$  - translation invariant





# The Covariance Function

- It specifies the covariance between pairs of random variables:

$$\text{Cov}(f(\mathbf{x}^{(p)}), f(\mathbf{x}^{(q)})) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)})$$

- ▶ Covariance between outputs as a function of the inputs
  - ▶ A crucial component in GPs
  - ▶ Intuitively, it describes the notion of **similarity**
  - ▶ It can be parametrized and we can learn its **hyperparameters** from data
- The matrix  $\mathbf{K}$  such that  $K_{i,j} = \kappa(\mathbf{x}^{(i)}, \mathbf{x}^{(j)})$  at all pairwise input points is known as the covariance matrix or **Gram matrix**.
- it must generate a positive semidefinite (PSD) matrix at any subset of points, i.e.  $\mathbf{b}^T \mathbf{K} \mathbf{b} \geq 0, \forall \mathbf{b} \in \mathbb{R}^N$
- **Stationary**:  $\varphi(\mathbf{x} - \mathbf{x}')$  - translation invariant
- **Isotropic**:  $\varphi(\|\mathbf{x} - \mathbf{x}'\|)$


# Samples from a Gaussian Process






# Computing with Infinite Vectors


GP prior



GP regression example



Inference result



$$K_{\infty} =$$

A diagram representing the kernel matrix  $K_{\infty}$ . It consists of a grid of dashed red lines, forming a cross-like pattern that extends infinitely in all directions, representing the structure of the kernel matrix for an infinite-dimensional space.

$$K_{\infty} =$$

A diagram representing the kernel matrix  $K_{\infty}$ . It consists of a 4x4 grid of orange squares, representing the structure of the kernel matrix for a finite-dimensional space.

$$K_y =$$

A diagram representing the kernel matrix  $K_y$ . It consists of a 4x4 grid of orange squares, representing the structure of the kernel matrix for a finite-dimensional space.

# The Squared Exponential (SE) Covariance Function

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \sigma_s^2 \exp \left( -\frac{1}{2} (\mathbf{x} - \mathbf{x}')^T \mathbf{C} (\mathbf{x} - \mathbf{x}') \right)$$

- $\sigma_s^2$  is the signal variance

# The Squared Exponential (SE) Covariance Function

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \sigma_s^2 \exp \left( -\frac{1}{2} (\mathbf{x} - \mathbf{x}')^T \mathbf{C} (\mathbf{x} - \mathbf{x}') \right)$$

- $\sigma_s^2$  is the signal variance
- $\mathbf{C}$  is a symmetric matrix that can have different parameterizations

# The Squared Exponential (SE) Covariance Function

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \sigma_s^2 \exp \left( -\frac{1}{2} (\mathbf{x} - \mathbf{x}')^T \mathbf{C} (\mathbf{x} - \mathbf{x}') \right)$$

- $\sigma_s^2$  is the signal variance
- $\mathbf{C}$  is a symmetric matrix that can have different parameterizations
- $\mathbf{C} = \ell^{-2} \mathbf{I}$ : isotropic SE

# The Squared Exponential (SE) Covariance Function

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \sigma_s^2 \exp \left( -\frac{1}{2} (\mathbf{x} - \mathbf{x}')^T \mathbf{C} (\mathbf{x} - \mathbf{x}') \right)$$

- $\sigma_s^2$  is the signal variance
- $\mathbf{C}$  is a symmetric matrix that can have different parameterizations
- $\mathbf{C} = \ell^{-2} \mathbf{I}$ : isotropic SE
- $\mathbf{C} = \text{diag}(\ell)^{-2}$  with  $\ell = (\ell_1, \dots, \ell_D)$ : Automatic Relevance Determination (ARD)


# The Squared Exponential (SE) Covariance Function

$$\kappa(\mathbf{x}, \mathbf{x}'; \boldsymbol{\theta}) = \sigma_s^2 \exp \left( -\frac{1}{2} (\mathbf{x} - \mathbf{x}')^T \mathbf{C} (\mathbf{x} - \mathbf{x}') \right)$$

- $\sigma_s^2$  is the signal variance
- $\mathbf{C}$  is a symmetric matrix that can have different parameterizations
- $\mathbf{C} = \ell^{-2} \mathbf{I}$ : isotropic SE
- $\mathbf{C} = \text{diag}(\ell)^{-2}$  with  $\ell = (\ell_1, \dots, \ell_D)$ : Automatic Relevance Determination (ARD)
- Each  $\ell_j$  is known as the characteristic length-scale: distance for which the function values are expected to vary significantly

# The Squared Exponential (SE) Covariance Function


## Example




$$\ell = 1, \sigma_s^2 = 1$$

# The Squared Exponential (SE) Covariance Function

## Example



$$\ell = 1, \sigma_s^2 = 1$$




$$\ell = 0.1, \sigma_s^2 = 1$$




# The Squared Exponential (SE) Covariance Function


## Example



$$\ell = 1, \sigma_s^2 = 1$$




$$\ell = 0.1, \sigma_s^2 = 1$$




$$\ell = 1, \sigma_s^2 = 4$$

# The Squared Exponential (SE) Covariance Function


## Example




$$\ell = 1, \sigma_s^2 = 1$$



$$\ell = 0.1, \sigma_s^2 = 1$$



$$\ell = 1, \sigma_s^2 = 4$$



$$\ell = 0.1, \sigma_s^2 = 4$$

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

**Prior**  $f(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \kappa(\mathbf{x}, \mathbf{x}'))$

# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

**Prior**  $f(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \kappa(\mathbf{x}, \mathbf{x}'))$

**Noise**  $y = f(\mathbf{x}) + \eta \quad \eta \sim \mathcal{N}(0, \sigma_n^2)$

# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

**Prior**  $f(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \kappa(\mathbf{x}, \mathbf{x}'))$

**Noise**  $y = f(\mathbf{x}) + \eta \quad \eta \sim \mathcal{N}(0, \sigma_n^2)$

- The joint distribution of  $\mathbf{y}$  and  $f_*$  is a Gaussian

# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

**Prior**  $f(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \kappa(\mathbf{x}, \mathbf{x}'))$

**Noise**  $y = f(\mathbf{x}) + \eta \quad \eta \sim \mathcal{N}(0, \sigma_n^2)$

- The joint distribution of  $\mathbf{y}$  and  $f_*$  is a Gaussian
- We simply need to figure out the **covariance structure**:  
 $\text{Cov}(y^{(p)}, y^{(q)}) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)}) + \sigma_n^2 \delta_{pq} \rightarrow \text{Cov}(\mathbf{y}) = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$



# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N, \mathbf{x} \in \mathbb{R}^D, y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

**Prior**  $f(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \kappa(\mathbf{x}, \mathbf{x}'))$

**Noise**  $y = f(\mathbf{x}) + \eta \quad \eta \sim \mathcal{N}(0, \sigma_n^2)$

- The joint distribution of  $\mathbf{y}$  and  $f_*$  is a Gaussian
- We simply need to figure out the **covariance structure**:  
 $\text{Cov}(y^{(p)}, y^{(q)}) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)}) + \sigma_n^2 \delta_{pq} \rightarrow \text{Cov}(\mathbf{y}) = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$
- To get the posterior on  $f_*$  we need to constrain this distribution to agree with the observed data  $(\mathbf{X}, \mathbf{y})$

# Standard GP Regression Model: Predictions (1)

**Data** :  $\mathcal{D} = \{(\mathbf{x}^{(n)}, y^{(n)})\}_{n=1}^N$ ,  $\mathbf{x} \in \mathbb{R}^D$ ,  $y \in \mathbb{R}$

**Input** :  $(\mathbf{X})_{D \times N}$ , **Targets**:  $(\mathbf{y})_{N \times 1}$

**Goal** : Make predictions  $f_* = f(\mathbf{x}_*)$  at  $\mathbf{x}_*$

**Prior**  $f(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \kappa(\mathbf{x}, \mathbf{x}'))$

**Noise**  $y = f(\mathbf{x}) + \eta \quad \eta \sim \mathcal{N}(0, \sigma_n^2)$

- The joint distribution of  $\mathbf{y}$  and  $f_*$  is a Gaussian
- We simply need to figure out the **covariance structure**:  
 $\text{Cov}(y^{(p)}, y^{(q)}) = \kappa(\mathbf{x}^{(p)}, \mathbf{x}^{(q)}) + \sigma_n^2 \delta_{pq} \rightarrow \text{Cov}(\mathbf{y}) = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$
- To get the posterior on  $f_*$  we need to constrain this distribution to agree with the observed data  $(\mathbf{X}, \mathbf{y})$
- This is achieved simply by **conditioning**:  $p(f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_*)$

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & k(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_* \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{K}(\mathbf{X}, \mathbf{x}_*)$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{k}(\mathbf{x}_*, \mathbf{X})$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$  then:

$$f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_* \sim \mathcal{N}(\mathbb{E}[f_*], \mathbb{V}[f_*]),$$

$$\mathbb{E}[f_*] = \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y},$$

$$\mathbb{V}[f_*] = \kappa(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*.$$

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{K}(\mathbf{X}, \mathbf{x}_*)$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$  then:

$$f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_* \sim \mathcal{N}(\mathbb{E}[f_*], \mathbb{V}[f_*]),$$

$$\mathbb{E}[f_*] = \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y},$$

$$\mathbb{V}[f_*] = \kappa(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*.$$

- $\mathbb{E}[f_*]$ : Linear combination of  $N$  observations, i.e. **linear predictor**

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{K}(\mathbf{X}, \mathbf{x}_*)$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$  then:

$$f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_* \sim \mathcal{N}(\mathbb{E}[f_*], \mathbb{V}[f_*]),$$

$$\mathbb{E}[f_*] = \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y},$$

$$\mathbb{V}[f_*] = \kappa(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*.$$

- $\mathbb{E}[f_*]$ : Linear combination of  $N$  observations, i.e. **linear predictor**
- Say  $\boldsymbol{\alpha} = (\mathbf{K} + \sigma_n^2 \mathbf{I})^{-1} \mathbf{y}$  then:  $\mathbb{E}[f_*] = \sum_{n=1}^N \alpha_i \kappa(\mathbf{x}^{(n)}, \mathbf{x}_*)$  is a linear combination of  $N$  kernel functions: **Representer theorem**

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{K}(\mathbf{X}, \mathbf{x}_*)$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$  then:

$$f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_* \sim \mathcal{N}(\mathbb{E}[f_*], \mathbb{V}[f_*]),$$

$$\mathbb{E}[f_*] = \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y},$$

$$\mathbb{V}[f_*] = \kappa(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*.$$

- $\mathbb{E}[f_*]$ : Linear combination of  $N$  observations, i.e. **linear predictor**
- Say  $\boldsymbol{\alpha} = (\mathbf{K} + \sigma_n^2 \mathbf{I})^{-1} \mathbf{y}$  then:  $\mathbb{E}[f_*] = \sum_{n=1}^N \alpha_i \kappa(\mathbf{x}^{(n)}, \mathbf{x}_*)$  is a linear combination of  $N$  kernel functions: **Representer theorem**
- We encountered this predictive distribution before [▶ Go to Linear Model](#)



## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{K}(\mathbf{X}, \mathbf{x}_*)$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$  then:

$$f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_* \sim \mathcal{N}(\mathbb{E}[f_*], \mathbb{V}[f_*]),$$

$$\mathbb{E}[f_*] = \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y},$$

$$\mathbb{V}[f_*] = \kappa(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*.$$

- $\mathbb{E}[f_*]$ : Linear combination of  $N$  observations, i.e. **linear predictor**
- Say  $\boldsymbol{\alpha} = (\mathbf{K} + \sigma_n^2 \mathbf{I})^{-1} \mathbf{y}$  then:  $\mathbb{E}[f_*] = \sum_{n=1}^N \alpha_i \kappa(\mathbf{x}^{(n)}, \mathbf{x}_*)$  is a linear combination of  $N$  kernel functions: **Representer theorem**
- We encountered this predictive distribution before [▶ Go to Linear Model](#)
- $\mathbb{V}[f_*]$  does not depend on  $\mathbf{y}$

## Standard GP Regression Model: Predictions (2)

$$\begin{bmatrix} \mathbf{y} \\ f_* \end{bmatrix} \sim \mathcal{N} \left( \mathbf{0}, \begin{bmatrix} \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I} & \mathbf{k}(\mathbf{X}, \mathbf{x}_*) \\ \mathbf{k}(\mathbf{x}_*, \mathbf{X}) & \kappa(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

Denoting  $\mathbf{k}_* = \mathbf{K}(\mathbf{X}, \mathbf{x}_*)$  and  $\tilde{\mathbf{K}} = \mathbf{K}(\mathbf{X}, \mathbf{X}) + \sigma_n^2 \mathbf{I}$  then:

$$f_* | \mathbf{X}, \mathbf{y}, \mathbf{x}_* \sim \mathcal{N}(\mathbb{E}[f_*], \mathbb{V}[f_*]),$$

$$\mathbb{E}[f_*] = \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{y},$$

$$\mathbb{V}[f_*] = \kappa(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}_*^T \tilde{\mathbf{K}}^{-1} \mathbf{k}_*.$$

- $\mathbb{E}[f_*]$ : Linear combination of  $N$  observations, i.e. **linear predictor**
- Say  $\boldsymbol{\alpha} = (\mathbf{K} + \sigma_n^2 \mathbf{I})^{-1} \mathbf{y}$  then:  $\mathbb{E}[f_*] = \sum_{n=1}^N \alpha_i \kappa(\mathbf{x}^{(n)}, \mathbf{x}_*)$  is a linear combination of  $N$  kernel functions: **Representer theorem**
- We encountered this predictive distribution before [▶ Go to Linear Model](#)
- $\mathbb{V}[f_*]$  does not depend on  $\mathbf{y}$
- In fact we have a **Gaussian posterior process**

# The Graphical Model for GPs

# The Graphical Model for GPs




Figure from Carl Rasmussen's slides

# The Graphical Model for GPs




Figure from Carl Rasmussen's slides

- Observations  $y$  depend on their corresponding latent function  $f$

# The Graphical Model for GPs




Figure from Carl Rasmussen's slides

- Observations  $y$  depend on their corresponding latent function  $f$
- The **marginalization** property implies that adding a new  $x_i^*, f_i^*, y_i^*$  does not affect the distribution

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - **Model Selection**
  - Other Covariance Functions

# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the [hyper-parameters](#).



# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the **hyper-parameters**.
- E.g. for the SE:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma_s^2 \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{x}')^T \mathbf{C}(\mathbf{x} - \mathbf{x}')^T\right)$  the parameters are  $\sigma_s^2$  and the parameters of  $\mathbf{C}$

# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the **hyper-parameters**.
- E.g. for the SE:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma_s^2 \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{x}')^T \mathbf{C}(\mathbf{x} - \mathbf{x}')^T\right)$  the parameters are  $\sigma_s^2$  and the parameters of  $\mathbf{C}$
- However, we will refer to the set of hyper-parameters  $\theta$  as the parameters of the covariance and the noise variance  $\sigma_n^2$

# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the **hyper-parameters**.
- E.g. for the SE:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma_s^2 \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{x}')^T \mathbf{C}(\mathbf{x} - \mathbf{x}')^T\right)$  the parameters are  $\sigma_s^2$  and the parameters of  $\mathbf{C}$
- However, we will refer to the set of hyper-parameters  $\theta$  as the parameters of the covariance and the noise variance  $\sigma_n^2$
- We can do cross-validation (**potential problems?**)

# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the **hyper-parameters**.
- E.g. for the SE:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma_s^2 \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{x}')^T \mathbf{C}(\mathbf{x} - \mathbf{x}')^T\right)$  the parameters are  $\sigma_s^2$  and the parameters of  $\mathbf{C}$
- However, we will refer to the set of hyper-parameters  $\theta$  as the parameters of the covariance and the noise variance  $\sigma_n^2$
- We can do cross-validation (**potential problems?**)
- We focus here on the so-called type II maximum likelihood, i.e. we want to maximize the **marginal likelihood**.

# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the **hyper-parameters**.
- E.g. for the SE:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma_s^2 \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{x}')^T \mathbf{C}(\mathbf{x} - \mathbf{x}')^T\right)$  the parameters are  $\sigma_s^2$  and the parameters of  $\mathbf{C}$
- However, we will refer to the set of hyper-parameters  $\theta$  as the parameters of the covariance and the noise variance  $\sigma_n^2$
- We can do cross-validation (**potential problems?**)
- We focus here on the so-called type II maximum likelihood, i.e. we want to maximize the **marginal likelihood**.
- Integrate out the “parameters” of the GP: (**which parameters?**)

# Model Selection

- It includes the discrete choice of the functional form for the covariance function and the values for the **hyper-parameters**.
- E.g. for the SE:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma_s^2 \exp\left(-\frac{1}{2}(\mathbf{x} - \mathbf{x}')^T \mathbf{C}(\mathbf{x} - \mathbf{x}')^T\right)$  the parameters are  $\sigma_s^2$  and the parameters of  $\mathbf{C}$
- However, we will refer to the set of hyper-parameters  $\theta$  as the parameters of the covariance and the noise variance  $\sigma_n^2$
- We can do cross-validation (**potential problems?**)
- We focus here on the so-called type II maximum likelihood, i.e. we want to maximize the **marginal likelihood**.
- Integrate out the “parameters” of the GP: (**which parameters?**)

$$\begin{aligned} p(\mathbf{y}|\mathbf{X}, \theta) &= \int p(\mathbf{y}|\mathbf{f}, \mathbf{X}, \theta) p(\mathbf{f}|\mathbf{X}, \theta) d\mathbf{f} \\ &= \mathcal{N}(\mathbf{y}|\mathbf{0}, \mathbf{K} + \sigma^2 \mathbf{I}) \end{aligned}$$

# Log Marginal Likelihood

$$\begin{aligned}\mathcal{L} &= \log p(\mathbf{y}|\mathbf{X}, \boldsymbol{\theta}) \\ &= \underbrace{-\frac{1}{2}\mathbf{y}^T(\mathbf{K} + \sigma_n^2\mathbf{I})^{-1}\mathbf{y}}_{\text{data-fit}} - \underbrace{\frac{1}{2}\log|\mathbf{K} + \sigma_n^2\mathbf{I}|}_{\text{complexity}} - \underbrace{\frac{N}{2}\log 2\pi}_{\text{normaliz.}}\end{aligned}$$

# Log Marginal Likelihood

$$\begin{aligned}\mathcal{L} &= \log p(\mathbf{y}|\mathbf{X}, \boldsymbol{\theta}) \\ &= \underbrace{-\frac{1}{2}\mathbf{y}^T(\mathbf{K} + \sigma_n^2\mathbf{I})^{-1}\mathbf{y}}_{\text{data-fit}} - \underbrace{\frac{1}{2}\log|\mathbf{K} + \sigma_n^2\mathbf{I}|}_{\text{complexity}} - \underbrace{\frac{N}{2}\log 2\pi}_{\text{normaliz.}}\end{aligned}$$


- Isotropic SE
- $\sigma_s^2 = 1$ ,  $\sigma_n^2 = 0.01$
- $\ell = 1$
- $N = 20$



# Log Marginal Likelihood

$$\begin{aligned}\mathcal{L} &= \log p(\mathbf{y}|\mathbf{X}, \boldsymbol{\theta}) \\ &= \underbrace{-\frac{1}{2}\mathbf{y}^T(\mathbf{K} + \sigma_n^2\mathbf{I})^{-1}\mathbf{y}}_{\text{data-fit}} - \underbrace{\frac{1}{2}\log|\mathbf{K} + \sigma_n^2\mathbf{I}|}_{\text{complexity}} - \underbrace{\frac{N}{2}\log 2\pi}_{\text{normaliz.}}\end{aligned}$$

- Isotropic SE
- $\sigma_s^2 = 1$ ,  $\sigma_n^2 = 0.01$
- $\ell = 1$
- $N = 20$



# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

- Can use gradient-based optimization

# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

- Can use gradient-based optimization
- General approach and only needs derivatives of the covariance

# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

- Can use gradient-based optimization
- General approach and only needs derivatives of the covariance
- Such principled “kernel” learning does not exist in standard SVM

# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

- Can use gradient-based optimization
- General approach and only needs derivatives of the covariance
- Such principled “kernel” learning does not exist in standard SVM
- **Non-convex optimization**

# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

- Can use gradient-based optimization
- General approach and only needs derivatives of the covariance
- Such principled “kernel” learning does not exist in standard SVM
- **Non-convex optimization**
- Multiple local optima correspond to different explanations of the data

# Hyper-parameter Learning

Let  $\tilde{\mathbf{K}} = \mathbf{K} + \sigma_n^2 \mathbf{I}$ :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta_i} &= \frac{1}{2} \mathbf{y}^T \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \tilde{\mathbf{K}}^{-1} \mathbf{y} - \frac{1}{2} \text{tr} \left( \tilde{\mathbf{K}}^{-1} \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right) \\ &= \frac{1}{2} \text{tr} \left( (\boldsymbol{\alpha} \boldsymbol{\alpha}^T - \tilde{\mathbf{K}}^{-1}) \frac{\partial \tilde{\mathbf{K}}}{\partial \theta_i} \right)\end{aligned}$$

where  $\boldsymbol{\alpha} = \tilde{\mathbf{K}}^{-1} \mathbf{y}$ .

- Can use gradient-based optimization
- General approach and only needs derivatives of the covariance
- Such principled “kernel” learning does not exist in standard SVM
- **Non-convex optimization**
- Multiple local optima correspond to different explanations of the data
- **Computational Requirements?**



# Automatic Relevance Determination (ARD)


- Inverse of the length-scale determines the relevance of the dimension.

# Automatic Relevance Determination (ARD)

- Inverse of the length-scale determines the relevance of the dimension.
- The larger the length-scale the more irrelevant the corresponding input is.


# Automatic Relevance Determination (ARD)

- Inverse of the length-scale determines the relevance of the dimension.
- The larger the length-scale the more irrelevant the corresponding input is.



# Automatic Relevance Determination (ARD)

- Inverse of the length-scale determines the relevance of the dimension.
- The larger the length-scale the more irrelevant the corresponding input is.



Learned length-scale for irrelevant dimension:  $1.0557 \times 10^5$

- 1 The Gaussian Distribution Revisited
- 2 Standard Bayesian Linear Regression
- 3 Bayesian Regression with Non-linear Feature Spaces
- 4 Gaussian Processes for Regression
  - Function-space View
  - Predictions
  - Model Selection
  - Other Covariance Functions

## Other Covariance Functions: Matérn Covariance


$$\kappa(\mathbf{x}, \mathbf{x}') = \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)^\nu \mathcal{K}_\nu \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)$$

where  $\mathcal{K}_\nu$  is a modified Bessel function and  $\nu > 0$ ,  $\ell > 0$ .

# Other Covariance Functions: Matérn Covariance

$$\kappa(\mathbf{x}, \mathbf{x}') = \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)^\nu \mathcal{K}_\nu \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)$$

where  $\mathcal{K}_\nu$  is a modified Bessel function and  $\nu > 0$ ,  $\ell > 0$ .




$\ell = 1$

# Other Covariance Functions: Matérn Covariance

$$\kappa(\mathbf{x}, \mathbf{x}') = \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)^\nu \mathcal{K}_\nu \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)$$

where  $\mathcal{K}_\nu$  is a modified Bessel function and  $\nu > 0$ ,  $\ell > 0$ .



- Stationary, Isotropic


$\ell = 1$



# Other Covariance Functions: Matérn Covariance

$$\kappa(\mathbf{x}, \mathbf{x}') = \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)^\nu \mathcal{K}_\nu \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)$$

where  $\mathcal{K}_\nu$  is a modified Bessel function and  $\nu > 0$ ,  $\ell > 0$ .



$\ell = 1$

- Stationary, Isotropic

- $\nu = 1/2$ :


$$\kappa(\mathbf{x}, \mathbf{x}') = \exp\left(-\frac{\|\mathbf{x} - \mathbf{x}'\|}{\ell}\right)$$

- ▶ Very rough process
- ▶ Brownian motion
- ▶ Ornstein-Uhlenbeck (D=1)

# Other Covariance Functions: Matérn Covariance

$$\kappa(\mathbf{x}, \mathbf{x}') = \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)^\nu \mathcal{K}_\nu \left( \frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{x}'\|}{\ell} \right)$$

where  $\mathcal{K}_\nu$  is a modified Bessel function and  $\nu > 0$ ,  $\ell > 0$ .



$\ell = 1$

- Stationary, Isotropic
- $\nu = 1/2$ :
  - $\kappa(\mathbf{x}, \mathbf{x}') = \exp\left(-\frac{\|\mathbf{x} - \mathbf{x}'\|}{\ell}\right)$ 
    - ▶ Very rough process
    - ▶ Brownian motion
    - ▶ Ornstein-Uhlenbeck (D=1)
- $\nu \rightarrow \infty$ : SE covariance

## Other Covariance Functions: Rational Quadratic

$$\kappa(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2\alpha\ell^2}\right)^{-\alpha}$$

with  $\alpha > 0$ ,  $\ell > 0$ .


can be seen as an infinite sum of squared exponential (SE) covariance functions with different characteristic length-scales.

# Other Covariance Functions: Rational Quadratic

$$\kappa(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2\alpha\ell^2}\right)^{-\alpha}$$

with  $\alpha > 0$ ,  $\ell > 0$ .

can be seen as **an infinite sum of squared exponential** (SE) covariance functions with different characteristic length-scales.




# Other Covariance Functions: Rational Quadratic

$$\kappa(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2\alpha\ell^2}\right)^{-\alpha}$$

with  $\alpha > 0$ ,  $\ell > 0$ .

can be seen as **an infinite sum of squared exponential** (SE) covariance functions with different characteristic length-scales.




with  $\alpha \rightarrow \infty$  is the SE covariance with length-scale  $\ell$ .

# Other Covariance Functions: Neural Network Covariance

- Consider a neural network with **one hidden layer** and  $N_H$  hidden units.
- Under certain assumptions the corresponding stochastic process will converge to a Gaussian Process as  $N_H \rightarrow \infty$ .
- For a specific settings of the transfer function of the neural net:

$$\kappa(\mathbf{x}, \mathbf{x}') = \frac{2}{\pi} \sin^{-1} \left( \frac{2\tilde{\mathbf{x}}^T \mathbf{\Sigma} \tilde{\mathbf{x}'}}{\sqrt{(1 + 2\tilde{\mathbf{x}}^T \mathbf{\Sigma} \tilde{\mathbf{x}})(1 + 2\tilde{\mathbf{x}'}^T \mathbf{\Sigma} \tilde{\mathbf{x}'})}} \right)$$



# Other Covariance Functions: Periodic, Smooth Functions


We can create a distribution over **periodic functions** of  $x$  by using the mapping  $\mathbf{u}(x) = (\cos(x), \sin(x))$  and then use the SE covariance on  $\mathbf{u}$  space. This gives rise to:

$$\kappa(x, x') = \exp\left(-\frac{2 \sin^2\left(\frac{x-x'}{2}\right)}{\ell^2}\right)$$

# Other Covariance Functions: Periodic, Smooth Functions

We can create a distribution over **periodic functions** of  $x$  by using the mapping  $\mathbf{u}(x) = (\cos(x), \sin(x))$  and then use the SE covariance on  $\mathbf{u}$  space. This gives rise to:

$$\kappa(x, x') = \exp\left(-\frac{2\sin^2\left(\frac{x-x'}{2}\right)}{\ell^2}\right)$$






# Other Covariance Functions: Periodic, Smooth Functions

We can create a distribution over **periodic functions** of  $x$  by using the mapping  $\mathbf{u}(x) = (\cos(x), \sin(x))$  and then use the SE covariance on  $\mathbf{u}$  space. This gives rise to:

$$\kappa(x, x') = \exp\left(-\frac{2\sin^2\left(\frac{x-x'}{2}\right)}{\ell^2}\right)$$



This is called **warping** and can also be used to introduce non-stationarity.

# Conclusions

- Models based on Gaussian process (GP) priors are flexible non-parametric Bayesian approaches to non-linear regression.
- GP regression can be seen as a generalisation of Bayesian regression with non-linear feature spaces and infinite-dimensional feature maps.
- Inference in GP regression models is analytically tractable and is computable thanks to the marginalisation property underlying GPs.
- Hyper-parameter learning carried out via non-convex optimisation of the marginal likelihood.
- High computational cost in time and memory,  $\mathcal{O}(N^3)$  and  $\mathcal{O}(N^2)$
- Reading: Rasmussen & Williams (GPML, 2006): Ch. 1, 2, 4 (except Sec. 4.3, 4.4), Sec. 5.1, 5.2, 5.4.