

Problem Set #2

Quiz, 5 questions

4/5 points (80%)

**Congratulations! You passed!**

Next Item

1 / 1
point

1.

This question will give you further practice with the Master Method. Suppose the running time of an algorithm is governed by the recurrence $T(n) = 7 * T(n/3) + n^2$. What's the overall asymptotic running time (i.e., the value of $T(n)$)?

- ☐ $\theta(n \log n)$
- ☐ $\theta(n^{2.81})$
- ☐ $\theta(n^2 \log n)$
- ☒ $\theta(n^2)$

Correct

$a=7, b=3, d=2$. Since $b^d > a$, this is case 2 of the Master Method.

1 / 1
point

2.

This question will give you further practice with the Master Method. Suppose the running time of an algorithm is governed by the recurrence $T(n) = 9 * T(n/3) + n^2$. What's the overall asymptotic running time (i.e., the value of $T(n)$)?

- ☐ $\theta(n^2)$
- ☐ $\theta(n \log n)$
- ☐ $\theta(n^{3.17})$
- ☒ $\theta(n^2 \log n)$

Correct

$a = b^d = 9$, so this is case 1 of the Master Method.

Problem Set #2

4/5 points (80%)

Quiz, 5 questions

1 / 1
point

3.

This question will give you further practice with the Master Method. Suppose the running time of an algorithm is governed by the recurrence $T(n) = 5 * T(n/3) + 4n$. What's the overall asymptotic running time (i.e., the value of $T(n)$)?

☐ $\theta(n^{2.59})$ ☐ $\theta(n^{5/3})$ ☐ $\theta(n^2)$ ☐ $\theta(n \log(n))$ ☐ $\theta(n^{\frac{\log 3}{\log 5}})$ ☒ $\theta(n^{\log_3(5)})$ **Correct**

$a = 5$, $b = 3$, $d = 1$. Since $a > b^d$, this is case 3 of the Master Method.

0 / 1
point

4.

Consider the following pseudocode for calculating a^b (where a and b are positive integers)

```

1  FastPower(a,b) :
2    if b = 1
3      return a
4    else
5      c := a*a
6      ans := FastPower(c, [b/2])
7      if b is odd
8        return a*ans
9      else return ans
10 end

```

Here $[x]$ denotes the floor function, that is, the largest integer less than or equal to x .

Now assuming that you use a calculator that supports multiplication and division (i.e., you can do multiplications and divisions in constant time), what would be the overall asymptotic running time of the above algorithm (as a function of b)?

Problem Set #2

Quiz, 5 questions

4/5 points (80%)

- ☐ $\Theta(\sqrt{b})$
- ☐ $\Theta(\log(b))$
- ☐ $\Theta(b)$
- ☒ $\Theta(b \log(b))$

This should not be selected

This would be even worse than the naive method of multiplying a by itself b times...



1 / 1
point

5.

Choose the smallest correct upper bound on the solution to the following recurrence: $T(1) = 1$ and $T(n) \leq T(\lfloor \sqrt{n} \rfloor) + 1$ for $n > 1$. Here $\lfloor x \rfloor$ denotes the "floor" function, which rounds down to the nearest integer. (Note that the Master Method does not apply.)

- ☐ $O(\sqrt{n})$
- ☐ $O(\log n)$
- ☐ $O(1)$
- ☒ $O(\log \log n)$

Correct

Bingo! This answer may be easiest to see by writing n as $2^{\log n}$ and then noting that every square-root operation cuts the exponent in half.

