



THE UNIVERSITY OF TEXAS
AT AUSTIN

CS383C NUMERICAL ANALYSIS
HW07 Cholesky Factorization

Edited by L^AT_EX

Department of Computer Science

STUDENT

Jimmy Lin
xl5224

COURSE COORDINATOR

Robert A. van de Geijn

UNIQUE NUMBER

53180

RELEASE DATE

Oct. 24 2014

DUE DATE

Oct. 30 2014

TIME SPENT

2 hours

October 26, 2014

Exercises

- Exercise 3.** Prove that $A = B^H B$ is HPD. **2**
- Exercise 4.** Show that diagonal elements are real and positive. **2**
- Exercise 14.** Implement Cholesky Factorization **3**
- Exercise 15.** Relationship of Cholesky Factorization and QR **3**

Exercise 3. Prove that $A = B^H B$ is HPD.

To prove that $A = B^H B$ is HPD, we need to show both of the following:

- $A^H = A$.

Proof.

$$A^H = (B^H B)^H = B^H (B^H)^H = B^H B = A \quad (1)$$

□

- $\forall x \neq 0, x^H A x > 0$.

Proof. Let x be arbitrary non-zero vector in \mathbb{C}^n

$$x^H A x = x^H B^H B x = (B x)^H B x = \|B x\|_2^2 > 0 \quad (2)$$

Note that for $x \neq 0$, $B x = \sum_i B_i x_i \neq 0$, otherwise B is not linearly independent columns. □

Since two properties above are proven, then we can conclude that

If $B \in \mathbb{C}^{m \times n}$ has linearly independent columns, then $A = B^H B$ is HPD.

Exercise 4. Show that diagonal elements are real and positive.

- diagonal elements of $A \in \mathbb{C}^{m \times m}$ are real.

Proof. Since A is HPD, then $A^H = A$. Hence, for diagonal elements $\theta_0, \theta_1, \dots, \theta_{m-1}$, then

$$\forall i = 0, \dots, m-1, \theta_i^H = \theta_i \quad (3)$$

Let $\theta_i = x_i + y_i j$, where j denotes imaginary unit, then

$$\forall i = 0, \dots, m-1, -y_i = y_i \quad (4)$$

That tells us

$$\forall i = 0, \dots, m-1, y_i = 0 \quad (5)$$

Then it can be concluded that

all diagonal elements $\theta_0, \theta_1, \dots, \theta_{m-1}$ are real. (all imaginary part is zero.)

□

- diagonal elements of $A \in \mathbb{C}^{m \times m}$ are positive.

Proof. Since $A \in \mathbb{C}^{m \times m}$ is HPD, then

$$\forall x \neq 0, x^H A x > 0 \quad (6)$$

Let e_0, \dots, e_{m-1} denotes unit vector (whose imaginary part is zero) that spans through the whole \mathbb{C}^m . And let $\theta_0, \theta_1, \dots, \theta_{m-1}$ denotes diagonal elements of HPD matrix A .

$$\forall i = 0, \dots, m-1, \theta_i = e_i^H A e_i > 0 \quad (7)$$

Note that the $e_i^H A e_i \neq 0$ since $e_i \neq 0$. Hence, it can be conclude that

all diagonal elements $\theta_0, \theta_1, \dots, \theta_{m-1}$ are positive.

□

Exercise 14. Implement Cholesky Factorization

```
% Copyright 2014 The University of Texas at Austin
%
% For licensing information see
%     http://www.cs.utexas.edu/users/flame/license.html
%
% Programmed by: Jimmy Lin
%     linxin@gmail.com

function [ A_out ] = CHOL_unb( A )

[ ATL, ATR, ...
  ABL, ABR ] = FLA_Part_2x2( A, ...
                             0, 0, 'FLA_TL' );

while ( size( ATL, 1 ) < size( A, 1 ) )

    [ A00, a01, A02, ...
      a10t, alpha11, a12t, ...
      A20, a21, A22 ] = FLA_Repart_2x2_to_3x3( ATL, ATR, ...
                                                ABL, ABR, ...
                                                1, 1, 'FLA_BR' );

    %-----%

    a01 = zeros(size(a01));
    A02 = zeros(size(A02));
    a12t = zeros(size(a12t));
    alpha11 = sqrt(alpha11);
    a21 = a21 / alpha11;
    A22 = A22 - tril (a21 * a21');

    %-----%

    [ ATL, ATR, ...
      ABL, ABR ] = FLA_Cont_with_3x3_to_2x2( A00, a01, A02, ...
                                              a10t, alpha11, a12t, ...
                                              A20, a21, A22, ...
                                              'FLA_TL' );

end

A_out = [ ATL, ATR
         ABL, ABR ];

return
```

Exercise 15. Relationship of Cholesky Factorization and QR

Proof. For matrix $B \in \mathbb{C}^{m \times n}$ with linearly independent columns, it has a unique QR factorization such that $B = QR$, where $Q \in \mathbb{C}^{m \times n}$ and $R \in \mathbb{C}^{n \times n}$. And then for HPD matrix A , we have

$$A = B^H B = (QR)^H QR = R^H Q^H QR = \underbrace{R^H}_L \underbrace{R}_{L^H} \quad (8)$$

□