



THE UNIVERSITY OF TEXAS
AT AUSTIN

CS383C NUMERICAL ANALYSIS

Homework 04

Edited by L^AT_EX

Department of Computer Science

STUDENT

Jimmy Lin

xl5224

COURSE COORDINATOR

Robert A. Van De Geijn

UNIQUE NUMBER

53180

RELEASE DATE

Oct. 02 2014

DUE DATE

Oct. 10 2014

TIME SPENT

10 hours

October 10, 2014

Exercises

I	Exercises on Solving LLS Problems	2
Exercise 2.		2
Exercise 3.		2
Exercise 4.		3
Exercise 5.		4
II	Exercises on Conditioning	5
Exercise 1.		5
Exercise 2.		6
Exercise 3.		7
Exercise 4.		8
4.1	8
4.2	8
Exercise 5.		8

Part I

Exercises on Solving LLS Problems

Exercise 2.

We can prove the existence of LQ factorization for $A_{m \times n} (m < n)$ by induction on m .

- Base Case: $m = 1$

$$A = (a_0^H) = l_{00}q_0^H = \left(\begin{array}{c|c} l_{00} & 0 \end{array} \right) \left(\begin{array}{c} q_0^H \\ 0 \end{array} \right) \quad (1)$$

Let $L_L = l_{00}$ and $Q_T = q_0^H$, we have

$$A = \left(\begin{array}{c|c} L_L & 0 \end{array} \right) \left(\begin{array}{c} Q_T \\ 0 \end{array} \right) = L_L Q_T \quad (2)$$

holds for $m = 1$.

- Inductive Case: assume that existence of LQ factorization holds for $m - 1$, then we show that it also holds for m .

That is, we partition

$$A = \left(\begin{array}{c} A_0 \\ a_1^H \end{array} \right) \quad (3)$$

Assume that $A_0 = L_{00}Q_0$ holds, then we show that $\exists L_L, Q_T$, s.t. $A = L_L Q_T$ holds.

Let

$$L_L = \left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^H & \rho_{11} \end{array} \right), \quad Q_T = \left(\begin{array}{c} Q_0 \\ q_1^H \end{array} \right) \quad (4)$$

where $l_{10}^H = a_1^H Q_0^H$ and $\rho_{11} = \|a_1^H - l_{10}^H Q_0\|_2$. Then we have

$$L_L Q_T = \left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^H & \rho_{11} \end{array} \right) \left(\begin{array}{c} Q_0 \\ q_1^H \end{array} \right) = \left(\begin{array}{c} L_{00}Q_0 \\ l_{10}^H Q_0 + \rho_{11} q_1^H \end{array} \right) = \left(\begin{array}{c} L_{00}Q_0 \\ a_1^H \end{array} \right) = \left(\begin{array}{c} A_0 \\ a_1^H \end{array} \right) = A \quad (5)$$

- By principle of induction, $A = L_L Q_T$ holds for arbitrary m .

Exercise 3.

$$\min_z \|Az - y\|_2 = \min_z \|LQz - y\|_2 \quad (6)$$

$$= \min_{\{z=Q^H w\}} \|LQQ^H w - y\|_2 \quad (7)$$

$$= \min_{\{z=Q^H w\}} \|Lw - y\|_2 \quad (8)$$

$$= \min_{\{z=Q^H w\}} \left\| \left(\begin{array}{c|c} L_L & 0 \end{array} \right) \left(\begin{array}{c} w_T \\ w_B \end{array} \right) - y \right\|_2 \quad (9)$$

$$= \min_{\{z=Q^H w\}} \|L_L w_T - y\|_2 \quad (10)$$

which derives solution for $w_T = L_L^{-1}y$. And then the general solution is

$$z = Q^H w = \left(\begin{array}{c|c} Q_T^H & Q_B^H \end{array} \right) \left(\begin{array}{c} w_T \\ w_B \end{array} \right) = Q_T^H w_T + Q_B^H w_B = Q_T^H L_L^{-1}y + Q_B^H w_B \quad (11)$$

where w_B can be arbitrary vector in \mathbb{C}^{n-r} .

Exercise 4.

```

%% Homework 03: LQ Factorization
% Copyright 2014 The University of Texas at Austin
%
% For licensing information see
%     http://www.cs.utexas.edu/users/flame/license.html
%
% Programmed by: Jimmy Lin
%     jimmylin@utexas.edu
function [ A_out, L_out, Q_out ] = LG_CGS_unb( A, L, Q )
    [ AT, ...
      AB ] = FLA_Part_2x1( A, ...
                          0, 'FLA.TOP' );

    [ LTL, LTR, ...
      LBL, LBR ] = FLA_Part_2x2( L, ...
                                0, 0, 'FLA.TL' );

    [ QT, ...
      QB ] = FLA_Part_2x1( Q, ...
                          0, 'FLA.TOP' );
    while ( size( AT, 1 ) < size( A, 1 ) )
        [ A0, ...
          alt, ...
          A2 ] = FLA_Repart_2x1_to_3x1( AT, ...
                                       AB, ...
                                       1, 'FLA.BOTTOM' );

        [ L00,  l01,      L02,  ...
          l10t, lambda11, l12t, ...
          L20,  l21,      L22 ] = FLA_Repart_2x2_to_3x3( LTL, LTR, ...
                                                         LBL, LBR, ...
                                                         1, 1, 'FLA.BR' );

        [ Q0, ...
          qlt, ...
          Q2 ] = FLA_Repart_2x1_to_3x1( QT, ...
                                       QB, ...
                                       1, 'FLA.BOTTOM' );

        %-----%
        l10t = alt * Q0;
        temp = alt - l10t * Q0;
        lambda11 = norm(temp, 2);
        qlt = temp / lambda11;
        %-----%

        [ AT, ...
          AB ] = FLA_Cont_with_3x1_to_2x1( A0, ...
                                       alt, ...
                                       A2, ...
                                       'FLA.TOP' );

        [ LTL, LTR, ...
          LBL, LBR ] = FLA_Cont_with_3x3_to_2x2( L00,  l01,      L02,  ...
                                                  l10t, lambda11, l12t, ...
                                                  L20,  l21,      L22, ...
                                                  'FLA.TL' );

        [ QT, ...
          QB ] = FLA_Cont_with_3x1_to_2x1( Q0, ...
                                       qlt, ...
                                       Q2, ...
                                       'FLA.TOP' );

    end
    A_out = [ AT
              AB ];
    L_out = [ LTL, LTR
              LBL, LBR ];
    Q_out = [ QT
              QB ];
return

```

Exercise 5.

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%% Homework 03: Householder LQ Transformation
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

% Copyright 2014 The University of Texas at Austin
%
% For licensing information see
%      http://www.cs.utexas.edu/users/flame/license.html
%
% Programmed by: Jimmy Lin
%      jimmylin@utexas.edu

function [ A_out, T_out ] = HLQ_unb( A, T )

[ ATL, ATR, ...
  ABL, ABR ] = FLA_Part_2x2( A, ...
                             0, 0, 'FLA_TL' );

[ TT, ...
  TB ] = FLA_Part_2x1( T, ...
                       0, 'FLA_TOP' );

while ( size( ATL, 1 ) < size( A, 1 ) )

    [ A00, a01,      A02, ...
      a10t, alpha11, a12t, ...
      A20, a21,      A22 ] = FLA_Repart_2x2_to_3x3( ATL, ATR, ...
                                                    ABL, ABR, ...
                                                    1, 1, 'FLA_BR' );

    [ T0, ...
      t1t, ...
      T2 ] = FLA_Repart_2x1_to_3x1( TT, ...
                                    TB, ...
                                    1, 'FLA_BOTTOM' );

    %-----%

    [ alpha11, temp, t1t ] = Housev( alpha11, a12t );
    a12t = temp;
    w21 = (a21 + A22 * a12t) / t1t;
    a21 = a21 - w21;
    A22 = A22 - w21 * a12t;

    %-----%

    [ ATL, ATR, ...
      ABL, ABR ] = FLA_Cont_with_3x3_to_2x2( A00, a01,      A02, ...
                                              a10t, alpha11, a12t, ...
                                              A20, a21,      A22, ...
                                              'FLA_TL' );

    [ TT, ...
      TB ] = FLA_Cont_with_3x1_to_2x1( T0, ...
                                       t1t, ...
                                       T2, ...
                                       'FLA_TOP' );

end

A_out = [ ATL, ATR
          ABL, ABR ];
T_out = [ TT
          TB ];

return

```

Part II

Exercises on Conditioning

Exercise 1.

Show that, for a consistent matrix norm, $\kappa(A) \geq 1$.

Proof.

$$\kappa(A) = \|A\| \cdot \|A^{-1}\| \geq \|AA^{-1}\| = \|I\| = 1 \quad (12)$$

Note that the above $\|\cdot\|$ was for arbitrary induced matrix norm. □

Lemma 1. For arbitrary matrix A and B , $\|AB\| \leq \|A\| \cdot \|B\|$.

Proof.

$$\|AB\| = \sup_{x \neq 0} \frac{\|ABx\|}{\|x\|} = \sup_{x \neq 0} \frac{\|A(Bx)\|}{\|x\|} \quad (13)$$

$$\leq \sup_{x \neq 0} \frac{\|A\| \cdot \|Bx\|}{\|x\|} \quad (14)$$

$$\leq \sup_{x \neq 0} \frac{\|A\| \cdot \|B\| \cdot \|x\|}{\|x\|} \quad (15)$$

$$= \|A\| \cdot \|B\| \quad (16)$$

Hence, it is concluded that $\|AB\| \leq \|A\| \cdot \|B\|$. □

Lemma 2. For arbitrary norm $\|\cdot\|$ and identity matrix I , $\|I\| = 1$.

Proof.

$$\|I\| = \sup_{x \neq 0} \frac{\|I \cdot x\|}{\|x\|} = \sup_{x \neq 0} \frac{\|x\|}{\|x\|} = 1 \quad (17)$$

□

Exercise 2.

If A has linearly independent columns, show that $\|(A^H A)^{-1} A^H\|_2 = \frac{1}{\sigma_{n-1}}$, where σ_{n-1} equals the smallest singular value of A .

Proof. Let U , Σ and V be singular value decomposition of A , such that $A = U\Sigma V^H$.

$$\|(A^H A)^{-1} A^H\|_2 = \|((U\Sigma V^H)^H U \Sigma V^H)^{-1} (U\Sigma V^H)^H\|_2 \quad (18)$$

$$= \|(V\Sigma^H U^H U \Sigma V^H)^{-1} V\Sigma^H U^H\|_2 \quad (19)$$

$$= \|(V\Sigma^H \Sigma V^H)^{-1} V\Sigma^H U^H\|_2 \quad (20)$$

$$= \|V^{-H} \Sigma^{-1} \Sigma^{-H} V^{-1} V\Sigma^H U^H\|_2 \quad (21)$$

$$= \|V^{-H} \Sigma^{-1} \Sigma^{-H} \Sigma^H U^H\|_2 \quad (22)$$

$$= \|V^{-H} \Sigma^{-1} U^H\|_2 \quad (23)$$

$$= \|V \Sigma^{-1} U^H\|_2 \quad (24)$$

$$= \|\Sigma^{-1}\|_2 \quad (25)$$

$$= \frac{1}{\sigma_{n-1}} \quad (26)$$

□

Lemma 3. (Unitary Invariance) For arbitrary unitary matrix U ,

$$\|UA\|_2 = \|AU\|_2 = \|A\|_2 \quad (27)$$

Lemma 4. For arbitrary diagonal matrix Σ ,

$$\|\Sigma^{-1}\|_2 = \frac{1}{\sigma_{n-1}} \quad (28)$$

where, σ_{n-1} is the least entry of Σ .

Note that above two lemmas have been proven in exercises of previous notes.

Exercise 3.

Let A have linearly independent columns. Show that $\kappa_2(A^H A) = \kappa_2(A)^2$.

Proof. We achieve the proof by employing SVD over A . Let unitary matrix U , diagonal matrix Σ and unitary matrix V be singular value decomposition of A , such that $A = U\Sigma V^H$. We start from the definition of condition number $\kappa_2(\cdot)$.

$$\kappa_2(A^H A) = \|A^H A\|_2 \cdot \|(A^H A)^{-1}\|_2 \quad (29)$$

Then we discuss the term $\|A^H A\|_2$ and $\|(A^H A)^{-1}\|_2$ respectively.

$$\|A^H A\|_2 = \|(U\Sigma V^H)^H U\Sigma V^H\|_2 \quad (30)$$

$$= \|V\Sigma^H U^H U\Sigma V^H\|_2 \quad (31)$$

$$= \|V\Sigma^H \Sigma V^H\|_2 \quad (32)$$

$$= \|\Sigma^H \Sigma\|_2 \quad (33)$$

$$= \sigma_0^2 \quad (34)$$

$$= \|A\|_2^2 \quad (35)$$

Note that σ_0 is the largest singular value of matrix A and also the largest entry of Σ .

$$\|(A^H A)^{-1}\|_2 = \|((U\Sigma V^H)^H U\Sigma V^H)^{-1}\|_2 \quad (36)$$

$$= \|(V\Sigma^H U^H U\Sigma V^H)^{-1}\|_2 \quad (37)$$

$$= \|(V\Sigma^H \Sigma V^H)^{-1}\|_2 \quad (38)$$

$$= \|V^{-H} \Sigma^{-1} \Sigma^{-H} V^{-1}\|_2 \quad (39)$$

$$= \|\Sigma^{-1} \Sigma^{-H}\|_2 \quad (40)$$

$$= \|\Sigma^{-1} \Sigma^{-1}\|_2 \quad (41)$$

$$= \frac{1}{\sigma_{n-1}^2} \quad (42)$$

$$= \|A^{-1}\|_2^2 \quad (43)$$

Now we have

$$\|A^H A\|_2 = \|A\|_2^2 \quad (44)$$

$$\|(A^H A)^{-1}\|_2 = \|A^{-1}\|_2^2 \quad (45)$$

Then

$$\kappa_2(A^H A) = \|A^H A\|_2 \cdot \|(A^H A)^{-1}\|_2 \quad (46)$$

$$= \|A\|_2^2 \cdot \|A^{-1}\|_2^2 \quad (47)$$

$$= (\|A\|_2 \cdot \|A^{-1}\|_2)^2 \quad (48)$$

$$= \kappa_2(A)^2 \quad (49)$$

Hence, it can be concluded that

$$\kappa_2(A^H A) = \kappa_2(A)^2 \quad (50)$$

□

Exercise 4.

4.1

(Only-If) If $Ax = y$, multiply both side with A^H and then we have $A^H Ax = A^H y$.

(If) If $A^H Ax = A^H y$, then multiply both side with $(A^H)^{-1}$ (inverse exists because A is full-rank) and we have $(A^H)^{-1} A^H Ax = (A^H)^{-1} A^H y$. Since $(A^H)^{-1} A^H = I$, it comes out $Ax = y$.

4.2

For normal equation method, the solution of LLS can be derived by

$$x = (A^H A)^{-1} A^H y \quad (51)$$

Let $B = (A^H A)^{-1} A^H$, then the condition number of normal equation program is

$$\kappa(B^{-1}) = \|((A^H A)^{-1} A^H)^{-1}\| \cdot \|(A^H A)^{-1} A^H\| \quad (52)$$

$$= \sigma_0 \cdot \frac{1}{\sigma_{n-1}} \quad (53)$$

$$= \frac{\sigma_0}{\sigma_{n-1}} \quad (54)$$

$$= \kappa(A) \quad (55)$$

By this, we can conclude that the condition number of normal equation program is not necessarily square of $\kappa(A)$.

Exercise 5.

Let $U \in \mathbb{C}^{n \times n}$ be unitary. Show that $\kappa_2(U) = 1$.

Proof.

$$\kappa_2(U) = \|U\|_2 \|U^{-1}\|_2 \quad (56)$$

$$= \sup_{x \neq 0} \frac{\|Ux\|_2}{\|x\|_2} \cdot \sup_{y \neq 0} \frac{\|U^{-1}y\|_2}{\|y\|_2} \quad (57)$$

$$= \sup_{x \neq 0} \frac{\|x\|_2}{\|x\|_2} \cdot \sup_{y \neq 0} \frac{\|y\|_2}{\|y\|_2} \quad (58)$$

$$= 1 \cdot 1 \quad (59)$$

$$= 1 \quad (60)$$

□

Lemma 5. For arbitrary unitary matrix U , its inverse U^{-1} is still unitary.

Proof.

$$UU^H = I \quad (61)$$

We multiply both side on the left with U^{-1} and U^{-H} , it comes

$$U^{-H} U^{-1} U U^H = U^{-H} U^{-1} \quad (62)$$

That is

$$(U^{-1})^H U^{-1} = I \quad (63)$$

Since U is unitary, then U is square matrix and so as U^{-1} . Then it is concluded that U^{-1} is unitary. □