



THE UNIVERSITY OF TEXAS
AT AUSTIN

CS383C NUMERICAL ANALYSIS
HW05 Numerical Stability

Edited by L^AT_EX

Department of Computer Science

STUDENT

Jimmy Lin
xl5224

COURSE COORDINATOR

Robert A. van de Geijn

UNIQUE NUMBER

53180

RELEASE DATE

Oct. 08 2014

DUE DATE

Oct. 14 2014

TIME SPENT

10 hours

October 11, 2014

Exercises

Exercise 3.	2
3.1	2
3.2	2
Exercise 10.	2
Exercise 12.	3
12.1	3
12.2	3
12.3	3
Exercise 13.	4
Exercise 15.	4
Exercise 18.	5
18.1	5
18.2	5
Exercise 19.	6
19.1 $k = 0$	6
19.2 $k > 0$	6
Exercise 23.	6
23.1	6
23.2	6
Exercise 25.	7
Exercise 27.	8

Exercise 3.

3.1

Write 1 as floating number.

$$.1 \underbrace{00 \cdots 0}_{t-1} \times 2^1 \quad (1)$$

3.2

Show that $\mathbf{u} = \frac{1}{2} \cdot 2^{1-t}$

Proof. Let $\chi = .\delta_0\delta_1 \cdots \delta_{t-1}\delta_t \cdots$ and $\check{\chi}$ to be the value stored in t-digit floating number with rounding mechanism. Then if $\delta_t = 0$, then $\chi = \check{\chi}$ and $|\delta\chi| = 0 \leq 2^{e-t-1}$. But if $\delta_t = 1$, due to the rounding mechanism, then $\chi < \check{\chi}$ and

$$|\delta\chi| = |\chi - \check{\chi}| = |.\delta_0\delta_1 \cdots \delta_{t-1}\delta_t \cdots \times 2^e - .\delta_0\delta_1 \cdots \delta'_{t-1} \times 2^e| \leq \underbrace{.00 \cdots 0}_t 1 \times 2^e = 2^{e-t-1} \quad (2)$$

For χ , since $\delta_0 = 1$ (normalized)

$$|\chi| = |.\delta_0\delta_1 \cdots \times 2^e| \geq .1 \times 2^e \geq 2^{e-1} \quad (3)$$

Thus,

$$\frac{|\delta\chi|}{|\chi|} \leq \frac{2^{e-t-1}}{2^{e-1}} = \frac{1}{2} \cdot 2^{1-t} \quad (4)$$

Then,

$$|\delta\chi| \leq \frac{1}{2} \cdot 2^{1-t} |\chi| \quad (5)$$

Now, we have

$$\mathbf{u} = \frac{1}{2} 2^{1-t} \quad (6)$$

□

Exercise 10.

Show that $|AB| \leq |A||B|$.

Proof. Let $C = AB$. And the (i, j) entry of $|C|$ is given by

$$|c_{i,j}| = \left| \sum_{p=0}^k a_{i,p} b_{p,j} \right| \leq \sum_{p=0}^k |a_{i,p} b_{p,j}| \leq \sum_{p=0}^k |a_{i,p}| |b_{p,j}| \quad (7)$$

which equals (i, j) entry of $|A||B|$. Hence, we have

$$|AB| \leq |A||B| \quad (8)$$

□

Exercise 12.

12.1

Show that if $|A| \leq |B|$, then $\|A\|_1 \leq \|B\|_1$.

Proof. Partition $A_{m \times n} = \left(\begin{array}{c|c|c|c} a_0 & a_1 & \dots & a_{n-1} \end{array} \right)$ where a_j indicates the j -th column of matrix A . Similarly, we partition $B_{m \times n} = \left(\begin{array}{c|c|c|c} b_0 & b_1 & \dots & b_{n-1} \end{array} \right)$ where b_j indicates the j -th column of matrix B . Then we use a_{ij} and b_{ij} to denote i -th element of a_j and b_j respectively.

$$\|A\|_1 = \max_{0 \leq j < n} \|a_j\|_1 = \max_{0 \leq j < n} \sum_{i=0}^{m-1} |a_{ij}| \leq \max_{0 \leq j < n} \sum_{i=0}^{m-1} |b_{ij}| = \max_{0 \leq j < n} \|b_j\|_1 = \|B\|_1 \quad (9)$$

□

Lemma 1. For arbitrary matrix $A = \left(\begin{array}{c|c|c|c} a_0 & a_1 & \dots & a_{n-1} \end{array} \right)$, $\|A\|_1 = \max_{0 \leq j < n} \|a_j\|_1$.

Proof. This lemma has been proved in Notes on Norms. □

12.2

Show that if $|A| \leq |B|$, then $\|A\|_\infty \leq \|B\|_\infty$.

Proof. Partition $A_{m \times n} = \left(\begin{array}{c} \frac{a_0}{a_1} \\ \vdots \\ \frac{a_{m-1}}{a_{m-1}} \end{array} \right)$, where a_i indicates the i -th row of matrix A . Similarly, we

partition $B_{m \times n} = \left(\begin{array}{c} \frac{b_0}{b_1} \\ \vdots \\ \frac{b_{m-1}}{b_{m-1}} \end{array} \right)$, where b_i indicates the i -th row of matrix B . Then we use a_{ij} and b_{ij}

to denote j -th element of a_i and b_i respectively.

$$\|A\|_\infty = \max_{0 \leq i < m} \|a_i\|_1 = \max_{0 \leq i < m} \sum_{j=0}^{n-1} |a_{ij}| \leq \max_{0 \leq i < m} \sum_{j=0}^{n-1} |b_{ij}| = \max_{0 \leq i < m} \|b_i\|_1 = \|B\|_\infty \quad (10)$$

□

Lemma 2. For arbitrary matrix $A = \left(\begin{array}{c} \frac{a_0}{a_1} \\ \vdots \\ \frac{a_{m-1}}{a_{m-1}} \end{array} \right)$, $\|A\|_\infty = \max_{0 \leq i < m} \|a_i\|_1$.

Proof. This lemma has been proved in Notes on Norms. □

12.3

Show that if $|A| \leq |B|$, then $\|A\|_F \leq \|B\|_F$. Let $A, B \in \mathbb{R}^{m \times n}$ and a_{ij} , b_{ij} be (i, j) entry of A , B respectively.

Proof.

$$\|A\|_F^2 = \sum_{i=0}^{m-1} \sum_{j=0}^{n-1} |a_{ij}|^2 \leq \sum_{i=0}^{m-1} \sum_{j=0}^{n-1} |b_{ij}|^2 = \|B\|_F^2 \quad (11)$$

□

Exercise 13.

$$\check{\kappa} = [(\chi_0\psi_0 + \chi_1\psi_1) + \chi_2\psi_2] \quad (12)$$

$$= [[\chi_0\psi_0 + \chi_1\psi_1] + [\chi_2\psi_2]] \quad (13)$$

$$= [[[\chi_0\psi_0] + [\chi_1\psi_1]] + [\chi_2\psi_2]] \quad (14)$$

$$= [[\chi_0\psi_0(1 + \epsilon_*^{(0)}) + \chi_1\psi_1(1 + \epsilon_*^{(1)})] + \chi_2\psi_2(1 + \epsilon_*^{(2)})] \quad (15)$$

$$= [(\chi_0\psi_0(1 + \epsilon_*^{(0)}) + \chi_1\psi_1(1 + \epsilon_*^{(1)}))(1 + \epsilon_+^{(1)}) + \chi_2\psi_2(1 + \epsilon_*^{(2)})] \quad (16)$$

$$= \left((\chi_0\psi_0(1 + \epsilon_*^{(0)}) + \chi_1\psi_1(1 + \epsilon_*^{(1)}))(1 + \epsilon_+^{(1)}) + \chi_2\psi_2(1 + \epsilon_*^{(2)}) \right) (1 + \epsilon_+^{(2)}) \quad (17)$$

$$= \chi_0\psi_0(1 + \epsilon_*^{(0)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) + \chi_1\psi_1(1 + \epsilon_*^{(1)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) + \chi_2\psi_2(1 + \epsilon_*^{(2)})(1 + \epsilon_+^{(2)}) \quad (18)$$

$$= \begin{pmatrix} \chi_0 \\ \chi_1 \\ \chi_2 \end{pmatrix}^T \begin{pmatrix} \epsilon_0(1 + \epsilon_*^{(0)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) \\ \epsilon_1(1 + \epsilon_*^{(1)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) \\ \epsilon_2(1 + \epsilon_*^{(2)})(1 + \epsilon_+^{(2)}) \end{pmatrix} \quad (19)$$

$$= \begin{pmatrix} \chi_0 \\ \chi_1 \\ \chi_2 \end{pmatrix}^T \begin{pmatrix} (1 + \epsilon_*^{(0)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) & 0 & 0 \\ 0 & (1 + \epsilon_*^{(1)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) & 0 \\ 0 & 0 & (1 + \epsilon_*^{(2)})(1 + \epsilon_+^{(2)}) \end{pmatrix} \begin{pmatrix} \epsilon_0 \\ \epsilon_1 \\ \epsilon_2 \end{pmatrix} \quad (20)$$

$$= \begin{pmatrix} \chi_0(1 + \epsilon_*^{(0)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) \\ \chi_1(1 + \epsilon_*^{(1)})(1 + \epsilon_+^{(1)})(1 + \epsilon_+^{(2)}) \\ \chi_2(1 + \epsilon_*^{(2)})(1 + \epsilon_+^{(2)}) \end{pmatrix}^T \begin{pmatrix} \epsilon_0 \\ \epsilon_1 \\ \epsilon_2 \end{pmatrix} \quad (21)$$

Exercise 15.

Now we complete the missing part for the Inductive Step Case 1 of Lemma 14.

Proof. Case 1: $\prod_{i=0}^n (1 + \epsilon)^{\pm 1} = \prod_{i=0}^{n-1} (1 + \epsilon)^{\pm 1} (1 + \epsilon_n)$.

By the inductive hypothesis, there exists a θ_n such that

$$(1 + \theta_n) = \prod_{i=0}^{n-1} (1 + \epsilon_i)^{\pm 1} \text{ and } |\theta_n| \leq n\mathbf{u}/(1 - n\mathbf{u}) \quad (22)$$

Then

$$\prod_{i=0}^n (1 + \epsilon)^{\pm 1} = \left(\prod_{i=0}^{n-1} (1 + \epsilon)^{\pm 1} \right) (1 + \epsilon_n) = (1 + \theta_n)(1 + \epsilon_n) = 1 + \underbrace{\theta_n + \epsilon_n + \theta_n \cdot \epsilon_n}_{\theta_{n+1}} \quad (23)$$

which tells us how to pick up θ_{n+1} . Then

$$|\theta_{n+1}| = |\theta_n + \epsilon_n + \theta_n \cdot \epsilon_n| \quad (24)$$

$$\leq |\theta_n| + |\epsilon_n| + |\theta_n| \cdot |\epsilon_n| \quad (25)$$

$$= \frac{n\mathbf{u}}{1 - n\mathbf{u}} + \mathbf{u} + \frac{n\mathbf{u}}{1 - n\mathbf{u}} \cdot \mathbf{u} \quad (26)$$

$$= \frac{n\mathbf{u} + \mathbf{u} - n\mathbf{u}^2 + n\mathbf{u}^2}{1 - n\mathbf{u}} \quad (27)$$

$$= \frac{(n+1)\mathbf{u}}{1 - n\mathbf{u}} \quad (28)$$

$$\leq \frac{(n+1)\mathbf{u}}{1 - (n+1)\mathbf{u}} \quad (29)$$

□

Exercise 18.

18.1

Show that if $n, b \geq 1$, then $\gamma_n \leq \gamma_{n+b}$.

Proof.

$$\gamma_n = \frac{n\mathbf{u}}{1 - n\mathbf{u}} \leq \frac{n\mathbf{u}}{1 - (n+b)\mathbf{u}} \leq \frac{(n+b)\mathbf{u}}{1 - (n+b)\mathbf{u}} = \gamma_{n+b} \quad (30)$$

Note that since \mathbf{u} is extremely small, then $1 - n\mathbf{u} > 0$ and $1 - (n+b)\mathbf{u} > 0$. □

18.2

Show that if $n, b \geq 1$, then $\gamma_n + \gamma_b + \gamma_n\gamma_b \leq \gamma_{n+b}$.

Proof.

$$\gamma_n + \gamma_b + \gamma_n\gamma_b \quad (31)$$

$$= \frac{n\mathbf{u}}{1 - n\mathbf{u}} + \frac{b\mathbf{u}}{1 - b\mathbf{u}} + \frac{n\mathbf{u}}{1 - n\mathbf{u}} \cdot \frac{b\mathbf{u}}{1 - b\mathbf{u}} \quad (32)$$

$$= \frac{n\mathbf{u} - nb\mathbf{u}^2 + b\mathbf{u} - nb\mathbf{u}^2 + nb\mathbf{u}^2}{(1 - n\mathbf{u})(1 - b\mathbf{u})} \quad (33)$$

$$= \frac{n\mathbf{u} - nb\mathbf{u}^2 + b\mathbf{u}}{1 - n\mathbf{u} - b\mathbf{u} + nb\mathbf{u}^2} \quad (34)$$

$$\leq \frac{n\mathbf{u} + b\mathbf{u}}{1 - n\mathbf{u} - b\mathbf{u} + nb\mathbf{u}^2} \quad (35)$$

$$\leq \frac{n\mathbf{u} + b\mathbf{u}}{1 - n\mathbf{u} - b\mathbf{u}} \quad (36)$$

$$= \frac{(n+b)\mathbf{u}}{1 - (n+b)\mathbf{u}} \quad (37)$$

$$= \gamma_{n+b} \quad (38)$$

Note that since \mathbf{u} is extremely small, then $1 - n\mathbf{u} > 0$ and $1 - (n+b)\mathbf{u} > 0$. Also, note that $nb\mathbf{u}^2 \geq 0$. □

Exercise 19.**19.1** $k = 0$

Show that $\left(\frac{I + \Sigma^{(k)}}{0} \middle| \frac{0}{(1 + \epsilon_1)} \right) (1 + \epsilon_2) = I + \Sigma^{(k+1)}$

Proof. Given that if $k = 0$, then $\epsilon_1 = 0$ and Σ^0 is 0×0 matrix, we have

$$(1 + 0) \cdot (1 + \epsilon_2) = \underbrace{1}_I + \underbrace{\epsilon_2}_{\Sigma^{(1)}} \quad (39)$$

Thus, $\left(\frac{I + \Sigma^{(k)}}{0} \middle| \frac{0}{(1 + \epsilon_1)} \right) (1 + \epsilon_2) = I + \Sigma^{(k+1)}$ holds for $k = 0$. \square

19.2 $k > 0$

Proof. For arbitrary $k > 0$,

$$\left(\frac{I + \Sigma^{(k)}}{0} \middle| \frac{0}{(1 + \epsilon_1)} \right) (1 + \epsilon_2) \quad (40)$$

$$= \left(\frac{(I + \Sigma^{(k)})(1 + \epsilon_2)}{0} \middle| \frac{0}{(1 + \epsilon_1)(1 + \epsilon_2)} \right) \quad (41)$$

$$= \left(\frac{I + \epsilon_2 I + \Sigma^{(k)} + \epsilon_2 \Sigma^{(k)}}{0} \middle| \frac{0}{1 + \epsilon_1 + \epsilon_2 + \epsilon_1 \epsilon_2} \right) \quad (42)$$

$$= \underbrace{\left(\frac{I}{0} \middle| \frac{0}{1} \right)}_I + \underbrace{\left(\frac{\epsilon_2 I + \Sigma^{(k)} + \epsilon_2 \Sigma^{(k)}}{0} \middle| \frac{0}{\epsilon_1 + \epsilon_2 + \epsilon_1 \epsilon_2} \right)}_{\Sigma^{(k+1)}} \quad (43)$$

which tells us that $\left(\frac{I + \Sigma^{(k)}}{0} \middle| \frac{0}{(1 + \epsilon_1)} \right) (1 + \epsilon_2) = I + \Sigma^{(k+1)}$ holds for $k > 0$. \square

Exercise 23.**23.1**

Show that $\check{\kappa} = (x + \delta x)^T y$, where $|\delta x| \leq \gamma_n |x|$.

Proof. Let $\delta x = \Sigma^{(n)} x$, where $\Sigma^{(n)}$ is as in Theorem 20.

$$|\delta x| = |\Sigma^{(n)} x| = \begin{pmatrix} |\theta_n \chi_0| \\ |\theta_n \chi_1| \\ \vdots \\ |\theta_2 \chi_{n-1}| \end{pmatrix} \leq \begin{pmatrix} |\theta_n| |\chi_0| \\ |\theta_n| |\chi_1| \\ \vdots \\ |\theta_2| |\chi_{n-1}| \end{pmatrix} \leq |\theta_n| \begin{pmatrix} |\chi_0| \\ |\chi_1| \\ \vdots \\ |\chi_{n-1}| \end{pmatrix} \leq \gamma_n \begin{pmatrix} |\chi_0| \\ |\chi_1| \\ \vdots \\ |\chi_{n-1}| \end{pmatrix} = \gamma_n |x| \quad (44)$$

Thus, it can be concluded for the backward analysis that

$$|\delta x| \leq \gamma_n |x| \quad (45)$$

\square

23.2

Show that $\check{\kappa} = x^T (y + \delta y)$, where $|\delta y| \leq \gamma_n |y|$.

Proof. The proof for perturbation on input y is the same as that of perturbation on input x . \square

Exercise 25.

Exercise 27.