



THE UNIVERSITY OF TEXAS
AT AUSTIN

CS383C NUMERICAL ANALYSIS

Midterm Exam

Edited by L^AT_EX

Department of Computer Science

STUDENT

Jimmy Lin

xl5224

COURSE COORDINATOR

Robert A. van de Geijn

UNIQUE NUMBER

53180

RELEASE DATE

Oct. 16 2014

DUE DATE

Oct. 21 2014

TIME SPENT

10 hours

October 17, 2014

Exercises

Exercise 1.	2
(a) Prove that $\ \cdot\ $ is a vector norm.	2
(b) Show that $\ \cdot\ $ is a matrix norm.	3
Exercise 2.	4
(a) Show that $(..)^{-1} = (..)$	4
(b) Show that $H_1 H_0 = ..$	4
(c) Show that $(1 - \frac{1}{\tau_1 u_1 u_1^T}).. = I - ..$	4
(d) Show that $H_{k-1} \cdots H_1 H_0 = I - U L^{-1} U^H$	5
Exercise 3.	6
(a) Relationship between m , n and r	6
(b) Reduced SVD in terms of U , V and Σ	6
Exercise 4.	7
(a) $\psi_1 = (v_{11} - \delta v_{11})\check{\chi}_1$ where δv_{11} is small.	7
(b)	7
(c) Induction for the desired result	8

Exercise 1.

(a) Prove that $\|\cdot\|$ is a vector norm.

To prove $\|\cdot\|$ is a vector norm, we have to show all the following three property holds:

- If $x \neq 0$, then $\|x\| > 0$.
- For arbitrary scalar α and vector x , $\|\alpha x\| = |\alpha| \cdot \|x\|$.
- For arbitrary vector x and y , $\|x + y\| \leq \|x\| + \|y\|$.

Now we turn to prove each of them individually.

Proof. Since D is diagonal with positive diagonal elements, then $\forall i, \delta_i > 0$ holds. Besides, since $x \neq 0$, then $\exists i, x_i^2 > 0$.

$$\|x\| = \sqrt{x^T D x} = \sqrt{\sum_{i=0}^{m-1} \delta_i x_i^2} > 0 \quad x \neq 0 \quad (1)$$

Therefore, we proved that if $x \neq 0$, then $\|x\| > 0$. \square

Proof.

$$\|\alpha x\| = \sqrt{(\alpha x)^T D (\alpha x)} = \sqrt{\sum_{i=0}^{m-1} \alpha^2 \delta_i x_i^2} = \sqrt{\alpha^2 \sum_{i=0}^{m-1} \delta_i x_i^2} = |\alpha| \sqrt{\sum_{i=0}^{m-1} \delta_i x_i^2} = |\alpha| \cdot \|x\| \quad (2)$$

\square

Proof.

$$\|x + y\|^2 = (\sqrt{(x + y)^T D (x + y)})^2 = x^T D x + x^T D y + 2x^T D y \quad (3)$$

$$(\|x\| + \|y\|)^2 = (\sqrt{x^T D x})^2 + (\sqrt{y^T D y})^2 = x^T D x + y^T D y + 2\sqrt{x^T D x} \sqrt{y^T D y} \quad (4)$$

Now we show that $x^T D y \leq \sqrt{x^T D x} \sqrt{y^T D y}$, which is equivalent to

$$\left(\sum_{i=0}^{m-1} \delta_i x_i y_i\right)^2 \leq \left(\sum_{i=0}^{m-1} \delta_i x_i^2\right) \left(\sum_{i=0}^{m-1} \delta_i y_i^2\right) \quad (5)$$

$$\iff \sum_{i=0}^{m-1} \sum_{j=0}^{m-1} \delta_i \delta_j x_i y_i x_j y_j \leq \sum_{i=0}^{m-1} \sum_{j=0}^{m-1} \delta_i \delta_j x_i^2 y_j^2 \quad (6)$$

$$\iff \sum_{i=0}^{m-1} \sum_{j=0}^{i-1} \delta_i \delta_j (2x_i y_i x_j y_j) + \sum_{i=0}^{m-1} \delta_i^2 x_i y_i x_i y_i \leq \sum_{i=0}^{m-1} \sum_{j=0}^{i-1} \delta_i \delta_j (x_i^2 y_j^2 + x_j^2 y_i^2) + \sum_{i=0}^{m-1} \delta_i^2 x_i^2 y_i^2 \quad (7)$$

$$\iff \sum_{i=0}^{m-1} \sum_{j=0}^{i-1} \delta_i \delta_j (2x_i y_i x_j y_j) \leq \sum_{i=0}^{m-1} \sum_{j=0}^{i-1} \delta_i \delta_j (x_i^2 y_j^2 + x_j^2 y_i^2) \quad (8)$$

which obviously holds if we apply basic inequality on each (i, j) pair,

$$2x_i y_i x_j y_j = 2(x_i y_j)(x_j y_i) \leq x_i^2 y_j^2 + x_j^2 y_i^2 \quad (9)$$

Since we have proved $x^T D y \leq \sqrt{x^T D x} \sqrt{y^T D y}$, then

$$\|x + y\|^2 \leq (\|x\| + \|y\|)^2 \quad (10)$$

which leads to desired triangle inequality since $\|\cdot\| \geq 0$,

$$\|x + y\| \leq \|x\| + \|y\| \quad (11)$$

\square

Since all three property have been proved, it can be concluded that

$$\|\cdot\| \text{ defined by } \|x\| = \sqrt{x^T D x} \text{ is a norm.} \quad (12)$$

(b) Show that $\|\cdot\|$ is a matrix norm.

To prove $\|\cdot\|$ is a matrix norm, we have to show all the following three property holds:

- If $A \neq 0$, then $\|A\| > 0$.
- For arbitrary scalar α and matrix A , $\|\alpha A\| = |\alpha| \cdot \|A\|$.
- For arbitrary matrix A and B , $\|A + B\| \leq \|A\| + \|B\|$.

Now we turn to prove each of them individually.

Proof. Since D is diagonal with positive diagonal elements, then $\forall i, \delta_i > 0$ holds. And Since $A \neq 0$, then there is at least one column vector that is not zero vector. Let us assume it is the j -th column, a_j , that is non-zero. Then

$$\|A\| = \|DA\|_2 = \max_{\|x\|_2 \neq 0} \frac{\|DAx\|_2}{\|x\|_2} \geq \frac{\|DAe_j\|_2}{\|e_j\|_2} > 0 \quad (13)$$

Therefore, we proved that if $A \neq 0$, then $\|A\| > 0$. □

Proof.

$$\|\alpha A\| = \|D(\alpha A)\|_2 \quad (14)$$

$$= \|\alpha DA\|_2 \quad (15)$$

$$= |\alpha| \cdot \|DA\|_2 \quad \|\cdot\|_2 \text{ is a norm.} \quad (16)$$

$$= |\alpha| \cdot \|A\| \quad (17)$$

Thus, we proved that $\|\alpha A\| = |\alpha| \cdot \|A\|$. □

Proof.

$$\|A + B\| = \|D(A + B)\|_2 \quad (18)$$

$$= \max_{x \neq 0} \frac{\|DAx + DBx\|_2}{\|x\|_2} \quad (19)$$

$$\leq \max_{x \neq 0} \frac{\|DAx\|_2 + \|DBx\|_2}{\|x\|_2} \quad (20)$$

$$= \max_{x \neq 0} \left(\frac{\|DAx\|_2}{\|x\|_2} + \frac{\|DBx\|_2}{\|x\|_2} \right) \quad (21)$$

$$\leq \max_{x \neq 0} \frac{\|DAx\|_2}{\|x\|_2} + \max_{y \neq 0} \frac{\|DBy\|_2}{\|y\|_2} \quad (22)$$

$$= \|DA\|_2 + \|DB\|_2 = \|A\| + \|B\| \quad (23)$$

Hence, we proved that $\|A + B\| \leq \|A\| + \|B\|$. □

Since all three property have been proved, it can be concluded that

$$\|\cdot\| \text{ defined by } \|A\| = \|DA\|_2 \text{ is a norm.} \quad (24)$$

Exercise 2.**(a) Show that** $(..)^{-1} = (..)$.*Proof.*

$$\left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^T & \lambda_{11} \end{array} \right) \left(\begin{array}{c|c} L_{00}^{-1} & 0 \\ \hline -\frac{l_{10}^T L_{00}^{-1}}{\lambda_{11}} & \frac{1}{\lambda_{11}} \end{array} \right) = \left(\begin{array}{c|c} L_{00} L_{00}^{-1} & 0 \\ \hline l_{10}^T L_{00}^{-1} - \frac{l_{10}^T L_{00}^{-1}}{\lambda_{11}} \cdot \lambda_{11} & \frac{1}{\lambda_{11}} \cdot \lambda_{11} \end{array} \right) = \left(\begin{array}{c|c} I & 0 \\ \hline 0 & 1 \end{array} \right) = I \quad (25)$$

Hence, it is concluded that (assume all inversion exists)

$$\left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^T & \lambda_{11} \end{array} \right)^{-1} = \left(\begin{array}{c|c} L_{00}^{-1} & 0 \\ \hline -\frac{l_{10}^T L_{00}^{-1}}{\lambda_{11}} & \frac{1}{\lambda_{11}} \end{array} \right) \quad (26)$$

□

(b) Show that $H_1 H_0 = ..$ According to given formula for H_i , we have

$$H_0 = I - \frac{1}{\tau_0} u_0 u_0^T \quad (27)$$

$$H_1 = I - \frac{1}{\tau_1} u_1 u_1^T \quad (28)$$

Now we manipulate $H_1 H_0$,

$$H_1 H_0 = \left(I - \frac{1}{\tau_1} u_1 u_1^T \right) \left(I - \frac{1}{\tau_0} u_0 u_0^T \right) \quad (29)$$

$$= I - \frac{1}{\tau_1} u_1 u_1^T - \frac{1}{\tau_0} u_0 u_0^T + \frac{1}{\tau_0 \tau_1} u_1 u_1^T u_0 u_0^T \quad (30)$$

$$= I - (u_0 | u_1) \left(\begin{array}{c|c} \frac{1}{\tau_0} & 0 \\ \hline -\frac{u_0^T u_1}{\tau_0 \tau_1} & \frac{1}{\tau_1} \end{array} \right) (u_0 | u_1)^T \quad (31)$$

$$= I - (u_0 | u_1) \left(\begin{array}{c|c} \tau_0 & 0 \\ \hline u_0^T u_1 & \tau_1 \end{array} \right)^{-1} (u_0 | u_1)^T \quad (32)$$

Hence, it is concluded that (assume all inversion exists)

$$H_1 H_0 = I - (u_0 | u_1) \left(\begin{array}{c|c} \tau_0 & 0 \\ \hline u_0^T u_1 & \tau_1 \end{array} \right)^{-1} (u_0 | u_1)^T \quad (33)$$

(c) Show that $(1 - \frac{1}{\tau_1 u_1 u_1^T}).. = I - ...$

$$\left(I - \frac{1}{\tau_1} u_1 u_1^T \right) \left(I - U_0 L_{00}^T U_0^T \right) = I - \frac{1}{\tau_1} u_1 u_1^T - U_0 L_{00}^T U_0^T + \frac{1}{\tau_1} u_1 u_1^T U_0 L_{00}^T U_0^T \quad (34)$$

$$= I - (U_0 | u_1) \left(\begin{array}{c|c} L_{00}^{-1} & 0 \\ \hline -u_1^T U_0 L_{00}^{-1} / \tau_1 & \frac{1}{\tau_1} \end{array} \right) (U_0 | u_1)^T \quad (35)$$

$$= I - (U_0 | u_1) \left(\begin{array}{c|c} L_{00} & 0 \\ \hline u_1^T U_0 & \tau_1 \end{array} \right)^{-1} (U_0 | u_1)^T \quad (36)$$

Thus, it is concluded that (assume all inversion exists)

$$\left(I - \frac{1}{\tau_1} u_1 u_1^T \right) \left(I - U_0 L_{00}^T U_0^T \right) = I - (U_0 | u_1) \left(\begin{array}{c|c} L_{00} & 0 \\ \hline u_1^T U_0 & \tau_1 \end{array} \right)^{-1} (U_0 | u_1)^T \quad (37)$$

(d) **Show that** $H_{k-1} \cdots H_1 H_0 = I - UL^{-1}U^H$

We can do mathematical induction on m . We separate the discussion of $k = 1$ and $k > 1$.

- Particular Case: $k = 1$,

Proof.

$$H_0 = I - \frac{1}{\tau_0} u_0 u_0^T = I - \underbrace{(u_0)}_{U_{(0)}} \underbrace{\left(\frac{1}{\tau_0}\right)}_{L_{(0)}^{-1}} \underbrace{(u_0)^T}_{U_{(0)}^T} \quad (38)$$

Hence, $H_{k-1} \cdots H_1 H_0 = I - UL^{-1}U^H$ holds for $k = 1$. \square

Note that this case is particular because it derives matrix which is indeed scalar.

- Base Case: $k = 2$, this can be proved by using result of (b).
- Inductive Case: assume that (I.H.)

$$H_{k-1} \cdots H_1 H_0 = I - U_{(k-1)} L_{(k-1)}^{-1} U_{(k-1)}^H \quad (39)$$

show that

$$H_k H_{k-1} \cdots H_1 H_0 = I - U_{(k)} L_{(k)}^{-1} U_{(k)}^H \quad (40)$$

holds for lower triangular matrix $L_{(k)}$.

Proof. With application of the result of (c), we have

$$H_k(H_{k-1} H_{k-2} \cdots H_1 H_0) = \left(1 - \frac{1}{\tau_k} u_k u_k^T\right) (I - U_{(k-1)} L_{(k-1)}^{-1} U_{(k-1)}^H) \quad (41)$$

$$= I - \underbrace{(U_{(k-1)} | u_k)}_{U_{(k)}} \underbrace{\left(\begin{array}{c|c} L_{(k-1)} & 0 \\ \hline u_k^T U_{(k-1)} & \tau_k \end{array} \right)^{-1}}_{L_{(k)}^{-1}} \underbrace{(U_{(k-1)} | u_k)^T}_{U_{(k)}^T} \quad (42)$$

$$= I - U_{(k)} L_{(k)}^{-1} U_{(k)}^T \quad (43)$$

\square

- By the principle of mathematical induction, it can be concluded that

$$H_{k-1} \cdots H_1 H_0 = I - UL^{-1}U^H \quad (44)$$

holds for all $k > 0$.

Exercise 3.**(a) Relationship between m , n and r .**

$$\text{rank}(A) = r \leq \min(m, n) \quad (45)$$

(b) Reduced SVD in terms of U , V and Σ .

- $A = U_r \Sigma_r V_r^T$, where $U_r = U_L$, $\Sigma_r = \Sigma_{TL}$ and $V_r = V_L$.

$$A = U \Sigma V^T = (U_L | U_R) \left(\begin{array}{c|c} \Sigma_{TL} & 0 \\ \hline 0 & 0 \end{array} \right) (V_L | V_R)^T = U_L \Sigma_{TL} V_L^T \quad (46)$$

- $(A^T A)^{-1} = U_r \Sigma_r V_r^T$, where $U_r = V_L^{-T}$, $\Sigma_r = \Sigma_{TL}^{-1} \Sigma_{TL}^{-1}$ and $V_r = V_L^{-T}$.

$$(A^T A)^{-1} = ((U \Sigma V^T)^T U \Sigma V^T)^{-1} \quad (47)$$

$$= (V \Sigma^T U^T U \Sigma V^T)^{-1} \quad (48)$$

$$= (V \Sigma \Sigma V^T)^{-1} \quad (49)$$

$$= \left((V_L | V_R) \left(\begin{array}{c|c} \Sigma_{TL} & 0 \\ \hline 0 & 0 \end{array} \right) \left(\begin{array}{c|c} \Sigma_{TL} & 0 \\ \hline 0 & 0 \end{array} \right) (V_L | V_R)^T \right)^{-1} \quad (50)$$

$$= (V_L \Sigma_{TL} \Sigma_{TL}^T V_L^T)^{-1} \quad (51)$$

$$= V_L^{-T} \Sigma_{TL}^{-1} \Sigma_{TL}^{-1} V_L^{-1} \quad (52)$$

- $(A^T A)^{-1} A^T = U_r \Sigma_r V_r^T$, where $U_r = V_L^{-T}$, $\Sigma_r = \Sigma_{TL}^{-1}$ and $V_r = U_L$.

$$(A^T A)^{-1} A^T = V_L^{-T} \Sigma_{TL}^{-1} \Sigma_{TL}^{-1} V_L^{-1} V_L \Sigma_{TL}^T U_L^T \quad (53)$$

$$= V_L^{-T} \Sigma_{TL}^{-1} \Sigma_{TL}^{-1} \Sigma_{TL}^T U_L^T \quad (54)$$

$$= V_L^{-T} \Sigma_{TL}^{-1} U_L^T \quad (55)$$

- $A(A^T A)^{-1} = U_r \Sigma_r V_r^T$, where $U_r = U_L$, $\Sigma_r = \Sigma_{TL}^{-1} \Sigma_{TL}^{-1}$ and $V_r = V_L^{-T}$.

$$A(A^T A)^{-1} = U_L \Sigma_{TL} V_L^T V_L^{-T} \Sigma_{TL}^{-1} \Sigma_{TL}^{-1} V_L^{-1} \quad (56)$$

$$= U_L \Sigma_{TL} \Sigma_{TL}^{-1} \Sigma_{TL}^{-1} V_L^{-1} \quad (57)$$

$$= U_L \Sigma_{TL}^{-1} V_L^{-1} \quad (58)$$

- $A(A^T A)^{-1} A^T = U_r \Sigma_r V_r^T$, where $U_r = I^{m \times r}$, $\Sigma_r = I^{r \times r}$ and $V_r = I^{n \times r}$.

$$A(A^T A)^{-1} A^T = (U_L \Sigma_{TL}^{-1} V_L^{-1}) V_L \Sigma_{TL}^T U_L^T \quad (59)$$

$$= U_L \Sigma_{TL}^{-1} \Sigma_{TL}^T U_L^T \quad (60)$$

$$= U_L U_L^T \quad (61)$$

$$= I \quad (62)$$

Exercise 4.**(a)** $\psi_1 = (v_{11} - \delta v_{11})\check{\chi}_1$ **where** δv_{11} **is small.***Proof.*

$$\psi_1 = [v_{11}\check{\chi}_1]_* = \frac{v_{11}\check{\chi}_1}{1 + \epsilon_*} = (1 - \frac{\epsilon_*}{1 + \epsilon_*})v_{11}\check{\chi}_1 = (v_{11} - \frac{\epsilon_* v_{11}}{1 + \epsilon_*})\check{\chi}_1 \equiv (v_{11} + \delta v_{11})\check{\chi}_1 \quad (63)$$

where

$$\delta v_{11} = -\frac{\epsilon_* v_{11}}{1 + \epsilon_*}, \text{ where } |\epsilon_*| \leq \mathbf{u} \quad (64)$$

Then

$$|\delta v_{11}| = |-\frac{\epsilon_*}{1 + \epsilon_*}v_{11}| \quad (65)$$

$$= \frac{|\epsilon_*|}{|1 + \epsilon_*|}|v_{11}| \quad (66)$$

$$\leq \frac{\mathbf{u}}{|1 + \epsilon_*|}|v_{11}| \quad (67)$$

$$\leq \frac{\mathbf{u}}{1 + |\epsilon_*|}|v_{11}| \quad (68)$$

$$\leq \frac{\mathbf{u}}{1 + |\epsilon_*| - |\epsilon_*| - \mathbf{u}}|v_{11}| \quad (69)$$

$$= \frac{\mathbf{u}}{1 - \mathbf{u}}|v_{11}| = \gamma_1|v_{11}| \quad (70)$$

Hence, we proved that

$$\psi_1 = (v_{11} - \delta v_{11})\check{\chi}_1 \text{ where } |\delta v_{11}| = \gamma_1|v_{11}| \quad (71)$$

□

(b)*Proof.* Start from dot product. We directly use the corollary in notes.

$$d \triangleq [u_{12}^T \check{x}_2] = (u_{12}^T + \delta u_{12}^T)\check{x}_2, \text{ where } \delta u_{12}^T \leq \gamma_k|u_{12}^T| \leq \gamma_{k+1}|u_{12}^T| \quad (72)$$

Then we consider the subtraction step with Standard Computational Model (SCM).

$$s \triangleq [\psi_1 - d] = (\psi_1 - d)(1 + \epsilon_-) = (\psi_1 - d) + (\psi_1 - d)\epsilon_- \equiv (\psi_1 - d) + \delta\psi_1 = (\psi_1 + \delta\psi_1) - d \quad (73)$$

where $|\delta\psi_1| = |(\psi_1 - d)\epsilon_-| \leq |\psi_1\epsilon_-| = |\epsilon_-||\psi_1| \leq \mathbf{u}|\psi_1|$. Then we consider the division step with Alternative Computational Model (ACM).

$$\check{\chi}_1 = [\frac{s}{v_{11}}] = \frac{s}{v_{11}}(1 + \epsilon_+)^{-1} = \frac{s}{v_{11}(1 + \epsilon_+)} \quad (74)$$

which can be rewritten as

$$(v_{11} + \delta v_{11})\check{\chi}_1 = s \quad (75)$$

where $|\delta v_{11}| = |\epsilon_+ v_{11}| \leq |\epsilon_+| \cdot |v_{11}| \leq \mathbf{u}|v_{11}|$. Then we have

$$(v_{11} + \delta v_{11})\check{\chi}_1 = (\psi_1 + \delta\psi_1) - d = (\psi_1 + \delta\psi_1) - (u_{12}^T + \delta u_{12}^T)\check{x}_2 \quad (76)$$

where

$$|\delta v_{11}| \leq \mathbf{u}|v_{11}|, \quad |\delta\psi_1| \leq \mathbf{u}|\psi_1|, \quad |\delta u_{12}^T| \leq \gamma_{k+1}|u_{12}^T| \quad (77)$$

□

(c) Induction for the desired result

To prove the desired result, let us do mathematical induction on k (the iteration of algorithm).

- Base Case: $k = 0$

Proof. In this case, $U^{0 \times 0}$ does not participate in computation, v_{11} is the right-bottom-most element, and no pre-computed \tilde{x}_2 . Hence, it is natural that

$$(v_{11} + \delta v_{11})\tilde{\chi}_1 = (\psi_1 + \delta\psi_1) \quad (78)$$

which indicates that $(U + \Delta U)\tilde{x} = (y + \delta y)$ holds for $k = 0$. (first iteration) \square

- Inductive case: Assume that at p -th iteration ($p > 0$), pre-computed $\tilde{x}_{2(p)}$ satisfy

$$(U_{(p)} + \Delta U_{(p)})\tilde{x}_{2(p)} = (y_{2(p)} + \Delta y_{2(p)}) \quad (79)$$

where $|\Delta U_{(p)}| \leq \gamma_n |U_{(p)}|$ and $|\delta y_{2(p)}| \leq \mathbf{u} |y_{2(p)}|$, (I.H.) holds.

Show that for the pre-computed \tilde{x}_2 at next iteration $\tilde{x}_{2(p+1)} = \left(\frac{\tilde{\chi}_{1(p)}}{\tilde{x}_{2(p)}} \right)$ and $y_{2(p+1)} = \left(\frac{\psi_{1(p)}}{y_{2(p)}} \right)$

$$(U_{(p+1)} + \Delta U_{(p+1)})\tilde{x}_{2(p+1)} = (y_{(p+1)} + \Delta y_{(p+1)}) \quad (80)$$

where $|\Delta U_{(p+1)}| \leq \gamma_n |U_{(p+1)}|$ and $|\delta y_{2(p+1)}| \leq \mathbf{u} |y_{2(p+1)}|$, holds for some upper triangle $U_{(p+1)}$.

Proof. In terms of result of (b), we have

$$(v_{11(p)} + \delta v_{11(p)})\tilde{\chi}_{1(p)} + (u_{12(p)}^T + \delta u_{12(p)}^T)\tilde{x}_{2(p)} = (\psi_{1(p)} + \delta\psi_{1(p)}) \quad (81)$$

where $\delta v_{11(p)} \leq \mathbf{u} |v_{11(p)}|$, $\delta\psi_{1(p)} \leq \mathbf{u} |\psi_{1(p)}|$, and $\delta u_{12(p)}^T \leq \gamma_{p+1} |u_{12(p)}^T|$.
Combined with (I.H.)

$$(U_{(p)} + \Delta U_{(p)})\tilde{x}_{2(p)} = (y_{2(p)} + \Delta y_{2(p)}) \quad (82)$$

we have

$$\left(\frac{v_{11(p)} + \delta v_{11(p)}}{0} \middle| \frac{u_{12(p)}^T + \delta u_{12(p)}^T}{U_{(p)} + \Delta U_{(p)}} \right) \left(\frac{\tilde{\chi}_{1(p)}}{\tilde{x}_{2(p)}} \right) = \left(\frac{\psi_{1(p)} + \delta\psi_{1(p)}}{y_{2(p)} + \Delta y_{2(p)}} \right) \quad (83)$$

where $|\Delta U_{(p)}| \leq \gamma_n |U_{(p)}|$, $\delta v_{11(p)} \leq \mathbf{u} |v_{11(p)}| \leq \gamma_n |v_{11(p)}|$, $\delta u_{12(p)}^T \leq \gamma_{p+1} |u_{12(p)}^T| \leq \gamma_n |u_{12(p)}^T|$
and $\delta\psi_{1(p)} \leq \mathbf{u} |\psi_{1(p)}|$, $|\delta y_{2(p)}| \leq \mathbf{u} |y_{2(p)}|$. (Note that $0 \leq p \leq n-1$)
which is equivalent to

$$\left(\underbrace{\left(\frac{v_{11(p)}}{0} \middle| \frac{u_{12(p)}^T}{U_{(p)}} \right)}_{U_{(p+1)}} + \underbrace{\left(\frac{\delta v_{11(p)}}{0} \middle| \frac{\delta u_{12(p)}^T}{\Delta U_{(p)}} \right)}_{\Delta U_{(p+1)}} \right) \underbrace{\left(\frac{\tilde{\chi}_{1(p)}}{\tilde{x}_{2(p)}} \right)}_{\tilde{x}_{2(p+1)}} = \left(\underbrace{\left(\frac{\psi_{1(p)}}{y_{2(p)}} \right)}_{y_{2(p+1)}} + \underbrace{\left(\frac{\delta\psi_{1(p)}}{\Delta y_{2(p)}} \right)}_{\delta y_{2(p+1)}} \right) \quad (84)$$

$$(U_{(p+1)} + \Delta U_{(p+1)})\tilde{x}_{2(p+1)} = (y_{2(p+1)} + \delta y_{2(p+1)}) \quad (85)$$

where $|\Delta U_{(p+1)}| \leq \gamma_n |U_{(p+1)}|$ and $|\delta y_{2(p+1)}| \leq \mathbf{u} |y_{2(p+1)}|$.
Hence, we successfully proved (80). \square

- By principle of mathematical induction, it is proved w.l.o.g. that

$$(U + \Delta U)\tilde{x} = (y + \delta y), \text{ where } |\Delta U| \leq \gamma_n |U| \text{ and } |\delta y| \leq \mathbf{u} |y| \quad (86)$$