



THE UNIVERSITY OF TEXAS
AT AUSTIN

EE381V LARGE SCALE OPTIMIZATION

Problem Set 1

Edited by L^AT_EX

Department of Computer Science

STUDENT

Jimmy Lin

xl5224

COURSE COORDINATOR

Sujay Sanghavi

UNIQUE NUMBER

17350

RELEASE DATE

September 12, 2014

DUE DATE

September 18, 2014

TIME SPENT

10 hours

September 15, 2014

Table of Contents

I	Matlab and Computational Assignment	2
1	Gradient Descent on three matrices	2
1.1	$X1, b1$	2
1.2	$X2, b2$	3
1.3	$X3, b3$	4
2	$\gamma = 1$ for the second matrix	5
II	Written Problems	6
1	Othorognal Subspace	6
(a)	Show that if U is a subspace, then so is U^\perp	6
(b)	Show that $(U^\perp)^\perp = U$	6
(c)	Show that if $U, W \subseteq V$ are subspaces of V , then $U \subseteq W \Leftrightarrow U^\perp \supseteq W^\perp$	6
(d)	Show that X^\perp makes sense, X^\perp is a subspace and $(X^\perp)^\perp \supseteq X$	7
(e)	Show that any $v \in V$ can be written uniquely as $v = u + u^\perp$	7
2	Boyd and Vandenberghe, Ex. 2.10	8
(a)	Show that if $A \in \mathbb{S}_+^n$ then the set C is convex	8
(b)	Show that C_1 is convex if there exists $\lambda \in \mathbb{R}$ such that $(A + \lambda gg^T) \in \mathbb{S}_+^n$	8
3	Boyd and Vandenberghe, Ex. 2.21	9
7	Form a Half-Space	10
8	Exists C such that $CA = B$	11
A	Codes Printout	12
(a)	Gradient Descent Routine	12
(b)	Running Script	13

List of Figures

1	Illustration for gradient descent on $X1$, staring with $b1$ by $\gamma = 0.5$ and 3.0	2
2	Illustration for gradient descent on $X2$, starting with $b2$ by $\gamma = 1.5$ and 3.0	3
3	Illustration for gradient descent on $X3$ staring with $b3$ by $\gamma = 0.005$ and 0.05	4
4	Plotting figure for gradient descent with $\gamma = 1$ on the second matrix	5

Part I

Matlab and Computational Assignment

1 Gradient Descent on three matrices

Command to get executed:

```
>> gd_run_script()
```

1.1 $X1, b1$

- Range of γ that leads to convergence: $(0, 2)$
- Range of γ that leads to divergence: $(2, +\infty)$
- Explanation: if $\gamma = 2$, the program indicates that

$$\forall k, f(x^{k+1}) = f(x^k)$$

Since the above equation is constantly true (independent of the minima), we can conclude that gradient descent with $\gamma = 2$ goes to the opposite side of that quadratic curve. Intuitively, the program will diverge if we set larger ($\gamma > 2$) and converge if we set smaller ($\gamma < 2$).

- Two illustrative examples: $\gamma = 0.5$ and $\gamma = 3.0$




Figure 1: Illustration for gradient descent on $X1$, starting with $b1$ by $\gamma = 0.5$ and 3.0

1.2 $X2, b2$

- Range of γ that leads to convergence: $(0, 2)$
- Range of γ that leads to divergence: $(2, +\infty)$
- Explanation: if $\gamma = 2$, the program indicates that

$$\forall k, f(x^{k+1}) = f(x^k)$$

Since the above equation is constantly true (independent of the minima), we can conclude that gradient descent with $\gamma = 2$ goes to the opposite side of that quadratic curve. Intuitively, the program will diverge if we set larger ($\gamma > 2$) and converge if we set smaller ($\gamma < 2$).

- Two illustrative examples: $\gamma = 1.5$ and $\gamma = 3.0$




Figure 2: Illustration for gradient descent on $X2$, starting with $b2$ by $\gamma = 1.5$ and 3.0

1.3 $X3, b3$

- Range of γ that leads to convergence: $(0, 0.02)$
- Range of γ that leads to divergence: $(0.02, +\infty)$
- Explanation: if $\gamma = 0.02$, the program indicates that

$$\forall k, f(x^{k+1}) = f(x^k)$$

Since the above equation is constantly true (independent of the minima), we can conclude that gradient descent with $\gamma = 0.02$ goes to the opposite side of that quadratic curve. Intuitively, the program will diverge if we set larger ($\gamma > 0.02$) and converge if we set smaller ($\gamma < 0.02$).

- Two illustrative examples: $\gamma = 0.005$ and $\gamma = 0.05$




Figure 3: Illustration for gradient descent on $X3$ starting with $b3$ by $\gamma = 0.005$ and 0.05

2 $\gamma = 1$ for the second matrix

Command to get executed:

```
>> gamma = 1;
>> [b2_opt, iters, all_costs] = gd (X2, b2, gamma);
```

Plotting: figure for $\gamma = 1$




Figure 4: Plotting figure for gradient descent with $\gamma = 1$ on the second matrix

Explanation: Through the smooth plotted curve, we guess that the gradient descent method got linear convergence when $\gamma = 1$ on X_2 . Hence, we trace convergence rate $\text{conv_rate} = f(x^k)/f(x^{k-1})$ as follows:

```
Iter: 2, Cost: 2.254428e-01, Conv_Rate: 0.980100
Iter: 3, Cost: 2.209565e-01, Conv_Rate: 0.980100
Iter: 4, Cost: 2.165594e-01, Conv_Rate: 0.980100
Iter: 5, Cost: 2.122499e-01, Conv_Rate: 0.980100
Iter: 6, Cost: 2.080261e-01, Conv_Rate: 0.980100
Iter: 7, Cost: 2.038864e-01, Conv_Rate: 0.980100
...
...
Iter: 381, Cost: 1.107934e-04, Conv_Rate: 0.980100
Iter: 382, Cost: 1.085886e-04, Conv_Rate: 0.980100
Iter: 383, Cost: 1.064277e-04, Conv_Rate: 0.980100
Iter: 384, Cost: 1.043098e-04, Conv_Rate: 0.980100
Iter: 385, Cost: 1.022340e-04, Conv_Rate: 0.980100
Iter: 386, Cost: 1.001996e-04, Conv_Rate: 0.980100
Iter: 387, Cost: 9.820558e-05, Conv_Rate: 0.980100
Converged to zeros!
```

In terms of above dumps and the fact that $f(x^*) = 0$, we can conclude that when $\gamma = 1$

$$f(x^{k+1}) - f(x^*) = 0.9801 \cdot (f(x^k) - f(x^*))$$

which supports our previous guess that

Gradient Descent with $\gamma = 1$ on second matrix leads to **linear convergence**.

Part II

Written Problems

1 Othorognal Subspace

(a) Show that if U is a subspace, then so is U^\perp

Proof. Since U is a subspace, then we have U satisfying all three properties shown below:

- $\mathbf{0} \in U$
- $\forall \mathbf{u}_1, \mathbf{u}_2 \in U, \mathbf{u}_1 + \mathbf{u}_2 \in U$
- $\forall \mathbf{u} \in U, \alpha \in \mathbb{R}, \alpha \mathbf{u} \in U$

Now we show that U^\perp is also a subspace by indicating U^\perp satisfies all three properties as above U do.

- Since $\forall \mathbf{u} \in U, \langle \mathbf{0}, \mathbf{u} \rangle = 0$ and $\mathbf{0} \in V$ ($\mathbf{0} \in U \subseteq V$), then it turned out that $\mathbf{0} \in U^\perp$.
- Let \mathbf{u} be arbitrary vector s.t. $\mathbf{u} \in U$, and $\mathbf{x}_1, \mathbf{x}_2$ to be distinct vector s.t. $\mathbf{x}_1 \in U^\perp$ and $\mathbf{x}_2 \in U^\perp$. By definition of U^\perp , we have $\langle \mathbf{x}_1, \mathbf{u} \rangle = 0$ and $\langle \mathbf{x}_2, \mathbf{u} \rangle = 0$. Then it is obvious that $\langle \mathbf{x}_1 + \mathbf{x}_2, \mathbf{u} \rangle = 0$. That is $\mathbf{x}_1 + \mathbf{x}_2 \in U^\perp$. Therefore, $\forall \mathbf{x}_1, \mathbf{x}_2 \in U^\perp, \mathbf{x}_1 + \mathbf{x}_2 \in U^\perp$ was proved.
- Let \mathbf{x} be arbitrary vector s.t. $\mathbf{x} \in U^\perp$, \mathbf{u} be arbitrary vector s.t. $\mathbf{u} \in U$ and arbitrary $\alpha \in \mathbb{R}$. By definition of U^\perp , we have $\langle \mathbf{x}_1, \mathbf{u} \rangle = 0$. Since inner product is linear operator, it is obvious that $\langle \alpha \mathbf{x}_1, \mathbf{u} \rangle = 0$. That is $\alpha \mathbf{x}_1 \in U^\perp$. Therefore, $\forall \mathbf{x} \in U^\perp, \alpha \in \mathbb{R}, \alpha \mathbf{x} \in U^\perp$ was proved.

Since U^\perp contains $\mathbf{0}$, and is closed under addition and scalar multiplication, it turned out that U^\perp is a subspace. Therefore, the statement that if U is a subspace, then so is U^\perp was proved. \square

(b) Show that $(U^\perp)^\perp = U$

Proof. By contradiction. Assume that $(U^\perp)^\perp \neq U$ and then show the contradiction. By definition of U^\perp , we have $U^\perp = \{\mathbf{v} \in V, \langle \mathbf{v}, \mathbf{u} \rangle = 0, \forall \mathbf{u} \in U\}$ and $(U^\perp)^\perp = \{\mathbf{x} \in V, \langle \mathbf{x}, \mathbf{v} \rangle = 0, \forall \mathbf{v} \in U^\perp\}$. Since $(U^\perp)^\perp \neq U$, then we can say that $\exists \mathbf{x} \notin U, \langle \mathbf{x}, \mathbf{v} \rangle = 0, \forall \mathbf{v} \in U^\perp$. That is to say, $\exists \mathbf{x} \in V$ but $\notin U, \langle \mathbf{x}, \mathbf{v} \rangle = 0, \forall \mathbf{v} \in U^\perp$ s.t. $\langle \mathbf{v}, \mathbf{u} \rangle = 0, \forall \mathbf{u} \in U$. However, such \mathbf{x} does not exist. Hence, we reject initial assumption and conclude that $(U^\perp)^\perp = U$. \square

(c) Show that if $U, W \subseteq V$ are subspaces of V , then $U \subseteq W \Leftrightarrow U^\perp \supseteq W^\perp$

Proof of $U \subseteq W \Rightarrow U^\perp \supseteq W^\perp$. $U^\perp = \{\mathbf{v} \in V, \langle \mathbf{v}, \mathbf{u} \rangle = 0, \forall \mathbf{u} \in U\}$ and $W^\perp = \{\mathbf{v} \in V, \langle \mathbf{v}, \mathbf{w} \rangle = 0, \forall \mathbf{w} \in W\} = \{\mathbf{v} \in V, \langle \mathbf{v}, \mathbf{w} \rangle = 0, \forall \mathbf{w} \in W \text{ and } \langle \mathbf{v}, \mathbf{u} \rangle = 0, \forall \mathbf{u} \in U\}$ (This is valid because $\forall U \subseteq W$ and then $\mathbf{u} \in W$). Now since membership of W^\perp requires one more condition, then it is obvious that $\mathbf{v} \in W^\perp \Rightarrow \mathbf{v} \in U^\perp$, and $\mathbf{v} \in U^\perp \not\Rightarrow \mathbf{v} \in W^\perp$ hold for arbitrary \mathbf{v} . Therefore, we can conclude that $U^\perp \supseteq W^\perp$. \square

Proof of $U^\perp \supseteq W^\perp \Rightarrow U \subseteq W$. By definition, we have $U^\perp = \{\mathbf{v} \in V | \langle \mathbf{v}, \mathbf{u} \rangle = 0, \forall \mathbf{u} \in U\}$ and $W^\perp = \{\mathbf{v} \in V | \langle \mathbf{v}, \mathbf{w} \rangle = 0, \forall \mathbf{w} \in W\}$. Since $U^\perp \supseteq W^\perp$, then we have $U^\perp \cap W^\perp = W^\perp$. Then $U^\perp \cap W^\perp = \{\mathbf{v} \in V | \langle \mathbf{v}, \mathbf{u} \rangle = 0, \forall \mathbf{u} \in U \text{ and } \langle \mathbf{v}, \mathbf{w} \rangle = 0, \forall \mathbf{w} \in W\} = \{\mathbf{v} \in V | \langle \mathbf{v}, \mathbf{x} \rangle = 0, \forall \mathbf{x} \in W \cup U\} = W^\perp$. Then we have $W \cup U = W$, which naturally derives $U \subseteq W$. \square

- (d) Show that X^\perp makes sense, X^\perp is a subspace and $(X^\perp)^\perp \supseteq X$
- (e) Show that any $v \in V$ can be written uniquely as $v = u + u^\perp$

2 Boyd and Vandenberghe, Ex. 2.10

- (a) Show that if $A \in \mathbb{S}_+^n$ then the set \mathbf{C} is convex
- (b) Show that C_1 is convex if there exists $\lambda \in \mathbb{R}$ such that $(A + \lambda gg^T) \in \mathbb{S}_+^n$

3 Boyd and Vandenberghe, Ex. 2.21

7 Form a Half-Space

8 Exists C such that $CA = B$

A Codes Printout

(a) Gradient Descent Routine

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%% Problem set 1: Standard Gradient Descent with fixed step size
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
% Usage:
%   [b, iter, all_costs] = gd (X, b_init, gamma)
% Parameters:
%   X: matrix in quadratic optimization
%   b_init: starting vector of variable b
%   gamma: fixed step size
% Note that stopping criteria is set by absolute eps = 10e-5.

function [b, iter, all_costs] = gd (X, b_init, gamma)
eps = 10e-5;
b = b_init;
last_cost = 0.5 * b' * X * b;
iter = 1;
all_costs = [];
while true,
    %% compute essential numerics and do gradient descent
    gradient = X * b;
    b = b - gamma * gradient;
    cost = 0.5 * b' * X * b;
    rate = (cost / last_cost);
    all_costs = [all_costs cost];
    %% output numeric information of this iteration
    disp(sprintf('Iter: %d, Cost: %e, Conv.Rate: %f', iter, cost, rate));
    %% quadratic optimization converges to zero
    if cost < eps,
        disp('Converged to zeros!')
        break
    end
    %% quadratic optimization diverges
    if cost >= last_cost && iter > 10,
        disp('Problem diverges!')
        break
    end
    %% prepare for next iteration
    last_cost = cost;
    iter = iter + 1;
end

%% uncomment following code for plotting individual gradient descent run
%% plot f(b^(n)) with regard to n
%% plot (1:iter, all_costs)
%% title (sprintf ('Gradient Descent with gamma=%f', gamma))
%% xlabel ('iteration: n')
%% ylabel ('cost of function: f(b^n)')

end

```

(b) Running Script

```

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%% Running scripts for applying gradient descent
%% on three given dataset
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%% for X1, b1
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
gamma1_one = 0.5; gamma2_two = 3;
[b1_opt_one, iter1_one, costs1_one] = Gradient_Descent(X1, b1, gamma1_one);
[b1_opt_two, iter1_two, costs1_two] = Gradient_Descent(X1, b1, gamma2_two);
subplot (1, 2, 1)
plot (1:iter1_one, costs1_one)
axis ([-0.1*iter1_one 1.1*iter1_one -0.05*max(costs1_one) max(costs1_one)])
title (sprintf ('Gradient Descent with gamma=%f', gamma1_one))
xlabel ('iteration: n')
ylabel ('cost of function: f(b^n)')
subplot (1, 2, 2)
plot (1:iter1_two, costs1_two)
axis ([-0.1*iter1_two 1.1*iter1_two -0.05*max(costs1_two) max(costs1_two)])
title (sprintf ('Gradient Descent with gamma=%f', gamma2_two))
xlabel ('iteration: n')
ylabel ('cost of function: f(b^n)')

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%% for X2, b2
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
gamma2_one = 1.5; gamma2_two = 3;
[b2_opt_one, iter2_one, costs2_one] = Gradient_Descent(X2, b2, gamma2_one);
[b2_opt_two, iter2_two, costs2_two] = Gradient_Descent(X2, b2, gamma2_two);
figure()
subplot (1, 2, 1)
plot (1:iter2_one, costs2_one)
axis ([-0.1*iter2_one 1.1*iter2_one -0.05*max(costs2_one) max(costs2_one)])
title (sprintf ('Gradient Descent with gamma=%f', gamma2_one))
xlabel ('iteration: n')
ylabel ('cost of function: f(b^n)')
subplot (1, 2, 2)
plot (1:iter2_two, costs2_two)
axis ([-0.1*iter2_two 1.1*iter2_two -0.05*max(costs2_two) max(costs2_two)])
title (sprintf ('Gradient Descent with gamma=%f', gamma2_two))
xlabel ('iteration: n')
ylabel ('cost of function: f(b^n)')

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%% for X3, b3
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
gamma3_one = 0.005; gamma3_two = 0.05;
[b3_opt_one, iter3_one, costs3_one] = Gradient_Descent(X3, b3, gamma3_one);
[b3_opt_two, iter3_two, costs3_two] = Gradient_Descent(X3, b3, gamma3_two);
figure()
subplot (1, 2, 1)
plot (1:iter3_one, costs3_one)
axis ([-0.1*iter3_one 1.1*iter3_one -0.05*max(costs3_one) max(costs3_one)])
title (sprintf ('Gradient Descent with gamma=%f', gamma3_one))
xlabel ('iteration: n')
ylabel ('cost of function: f(b^n)')
subplot (1, 2, 2)
plot (1:iter3_two, costs3_two)
axis ([-0.1*iter3_two 1.1*iter3_two -0.05*max(costs3_two) max(costs3_two)])
title (sprintf ('Gradient Descent with gamma=%f', gamma3_two))
xlabel ('iteration: n')
ylabel ('cost of function: f(b^n)')

```