

# Class 9

AUTHOR

Charlie Rezanka (A15837296)

## PBD Statistics

```
data <- read.csv("Data Export Summary.csv", row.names=1)
head(data)
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	154,766	10,155	12,187	191	72	32
Protein/Oligosaccharide	9,083	1,802	32	7	1	0
Protein/NA	8,110	3,176	283	6	0	0
Nucleic acid (only)	2,664	94	1,450	12	2	1
Other	163	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	177,403					
Protein/Oligosaccharide	10,925					
Protein/NA	11,575					
Nucleic acid (only)	4,223					
Other	204					
Oligosaccharide (only)	22					

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

```
#create a function with X as an input
sum_comma <- function(data) {
  #Substitute comma and convert to numeric
  sum(as.numeric(gsub(",", "", data)))
}
```

For X Ray

```
round(sum_comma(data$X.ray)/sum_comma(data$Total), 3)*100
```

```
[1] 85.5
```

For EM

```
round(sum_comma(data$EM)/sum_comma(data$Total), 3)*100
```


```
[1] 7.5
```

Q2: What proportion of structures in the PDB are protein?

```
round(sum_comma(data$Total[1])/sum_comma(data$Total), 3)*100
```

[1] 86.8

Q3: Insert your HIV image.



HIV-PR structure from MERK with critical residues highlighted. Protein structure is color coded to residue.

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

Hydrogen molecules are too small and abundant to be efficiently displayed

Q5: There is a critical "conserved" water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have

The water molecule binds to to ASP25 on each chain

#Working with Structures in R

```
library(bio3d)
pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
```

```
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNFP
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40
	segid	elesy	charge										
1	<NA>	N	<NA>										
2	<NA>	C	<NA>										
3	<NA>	C	<NA>										

4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

Q7: How many amino acid residues are there in this pdb object?

There are 190 residues.

Q8: Name one of the two non-protein residues?

Water (shown as HOH)

Q9: How many protein chains are in this structure?

There are 2 chains.

## Predicting Movements of a Structure

First, read the ADK structure

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```
MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLAAVKSGSELGKQAKDIMDAGKLV  
DELVIALVKERIAQEDCRNGFLLDGFPRTIPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```


```
+ attr: atom, xyz, seqres, helix, sheet,  
       calpha, remark, call
```

Perform a prediction of flexibility with a technique called NMA (normal mode analysis), which is meant to predict flexibility and motion.

```
m <- nma(adk)
```

```
Building Hessian...      Done in 0.1 seconds.  
Diagonalizing Hessian... Done in 0.81 seconds.
```

```
plot(m)
```



Write out a "movie" (trajectory) of the motion for viewing in MOLstar

```
mktrj(m, file="adk_m7.pdb")
```