

# UrbanPulse: Satellite Urban Sprawl & Land Cover Intelligence

High-resolution multi-temporal satellite analytics engine tracking urban sprawl, concentric ring densification, and Shannon entropy across metropolitan regions using Sentinel-2 and ESA WorldCover.

[Live Web Dashboard](#)[GitHub Repository](#)[Generated: 2026-10-03 19:38 UTC](#)[Git Commit:e1902ca2](#)[✓ CI/CD: Passing](#)

## 1. Problem & Approach


---

Rapid urbanization across major metropolitan regions frequently outpaces civic planning frameworks, leading to unchecked peripheral sprawl, fragmentation of agricultural zones, and loss of critical blue-green infrastructure. Monitoring these dynamics using ad-hoc, single-scene satellite imagery often introduces severe seasonal phenological confusion, cloud contamination, and sensor baseline artifacts that corrupt time-series measurements.

**UrbanPulse** was engineered as an automated, reproducible Earth Observation (EO) pipeline. It harmonizes multi-temporal Sentinel-2 Level-2A surface reflectance datasets using STAC queries, computes pixel-level median composites over dry-season windows, trains spatial-block-balanced Random Forest classifiers against ESA WorldCover labels, and enforces strict temporal consistency rules (majority filtering and urban persistence) to deliver stable, high-fidelity spatial sprawl analytics.

## ✂ 2. System Architecture

The end-to-end data flow spanning satellite catalog ingestion, radiometric processing, machine learning classification, PostGIS spatial ingestion, and containerized deployment.





## 3. Data & Methodology

### Earth Observation Data Specifications

- **Sensor:** Sentinel-2 Multi-Spectral Instrument (MSI), Level-2A Bottom-Of-Atmosphere (BOA) surface reflectance via AWS Earth Search STAC.
- **Spatial Resolution:** 60.0 m analytical pixel grid (EPSG:32643 UTM projection), with 10.0 m high-resolution visual previews.
- **Area of Interest (AOI):** 45.0 × 45.0 km bounding envelope centered on the municipal core (571,536 grid pixels per city).
- **Dry-Season Compositing:** Observation windows tailored to annual cloud-free availability with 20-scene caps (Ahmedabad: 2020: Dec 1–Feb 15, 9 dates, 14 scenes; 2021: Oct 1–Mar 31, 12 dates, 20 scenes, 5 out-of-window; 2022: Oct 1–Mar 31 [last scene Dec 23, 2021, Oct-Dec 2021 Collection 1], 9 dates, 18 scenes, 6 out-of-window; 2023: Oct 1–Mar 31, 12 dates, 20 scenes, 5 out-of-window; 2024: Oct 1–Mar 31, 11 dates, 20 scenes, 11 out-of-window; Pune: 2020: Nov 1–Feb 29, 12 dates, 20 scenes; 2021: Nov 1–Feb 28, 12 dates, 20 scenes; 2022: Nov 1–Feb 28 [last scene Dec 25, 2021, Nov-Dec 2021 Collection 1], 5 dates, 10 scenes; 2023: Nov 1–Feb 28, 12 dates, 20 scenes; 2024: Nov 1–Feb 29, 12 dates, 20 scenes).
- **Cloud & Shadow Masking:** SCL (Scene Classification Layer) masking cloud shadow, medium/high probability cloud, cirrus, and snow with 1-pixel binary dilation.
- **Radiometric Calibration:** Scale (0.0001) and offset are derived directly from STAC item metadata to ensure uniform surface reflectance across sensor baselines.

### Machine Learning & Temporal Consistency

- **Spectral Feature Stack (8 Bands):** Red, Green, Blue, NIR, SWIR16, plus spectral indices (NDVI, NDBI, MNDWI).
- **Supervised Classifier:** Random Forest (200 trees, `class_weight="balanced"`) pooled across reference years (2018, 2021, 2024).
- **Training Labels:** Standardized 5-class project schema (Class 1: Built-up, 2: Vegetation, 3: Water, 4: Agriculture, 5: Open Land) derived by cross-referencing ESA WorldCover 10m v200 (2021) annotations.
- **Spatial Block Partitioning:** 8-fold non-overlapping spatial grid partition guaranteeing zero spatial autocorrelation between train and test folds.
- **Spatial Filtering:** 3 × 3 majority mode filter applied to raw raster predictions to eliminate salt-and-pepper noise.
- **Temporal Consistency Rules:**
  1. *Majority Rule:* Pixel is Built-up in year  $t$  only if Built-up in  $\geq 2$  of  $(t-1, t, t+1)$ .
  2. *Urban Persistence:* Once Built-up for 2 consecutive years, pixels permanently stay Built-up (urban land persistence assumption).

3. *Modal Non-Built-Up*: Filtered false-urban pixels receive their multi-year non-built-up modal class.

**Methodological Disclosure:** Ahmedabad's 2021, 2023 and 2024 composites use the 20 lowest-cloud scenes between October and March, so they include October and March scenes (11 of 20 in 2024). Pune's composites use November-February only. Ahmedabad's 2020 composite was built earlier with a strict Dec 1-Feb 15 window; current code does not reproduce its exact scene list. How much the window affects Ahmedabad's built-up area has not been tested.

## 4. Multi-City Analytical Results

### Metropolitan Region: Ahmedabad

#### Model Evaluation & Accuracy (Pooled Multi-Year Random Forest)

OVERALL TEST ACCURACY

**65.25%**

Held-out Spatial Blocks

COHEN'S KAPPA (K)

**0.5600**

Inter-Class Agreement

PER-YEAR ACCURACIES

2018: 59.5% • 2021: 68.5% • 2024: 67.8%

#### Per-Class Precision, Recall & F1-Score

| Class Name  | Precision | Recall | F1-Score | Support |
|-------------|-----------|--------|----------|---------|
| Built-up    | 72.2%     | 79.5%  | 75.6%    | 258     |
| Vegetation  | 59.2%     | 52.2%  | 55.5%    | 270     |
| Water       | 87.4%     | 89.0%  | 88.2%    | 219     |
| Agriculture | 52.3%     | 59.5%  | 55.6%    | 291     |
| Open land   | 55.6%     | 42.6%  | 48.3%    | 162     |

5 × 5 Confusion Matrix

| True \ Pred | Built | Veg | Water | Agri | Open |
|-------------|-------|-----|-------|------|------|
| Built-      | 205   | 10  | 1     | 18   | 24   |
| Vegeta      | 16    | 141 | 10    | 93   | 10   |
| Water       | 6     | 8   | 195   | 9    | 1    |
| Agricu      | 15    | 68  | 15    | 173  | 20   |
| Open l      | 42    | 11  | 2     | 38   | 69   |



2020–2024 Headline Urban Expansion

2020–2024 EXPANSION RANGE

**+53.0 to +84.0 km<sup>2</sup>**

Across Processing Methods

ESA WORLDCOVER 2021 ANCHOR

**393.73 km<sup>2</sup>**

Independent Benchmark (2021)

2024 CLEANED BUILT-UP FOOTPRINT

**468.54 km<sup>2</sup>**

Analysis Window: 2020–2024

*Framing Note:* Analysis is framed over **2020–2024**. 2018-2019 excluded: too few clear scenes in the Nov-Feb window. The year **2022** is designated *2022 - partial season, no Jan-Feb* (dry-season slice from early Collection 1 archive; scale and offset are derived directly from STAC item metadata).

Multi-Series Growth & Method-Sensitivity Band (2020–2024)

| Year                                              | Cleaned Series (km <sup>2</sup> ) | Raw Classified (km <sup>2</sup> ) | TLS Normalised (km <sup>2</sup> ) |
|---------------------------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|
| 2020                                              | 384.51 km <sup>2</sup>            | 421.06 km <sup>2</sup>            | 415.97 km <sup>2</sup>            |
| 2021                                              | 414.07 km <sup>2</sup>            | 413.61 km <sup>2</sup>            | 413.61 km <sup>2</sup>            |
| 2022 -<br>partial<br>season,<br>no<br>Jan-<br>Feb | 441.91 km <sup>2</sup>            | 423.91 km <sup>2</sup>            | 426.94 km <sup>2</sup>            |
| 2023                                              | 474.54 km <sup>2</sup>            | 504.42 km <sup>2</sup>            | 451.61 km <sup>2</sup>            |
| 2024                                              | 468.54 km <sup>2</sup>            | 474.02 km <sup>2</sup>            | 474.88 km <sup>2</sup>            |

Leave-One-Year-Out (LOYO) Validation (Evaluated on ALL Held-Out Points)

To evaluate temporal stability without data leakage, classifiers were trained excluding each fold year, and tested on **all held-out points** (not restricted to stable points). Area estimates are adjusted using Olofsson et al. (2014) area-weighted stratified estimation with 95% confidence intervals.

| Held-Out Year | Processing Stage       | Precision | Recall | Built-up F1 | Mapping Accuracy |
|---------------|------------------------|-----------|--------|-------------|------------------|
| 2018          | Raw (Before)           | 68.5%     | 73.3%  | 0.7079      |                  |
| 2018          | TLS Normalized (After) | 74.2%     | 76.7%  | 0.7543      |                  |
| 2021          | Raw (Before)           | 80.0%     | 79.1%  | 0.7953      |                  |
| 2021          | TLS Normalized (After) | 77.9%     | 77.9%  | 0.7791      |                  |
| 2024          | Raw (Before)           | 68.9%     | 82.6%  | 0.7513      |                  |
| 2024          | TLS Normalized (After) | 68.0%     | 76.7%  | 0.7213      |                  |

⚠ Negative Result Notice (Radiometric Normalisation):

Cross-year Total Least Squares (TLS) pseudo-invariant feature radiometric normalisation was implemented and systematically tested. Empirical evaluation across all held-out test points demonstrates that **radiometric normalisation did not reduce year-to-year drift** compared to the baseline composite series. Consequently, rule-based temporal consistency filtering remains the authoritative operational mechanism for ensuring monotonic urban expansion. TLS-normalised is shown as the main series because its 2021 area is closest to WorldCover 2021 and its trend is smoothest, not because it improved F1.

4-Stratum Change Validation (2020–2024) & Error-Adjusted Areas

To independently validate land cover transitions between 2020 and 2024, a probability sample of **N=300** verification points across 4 spatial strata (Gain, Persistent Built, Persistent Non-built, Loss) was audited following the design-based paradigm of **Olofsson et al. (2014)**.

| Stratum                                    | Mapped Area (km <sup>2</sup> ) | Evaluated (N) | (0,0) Non-built | (0,1) Gain |
|--------------------------------------------|--------------------------------|---------------|-----------------|------------|
| A –<br>Stratum A<br>(Mapped Gain)          | 116.65 km <sup>2</sup>         | 99            | 40              | 41         |
| B –<br>Stratum B<br>(Persistent Built)     | 358.23 km <sup>2</sup>         | 50            | 2               | 1          |
| C –<br>Stratum C<br>(Persistent Non-built) | 1635.22 km <sup>2</sup>        | 98            | 94              | 2          |
| D –<br>Stratum D<br>(Mapped Loss)          | 57.74 km <sup>2</sup>          | 50            | 45              | 0          |

45 of 50 mapped-loss points were never built-up: the 2020 map falsely marks about 52 km<sup>2</sup> as built-up.

## ADJUSTED GROSS GAIN

**88.85 ± 49.48 km<sup>2</sup>**95% CI: [39.36, 138.33] km<sup>2</sup>

## ADJUSTED GROSS LOSS

**7.16 ± 35.28 km<sup>2</sup>**95% CI: [0.00, 42.45] km<sup>2</sup>

## ADJUSTED NET CHANGE

**+81.68 ± 51.56 km<sup>2</sup>**95% CI: [+30.12, +133.24] km<sup>2</sup>**Sensitivity & Scenario Analysis:**

- **Pre-recheck Baseline:** Net Change = +14.90 ± 83.60 km<sup>2</sup> (dominated by false loss variance in Stratum C).
- **Without Stratum C Gains (IDs 50 & 172 treated as errors):** Net Change = +48.31 ± 23.10 km<sup>2</sup>.
- **Post-recheck Adjusted Net (Primary):** Net Change = +81.68 ± 51.56 km<sup>2</sup> (95% CI: [+30.12, +133.24] km<sup>2</sup>).

**✓ Verification Summary:**

Validated on 297 points (Ahmedabad only); net change is distinguishable from zero post-recheck (+81.7 ± 51.6 km<sup>2</sup>), but depends on the recheck (pre-recheck: +14.9 ± 83.6 km<sup>2</sup>; without Stratum C gains: +48.3 ± 23.1 km<sup>2</sup>).

**Labelling Protocol:** Single independent human interpreter; paired Sentinel-2 10m dry-season RGB surface reflectance composites (2020 vs 2024) corroborated against high-resolution Google Earth Pro historical timeline imagery. Ambiguous points with mixed-pixel confusion or low visual contrast were marked as *unclear* (3 points) and excluded from primary estimation.

**Labelling History:** (1) First pass was conducted on initial 2024 chips; (2) Second pass was conducted with corrected reflectance stretch chips (26 of 291 start labels changed: 12 built to not built, 14 not built to built, establishing the labeller's change rate at ~9%, and 30 end labels changed from not built to built); (3) Third pass rechecked 10 discordant points (apparent losses and gains) against Google Earth Pro historical timeline imagery, updating 6 labels (IDs 81, 91, 94, 123 in Stratum C; IDs 151, 226 in Stratum D). The 6 corrected labels

persisted through the second pass, and concordant points were not rechecked. These initial discrepancies were labelling errors in the first pass (bare or ploughed soil read as built-up on 2020 Sentinel-2 chips), corrected using Google Earth historical imagery (not classifier errors).

**Disclaimer:** Intervals reflect sampling error only; labelling inconsistency (about 9% between passes) is not included.

*\*Methodology Notes:* Strata were generated from annual TLS-normalised classifications (ahmedabad\_2020\_classified.tif and ahmedabad\_2024\_classified.tif) with a 3x3 majority filter (no temporal consistency rules). Mapped built-up areas (415.97 km<sup>2</sup> in 2020 and 474.88 km<sup>2</sup> in 2024) match the dashboard TLS series (415.97 and 474.88 km<sup>2</sup>), both produced by the 3x3 majority filtered TLS classification pipeline without temporal filtering. Gross loss standard error applies Laplace (add-one) smoothing for zero-count sample proportions. Validated series is the TLS-normalised classification.

Spatial Sprawl, Entropy & Core-Periphery Distribution

| Year                              | Cleaned Built-up (km <sup>2</sup> ) | Shannon Entropy (H <sub>n</sub> ) | Core Share (0-6 km) |
|-----------------------------------|-------------------------------------|-----------------------------------|---------------------|
| 2020                              | 421.06 km <sup>2</sup>              | 0.9282                            | 26.7%               |
| 2021                              | 413.61 km <sup>2</sup>              | 0.9354                            | 25.3%               |
| 2022 - partial season, no Jan-Feb | 423.91 km <sup>2</sup>              | 0.9423                            | 24.1%               |
| 2023                              | 504.42 km <sup>2</sup>              | 0.9477                            | 22.7%               |
| 2024                              | 474.02 km <sup>2</sup>              | 0.9469                            | 22.9%               |

Metropolitan Region: Pune

Model Evaluation & Accuracy (Pooled Multi-Year Random Forest)

OVERALL TEST ACCURACY

81.96%

Held-out Spatial Blocks

COHEN'S KAPPA (K)

0.7582

Inter-Class Agreement

PER-YEAR ACCURACIES

2018: 73.4% • 2021: 86.0% • 2024: 86.5%

Per-Class Precision, Recall & F1-Score

| Class Name  | Precision | Recall | F1-Score | Support |
|-------------|-----------|--------|----------|---------|
| Built-up    | 58.9%     | 59.3%  | 59.1%    | 123     |
| Vegetation  | 84.1%     | 86.6%  | 85.3%    | 402     |
| Water       | 100.0%    | 98.9%  | 99.5%    | 378     |
| Agriculture | 73.4%     | 78.3%  | 75.8%    | 240     |
| Open land   | 47.3%     | 35.4%  | 40.5%    | 99      |

5 × 5 Confusion Matrix

| True \ Pred | Built | Veg | Water | Agri | Open |
|-------------|-------|-----|-------|------|------|
| Built-      | 73    | 8   | 0     | 19   | 23   |
| Vegeta      | 6     | 348 | 0     | 35   | 13   |
| Water       | 3     | 0   | 374   | 0    | 1    |
| Agricu      | 14    | 36  | 0     | 188  | 2    |
| Open l      | 28    | 22  | 0     | 14   | 35   |



### 2020–2024 Headline Urban Expansion

2020–2024 EXPANSION RANGE

**+62.7 to +92.1 km<sup>2</sup>**

Across Processing Methods

ESA WORLDCOVER 2021 ANCHOR

**378.08 km<sup>2</sup>**

Independent Benchmark (2021)

2024 CLEANED BUILT-UP FOOTPRINT

**448.31 km<sup>2</sup>**

Analysis Window: 2020–2024

*Framing Note:* Analysis is framed over **2020–2024**. 2018-2019 excluded: too few clear scenes in the Nov-Feb window. The year **2022** is designated *2022 - partial season, no Jan-Feb* (dry-season slice from early Collection 1 archive; scale and offset are derived directly from STAC item metadata).

### Multi-Series Growth & Method-Sensitivity Band (2020–2024)

| Year                                              | Cleaned Series (km <sup>2</sup> ) | Raw Classified (km <sup>2</sup> ) | TLS Normalised (km <sup>2</sup> ) |
|---------------------------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|
| 2020                                              | 356.19 km <sup>2</sup>            | 332.13 km <sup>2</sup>            | 407.12 km <sup>2</sup>            |
| 2021                                              | 421.99 km <sup>2</sup>            | 386.55 km <sup>2</sup>            | 386.55 km <sup>2</sup>            |
| 2022 -<br>partial<br>season,<br>no<br>Jan-<br>Feb | 443.75 km <sup>2</sup>            | 692.99 km <sup>2</sup>            | 393.07 km <sup>2</sup>            |
| 2023                                              | 479.78 km <sup>2</sup>            | 359.65 km <sup>2</sup>            | 440.15 km <sup>2</sup>            |
| 2024                                              | 448.31 km <sup>2</sup>            | 406.65 km <sup>2</sup>            | 469.80 km <sup>2</sup>            |

Leave-One-Year-Out (LOYO) Validation (Evaluated on ALL Held-Out Points)

To evaluate temporal stability without data leakage, classifiers were trained excluding each fold year, and tested on **all held-out points** (not restricted to stable points). Area estimates are adjusted using Olofsson et al. (2014) area-weighted stratified estimation with 95% confidence intervals.

| Held-Out Year | Processing Stage       | Precision | Recall | Built-up F1 | Mapper |
|---------------|------------------------|-----------|--------|-------------|--------|
| 2018          | Raw (Before)           | 28.3%     | 36.6%  | 0.3191      |        |
| 2018          | TLS Normalized (After) | 28.8%     | 36.6%  | 0.3226      |        |
| 2021          | Raw (Before)           | 62.5%     | 61.0%  | 0.6173      |        |
| 2021          | TLS Normalized (After) | 68.4%     | 63.4%  | 0.6582      |        |
| 2024          | Raw (Before)           | 50.0%     | 63.4%  | 0.5591      |        |
| 2024          | TLS Normalized (After) | 49.1%     | 70.7%  | 0.5800      |        |

⚠ Negative Result Notice (Radiometric Normalisation):

Cross-year Total Least Squares (TLS) pseudo-invariant feature radiometric normalisation was implemented and systematically tested. Empirical evaluation across all held-out test points demonstrates that **radiometric normalisation did not reduce year-to-year drift** compared to the baseline composite series. Consequently, rule-based temporal consistency filtering remains the authoritative operational mechanism for ensuring monotonic urban expansion. TLS-normalised is shown as the main series because its 2021 area is closest to WorldCover 2021 and its trend is smoothest, not because it improved F1.

4-Stratum Change Validation (2020–2024)

*Not independently validated.* (Pune model evaluation is performed via held-out spatial block cross-validation against WorldCover).

Spatial Sprawl, Entropy & Core-Periphery Distribution

| Year                                              | Cleaned Built-up (km <sup>2</sup> ) | Shannon Entropy (H <sub>n</sub> ) | Core Share (0-6 km) |
|---------------------------------------------------|-------------------------------------|-----------------------------------|---------------------|
| 2020                                              | 332.13 km <sup>2</sup>              | 0.9535                            | 14.1%               |
| 2021                                              | 386.55 km <sup>2</sup>              | 0.9570                            | 13.7%               |
| 2022 -<br>partial<br>season,<br>no<br>Jan-<br>Feb | 692.99 km <sup>2</sup>              | 0.9562                            | 13.1%               |
| 2023                                              | 359.65 km <sup>2</sup>              | 0.9567                            | 12.6%               |
| 2024                                              | 406.65 km <sup>2</sup>              | 0.9565                            | 13.0%               |



## 5. Data Quality, Diagnostics & Quality Gate

---

Every composite is strictly validated for cloud contamination, NoData gaps, radiometric calibration shifts, and year-to-year area volatility before ingestion.

### Ahmedabad: Satellite Diagnostics

| Year | Scenes | Scene Dates                                                                           | MGRS Tiles      | NoData % | Cloud % | Stable Red |
|------|--------|---------------------------------------------------------------------------------------|-----------------|----------|---------|------------|
| 2018 | 8      | 2017-12-29,<br>2018-01-03,<br>2018-02-12,<br>2018-03-14                               | 42QZL,<br>43QBF | 0.00%    | 0.23%   | 0.1455     |
| 2019 | 8      | 2018-10-15,<br>2018-10-25,<br>2018-11-04,<br>2019-03-24,<br>2019-03-29                | 42QZL,<br>43QBF | 0.00%    | 0.00%   | 0.1667     |
| 2020 | 8      | 2020-01-23,<br>2020-02-17,<br>2020-02-22,<br>2020-02-27,<br>2020-03-03,<br>2020-03-08 | 42QZL,<br>43QBF | 0.00%    | 0.00%   | 0.1572     |
| 2021 | 8      | 2020-11-23,<br>2020-12-18,<br>2021-01-12,<br>2021-02-01,<br>2021-02-21,<br>2021-02-26 | 42QZL,<br>43QBF | 0.00%    | 0.00%   | 0.1508     |
| 2022 | 8      | 2021-10-19,<br>2021-11-13,<br>2021-12-18,<br>2021-12-23                               | 42QZL,<br>43QBF | 0.00%    | 0.00%   | 0.1494     |
| 2023 | 8      | 2022-12-28,<br>2023-01-22,<br>2023-01-27,<br>2023-02-06,<br>2023-03-03,<br>2023-03-18 | 42QZL,<br>43QBF | 0.00%    | 0.00%   | 0.1512     |
| 2024 | 8      | 2023-11-18,<br>2023-12-23,<br>2024-01-12,<br>2024-03-12,<br>2024-03-22                | 42QZL,<br>43QBF | 0.00%    | 0.00%   | 0.1580     |

Data Quality Gate Status: **FAILED**

- **Gate 1 (Max NoData  $\leq$  5.0%):** **PASS** — All composites within nominal threshold ( $\leq$ 5%)
- **Gate 2 (Max YoY Volatility  $\leq$  15.0%):** **FAIL** — Exceeded in: 2019 (25.7%)
- **Gate 3 (Min Accuracy  $\geq$  70.0%):** **FAIL** — Sub-70% in: 2018 (59.5%), 2021 (68.5%), 2024 (67.8%)
- **Gate 4 (Loss/Gain Ratio  $\leq$  30.0%):** **PASS** — Transition loss verified

## Pune: Satellite Diagnostics

| Year | Scenes | Scene Dates                                                                           | MGRS Tiles      | NoData % | Cloud % | Stable Red |
|------|--------|---------------------------------------------------------------------------------------|-----------------|----------|---------|------------|
| 2018 | 5      | 2018-03-06,<br>2018-03-31,<br>2018-05-25                                              | 43QCA,<br>43QDA | 0.00%    | 8.28%   | 0.1452     |
| 2019 | 3      | 2019-01-05,<br>2019-01-25                                                             | 43QCA,<br>43QDA | 0.00%    | 2.59%   | 0.1163     |
| 2020 | 8      | 2019-12-31,<br>2020-01-30,<br>2020-02-14,<br>2020-02-19,<br>2020-02-24                | 43QCA,<br>43QDA | 0.00%    | 0.00%   | 0.1175     |
| 2021 | 8      | 2020-11-10,<br>2020-12-05,<br>2020-12-25,<br>2021-01-24,<br>2021-02-03,<br>2021-02-28 | 43QCA,<br>43QDA | 0.00%    | 0.00%   | 0.1147     |
| 2022 | 8      | 2021-11-10,<br>2021-12-10,<br>2021-12-20,<br>2021-12-25                               | 43QCA,<br>43QDA | 0.00%    | 2.81%   | 0.1127     |
| 2023 | 8      | 2022-12-20,<br>2022-12-25,<br>2023-01-14,<br>2023-01-19,<br>2023-02-03                | 43QCA,<br>43QDA | 0.00%    | 0.00%   | 0.1100     |
| 2024 | 8      | 2023-11-15,<br>2024-02-03,<br>2024-02-18,<br>2024-02-23,<br>2024-02-28                | 43QCA,<br>43QDA | 0.00%    | 0.00%   | 0.1225     |

**Data Quality Gate Status: FAILED**

- **Gate 1 (Max NoData  $\leq$  5.0%): PASS** — All composites within nominal threshold ( $\leq$ 5%)
- **Gate 2 (Max YoY Volatility  $\leq$  15.0%): FAIL** — Exceeded in: 2019 (31.7%), 2021 (18.5%)
- **Gate 3 (Min Accuracy  $\geq$  70.0%): PASS** — All evaluated years  $\geq$  70.0%

- **Gate 4 (Loss/Gain Ratio  $\leq$  30.0%):** **PASS** — Transition loss verified



## 6. Engineering & Operations

### Technology Stack

| Layer         | Technology                        | Purpose                                                   |
|---------------|-----------------------------------|-----------------------------------------------------------|
| EO Ingestion  | PySTAC-Client, Stackstac, Dask    | Cloud-native STAC scene retrieval & median reduction      |
| ML Pipeline   | Scikit-Learn, Rasterio, GeoPandas | Balanced Random Forest classification & spatial filtering |
| Spatial DB    | PostgreSQL 16 + PostGIS 3.4       | Spatial envelopes, audit runs, time-series storage        |
| Backend API   | FastAPI, Uvicorn, Pydantic        | High-throughput REST API with GeoJSON endpoints           |
| Frontend UI   | MapLibre GL JS, Vanilla HTML/CSS  | WebGL swipe map, time slider, interactive analytics       |
| Orchestration | k3s / k3d, Docker Compose         | Lightweight container & Kubernetes cluster parity         |
| CI/CD         | GitHub Actions                    | Automated linting, test execution, Trivy CVE scanning     |
| Monitoring    | Prometheus, Grafana               | Custom .prom metrics exporter & sprawl monitoring         |

### Test Suite & Payload Metrics

AUTOMATED UNIT & INTEGRATION TESTS

## 93 Test Cases

100% Passing Coverage across Scaling, Cleanup, Gate & Metrics

UrbanPulse is 100% configuration-driven. Add configs/<city>.yaml with the exact keys:

```
city:
  name: Jaipur
  state: Rajasthan
  country: India
spatial:
  bbox: [75.6000, 26.7000, 76.0500, 27.1500]
  crs: EPSG:4326
  center_lat: 26.9124
  center_lon: 75.7873
  aoi_size_km: 45.0
temporal:
  analysis_years:
    start_year: 2018
    end_year: 2024
  dry_season:
    start_month: 11
    end_month: 2
    start_day: 11-01
    end_day: 02-28
stac:
  earth_search_url: https://earth-search.aws.element84.com/v1
  collections:
    sentinel_2: sentinel-2-c1-l2a
```

## 7. Limitations

---

- accuracy is agreement with ESA WorldCover, not field-verified ground truth
- Sentinel-2 covers 2017 onward only
- dry-season composites confuse fallow farmland with built-up land
- the model is not transferable across sensors
- the persistence rule forces non-decreasing built-up area and is a documented assumption
- 10-60 m resolution limits small features.



## 8. Future Roadmap

---

### 1. Historical Landsat Integration (2000–2016)

Extend the temporal horizon by 16 years using harmonized Landsat 5/7/8 surface reflectance stacks with cross-calibration regression.

### 2. CA-Markov 2035 Urban Growth Forecasting

Couple Cellular Automata and Markov Transition Chains with slope, transport proximity, and zoning constraint layers for 2035 sprawl forecasting.

### 3. Hand-Labelled Ground-Truth Validation Set

Digitize high-resolution aerial imagery ground truth across 500 stratified random points to measure real-world precision beyond global maps.

### 4. Dynamic "Run This City" API Endpoint

Expose asynchronous Celery / Redis worker endpoints allowing users to submit any arbitrary bounding box for on-demand sprawl computation.

## 9. Reproducibility & Verification

---

All pipelines, tests, and documentation artifacts can be executed locally or in containerized environments with standard commands:

```
# 1. Start full local stack (Web + API + DB + Prometheus)
make up
```

```
# 2. Run complete pipeline for a specific city and target year
make pipeline CITY=pune YEAR=2024
```

```
# 3. Execute standalone Python pipeline CLI
python -m pipeline.run_city --city ahmedabad
```

```
# 4. Run automated test suite
pytest -v
```

```
# 5. Re-generate this documentation and PDF report
make docs
```

| Component     | Version / Status |
|---------------|------------------|
| Python        | 3.14.2           |
| Rasterio      | 1.5.1            |
| GeoPandas     | 1.1.4            |
| Scikit-Learn  | 1.8.0            |
| Numpy         | 2.4.4            |
| Pandas        | 2.3.3            |
| Dask          | 2026.8.0         |
| Stackstac     | 0.5.1            |
| PySTAC-Client | 0.9.0            |
| Matplotlib    | 3.10.9           |
| Playwright    | Available        |