

# DEDER GENERAL HOSPITAL EMERGENCY INJURY AND CRITICAL CARE DEPARTMENT



# CLINICAL AUDIT TO IMPROVE THE QUALITY OF CLINICAL CARE OF EMERGENCY MANAGEMENT OF POISONING PATIENTS

By: Murad Amin (BSc Nurse)- Emergency head

Dr.Samuel Shimelis (MD, Emergency Director)-Team leader

### <u>Advisors:</u>

\* HSQU Team

Dader, Oromia

December 2017E.C

## Emergency and critical care case team clinical Audit/QI members

| S/N | Full Name          | Status             | Role             |
|-----|--------------------|--------------------|------------------|
| 1.  | Dr.Samuel Shimelis | Emergency Director | Chairperson      |
| 2.  | Murad Amin         | Emergency Head     | Secretary        |
| 3.  | Wardi Usman        | Staff              | Deputy Secretary |
| 4.  | Dachas Shamsadin   | Staff              | Member           |
| 5.  | Zabib Abrahim      | Staff              | Member           |
| 6.  | Alamudin Sufiyan   | Staff              | Member           |
| 7.  | Yosef Tesfaye      | Staff              | Member           |
| 8.  | Buzu Seyum         | Staff              | Member           |

### **Table of Contents**

| Emergency and critical care case team clinical Audit/Q1 members | 2   |
|---|-----|
| Lists of Figures and Tables                                     | 4   |
| Implementation Status of Previous Action Plan                   | 5   |
| Background  | 5   |
| Aim   | 5   |
| Objectives  | 5   |
| Methods   | 6   |
| Study area & period   | 6   |
| Study design  | 6   |
| Source population   | 6   |
| Sampling technique  | 6   |
| Study Variables   | 6   |
| Data collection method  | 6   |
| Data Processing & analysis                                      | 6   |
| RESULT  | 7   |
| Recommendations   | 13  |
| References  | 1./ |

# **Lists of Figures and Tables**

| Figure 1: score for each criterion/standard for emergency management of poisoning patie | ents, |
|---|-------|
| Dec 2017E.C   | 9     |
| Figure 2: Appropriate evaluation and management   | 10    |
| Figure 3: Appropriate history and physical examination                                  | 11    |
| Figure 4: Appropriate history and physical examination                                  | 12    |
|   |       |
|   |       |
| Table 1: ACTUAL PERFORMANCE ANDV PERFORMANCE AGAINST TARGET                             | 8     |

#### Implementation Status of Previous Action Plan

The previous action plan aimed to address gaps in the emergency management of poisoning patients. Key interventions included training healthcare providers on standard poisoning care principles, improving documentation practices, and ensuring the availability of diagnostic resources. Implementation showed improvement in some areas, such as the identification information and disposition documentation, achieving 100% compliance. However, significant gaps remained in conducting relevant investigations (29%) and detailed history-taking (70%)

#### **Background**

Poisoning is a critical public health issue requiring timely and effective emergency care to prevent morbidity and mortality. The management of poisoning patients involves systematic identification, evaluation, treatment, and disposition based on evidence-based guidelines. Clinical audits are essential to assess adherence to standard care practices and identify areas for improvement. This audit was conducted to evaluate the quality of clinical care provided to poisoning patients in the emergency department.

#### Aim

• To improve the quality of clinical care for poisoning patients in the emergency department by identifying gaps in practice and implementing targeted quality improvement interventions.

#### **Objectives**

- To improve the quality of clinical care provided to a patient presenting to ED with poisoning
- To ensure poisoning patients presenting to the ED are resuscitated appropriately
- To ensure poisoning patients presenting to the ED are evaluated appropriately
- To ensure identification of exact poisoning or toxidrome identification is done in the ED
- To ensure appropriate initial treatment is started in the ED

#### Methods

#### Study area & period

The clinical audit was conducted in EOPD of Deder General Hospital from September 21-December 20, 2017EC

#### Study design

Retrospective cross-sectional study

#### Source population

All patients poison come to and cards are available during the study period.

#### Sampling technique

A total of 10 medical records (client chart) of the last reporting quarter should be sampled for the audit. The individual client charts were withdrawn by systematic random sampling. Clinical audit is not research. It is about evaluating compliance with standards rather than creating new knowledge, therefore sample sizes for data collection are often a compromise between the statistical validity of the results and pragmatic issues around data collection i.e., time, access to data, costs. The sample should be small enough to allow for speedy data collection but large enough to be representative. In some audits the sample will be time driven and in others it will be numerical.

#### **Study Variables**

#### **Dependent variables:**

Poison care

#### **Independent Variables**

Demographic poison, Emergency OPD

Data collection method

Data extraction sheet was adapted from National clinical audit tool

#### Data Processing & analysis

Data from extraction sheets was manually verified and entered into the SPSS version 25 software for analysis. The software checked data types, sizes, classifications, and allowable values. Corrections were made, and the findings were presented in tables and figures.

#### **RESULT**

The overall compliance rate was calculated at 79%, emphasizing the need for targeted interventions to bridge the remaining gaps (**figure 1**).

The audit revealed mixed compliance levels across various components of emergency care for poisoning patients. Identification information, diagnosis, treatment, and disposition all achieved 100% compliance, indicating that standard protocols for these aspects are well-established and adhered to effectively. Similarly, documentation of care provider identification and patient discharge outcomes also demonstrated perfect compliance, reflecting robust administrative practices in these areas. However, significant gaps were identified in conducting relevant investigations (29%) and detailed history-taking and physical examinations (70%). These deficiencies suggest potential limitations in resource availability, clinical workflow, or provider training. Evaluation and management of acute life-threatening injuries scored 96%, indicating minor lapses that require attention (**Table 1**).




Figure 1: Overall Performance of emergency Management Poisoning Patients, Dec, 2017E.C

Table 1: ACTUAL PERFORMANCE ANDV PERFORMANCE AGAINST TARGET

| Sno | Variables  | Target | Actual      | Performance    |  |
|-----|--|--------|-------------|----------------|--|
|     |  |        | Performance | against target |  |
| 1   | Identification information recorded                          | 100    | 100         | 0              |  |
| 2   | Evaluation and management of acute life-threatening injuries | 100    | 96          | 4              |  |
| 3   | Detailed history and physical examination                    | 100    | 70          | 20             |  |
| 4   | Relevant investigation (RFT, LFT, Coagulation profile, and   | 100    | 29          | 79             |  |
|     | ECG) conducted   |        |             |                |  |
| 5   | Accurate diagnosis   | 100    | 100         | О              |  |
| 6   | Appropriate treatment  | 100    | 100         | О              |  |
| 7   | Appropriate disposition:                                     | 100    | 100         | О              |  |
| 8   | Care provider identification documented                      | 100    | 100         | 0              |  |
| 9   | Patient discharged alive                                     | 80     | 100         | 0              |  |
|     | Total Percentage (%)   | 100    | 795/9=70    | 103            |  |

Graph showing score for each criterion/standard for emergency management of poisoning patients.




Figure 1: score for each criterion/standard for emergency management of poisoning patients, Dec 2017E.C

#### GRAPH SHOWING APPROPRIATE EVALUATION AND MANAGEMENT

The evaluation of acute life-threatening poisoning management demonstrates strong compliance with critical interventions, such as airway management, oxygen administration, IV access establishment, and decontamination, each achieving compliance rates of 95-100%. However, areas requiring improvement include Glasgow Coma Scale (GCS) assessment (60%) and Random Blood Sugar (RBS) monitoring (79%), which are crucial for neurological and metabolic support. These gaps suggest a need for enhanced training and reinforcement of protocols to ensure consistent application of these assessments. Overall, the results reflect a well-implemented poisoning care protocol with opportunities for targeted improvements to further optimize patient outcomes (figure2).




Figure 2: Appropriate evaluation and management

#### GRAPH SHOWING APPROPRIATE HISTORY AND PHYSICAL EXAMINATION

While several critical aspects of history-taking and examination were completed thoroughly (e.g., signs/symptoms, type of poisoning, time of exposure), some areas, such as allergy/medication history and last oral intake, were entirely omitted. Additionally, the inquiry into the reason for exposure was only partially addressed. (figure 3).




Figure 3: Appropriate history and physical examination

#### GRAPH SHOWING RELEVANT INVESTIGATION

The findings indicate a significant gap in performing investigations essential for managing a poisoning case. While only CBC investigations was completed, the following key investigations were not performed, even for patients with indications:

- Renal and Liver Function Tests (RFT and LFT):
- Serum Electrolytes (Na, K, Mg)
- Coagulation Profile (PT, PPT, INR).
- ECG (figure 4).




Figure 4: RELEVANT INVESTIGATION

#### Recommendations

- 1. Strengthen the process for ordering and performing investigations:
- 2. Conduct regular training on POISON management protocols.
- 3. Monitor and evaluate the implementation of corrective actions through periodic audits.
- 4. Improve interdepartmental coordination to expedite investigations.



# DEDER GENERAL HOSPITAL CLINICAL AUDIT QUALITY IMPROVEMENT PLAN/PDSA FORM

Clinical Audit Title: TO IMPROVE THE QUALITY OF CLINICAL CARE OF EMERGENCY MANAGEMENT OF POISONING PATIENTS

Clinical Audit Lead: <u>Dr.Samuel Shimelis (GP)</u>. Department /Team: E<u>OPD</u> Date: <u>28/4/2017E.C</u>

|  | Plan  |   |                                   |  |  | DO   | STUDY  | ACT   |
|--|---|---|-----------------------------------|--|--|--|--|---|
| Recommenda<br>tion   | Actions to address the recommendation/Change idea   | Person Responsible                              | Target<br>Date                    | Data collection plan   |  | plan. Record data,<br>observations and<br>modifications to<br>the plan. Use<br>visual<br>descriptions such | synthesis. Do the results align with the explicit criteria? Write the progress made in the implementation, the difficulties faced and actions taken to address them. | Decision: What action are we going to take as a result of this cycle (Adopt, Adapt, or Abandon)? Are we ready to implement? What other processes or systems might be affected by this |
|  | we need to try the change?  | Who will perform the test?<br>(Name or<br>Role) | When will<br>this be<br>complete? | data? (Checklist,<br>Chart audit,)                           | Who will<br>collect the<br>data? (Name or<br>Role) |  |  |   |
| Perform appropriate evaluation and management  | Write feedback to emergency OPD assigned physicians   | EOPD Director (Dr. Samuel<br>Sh)                | until Jan<br>20,<br>2017E.C       | Audit patient<br>records monthly<br>managed for<br>poisoning | QU (Abdi T+<br>Abdella A)                          |  |  |   |
| Facilitate Access to Essential Investigations ( RFT, LFT, serum electrolytes, coagulation profile, ECG). | <ul> <li>Procure necessary laboratory investigation.</li> <li>Order/do necessary lab investigation for all poisoning patients.</li> </ul> | CEO & MD (Nuredin &<br>Dr. Derese)              | Until Mar<br>20,<br>2017E.C       | Audit patient<br>records monthly<br>managed for<br>poisoning | Quality<br>U(Abdi T+<br>Abdella A)                 |  |  |   |

| Adapt (Modify this change and plan next PDSA cycle; loop back to "Plan") Abandon (Change didn't work/won't lead to improvement. Identify new change; plan new PDSA cycle; loop back to "Plan")  Adopt (Data revealed this change was effective and worked well; Next step, develop implementation plan) >>>> |               |           |                         |  |  |  |
|--|---------------|-----------|-------------------------|--|--|--|
|  | Completed by: | Sign off: | Date of review of PDSA: |  |  |  |

#### References

- 1. World Health Organization (WHO). Clinical Management of Acute Poisoning: Guidance for Health-Care Providers. WHO, 2020.
- 2. American College of Emergency Physicians (ACEP). Clinical Policy for the Initial Approach to Patients Presenting with Acute Toxic Ingestion. Annals of Emergency Medicine, 2017.
- National Poisons Information Service (NPIS). Management of Poisoning in Emergency Care. Clinical Toxicology Guidelines, 2019.
- 4. Goldfrank, L. R., et al. **Goldfrank's Toxicologic Emergencies** (11th Edition). McGraw-Hill Education, 2019.