



25.

a. min. = 0
 25% = 14.12
 med. = 33.15
 mean = 51.45
 75% = 69.44
 max. = 1645.56



b.

```
# advanced
summary(cex5$foodaway[cex5$advanced == 1])

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##    0.00   21.67   48.34   75.37   89.86  1179.00


# college
summary(cex5$foodaway[cex5$college == 1])

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##    0.00   14.98   36.11   54.98   72.22  1645.56

# no advanced or college
summary(cex5$foodaway[cex5$advanced == 0 & cex5$college == 0])

##    Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##    0.00    9.63   25.02   38.32   55.97   437.78
```


c.



d.

```
## Call:
## lm(formula = log(foodaway) ~ income, data = cex5[cex5$foodaway > 0, ])
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.3034 -0.6003  0.0654  0.6119  3.6262
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  3.1364599   0.0420567    74.58  <2e-16 ***
## income       0.0069283   0.0004898    14.15  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.898 on 2003 degrees of freedom
## Multiple R-squared:  0.09083,    Adjusted R-squared:  0.09038
## F-statistic: 200.1 on 1 and 2003 DF,  p-value: < 2.2e-16
```

e.



f.

