13.



Q13 Yes, it is a normal dist.
The y-axis is cumulative dist. function, and since most points are around the 50% area, it follows a normal dist.

Applying the Basics

20. Cerebral Blood Flow Cerebral blood flow (CBF) i the brains of healthy people is normally distributed wi a mean of 74. A random sample of 25 stroke patients resulted in an average CBF of $\bar{x} = 69.7$ with a standard deviation of $s = 16$. Test the hypothesis that the averag CBF for stroke patients is different from that of healthy people using $\alpha = .05$.

Q20

1) $H_0 : \mu = 74$

$H_1 : \mu \neq 74$

2) $\alpha = 0.05$

3) $t_{stat} = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \sim t(n-1)$

4) $t^* = \frac{69.7 - 74}{16/\sqrt{25}} = -1.34375$

5) $t_{0.025} = \pm 2.064$

6) Fail to rej H_0 since it lies within the critical values

p-value.

$p = 0.192 > 0.05$

$\therefore$ Do not rej H_0.

.00 g with probability equal to .95?

30. Bulimia An article in the *British Journal of Clinical Psychology* indicates that self-esteem was an important predictor of those who will benefit from treatment for bulimia.[4] The table gives the mean and standard deviation of self-esteem scores prior to treatment, at posttreatment, and during a follow-up:

	Pretreatment	Posttreatment	Follow-up
Sample Mean $\bar{x}$	20.3	26.6	27.7
Standard Deviation s	5.0	7.4	8.2
Sample Size n	21	21	20

a. Use a test of hypothesis to determine whether there is sufficient evidence to conclude that the true pretreatment mean is less than 25.

b. Construct a 95% confidence interval for the true posttreatment mean.

c. In Section 10.3, we will introduce small-sample techniques for making inferences about the difference between two population means. Without a formal statistical test, what might you conclude about the differences among the three sampled population means represented by the results in the table?

Q30 a) $\bar{x} = 20.3$

$s = 5.0$

$n = 21$

1) $H_0 : \mu = 25$

$H_1 : \mu < 25$

2) Assume $\alpha = 0.05$

3) $t_{stat} = \frac{\bar{x} - \mu}{s/\sqrt{n}} \sim t(n-1)$

4) $t^* = \frac{20.3 - 25}{5/\sqrt{21}}$

$= -4.38162$

$t_{0.05} = -1.725 > -4.38$

$\therefore$ rej H_0.

b) $\bar{x} = 26.6$

$s = 7.4$

$n = 21$

$C.I. = 26.6 \pm t_{0.025,20} \left(\frac{7.4}{\sqrt{21}}\right)$

$= 26.61 \pm 2.086 \times 1.61481$

$= (23.2722, 29.9281)$

c) After treatment, people had higher self-esteem as shown by the mean.

35. Sodium Chloride Measurements of water intake, obtained from a sample of 17 rats that had been injected with a sodium chloride solution, produced a mean and standard deviation of 31.0 and 6.2 cubic centimeters (cm^3), respectively. Suppose that the average water intake for noninjected rats observed over a comparable period of time is 22.0 cm^3.

a. Do the data indicate that injected rats drink more water than noninjected rats? Test at the 5% level of significance.

b. Find a 90% confidence interval for the mean water intake for injected rats.

Q 35 a) 1) $H_0 : \mu = 22$ $\quad \bar{x} = 31$, $s = 6.2$

$H_1 : \mu > 22$

2) $\alpha = 0.05$

3) $t_{stat} = \frac{\bar{x} - \mu}{s/\sqrt{n}}$

4) $t^* = \frac{31 - 22}{6.2/\sqrt{17}}$

$= 5.985$

5) R.R. $= t_{16, 0.05}$

$= 1.746$

6) Since $5.985 > 1.746$, do not rej H_0

$\therefore$ Indicates that injected rats drink more.

b) C.I. $= 31 \pm t_{0.05, 16} \left(\frac{s}{\sqrt{n}}\right)$

$= 31 \pm 1.746 \times \frac{6.2}{\sqrt{17}}$

$= 31 \pm 2.1125$

$= (28.775, 33.225)$

Two-Sample Tests *Use the information provided in Exercises 7–8. State the null and alternative hypotheses for testing for a significant difference in means, calculate the pooled estimate of σ^2, the associated degrees of freedom, and the observed value of the t statistic. What is the rejection region using $\alpha = .05$? What is the p-value for the test? What can you conclude from these data?*

7.

Population 1	12	3	8	5	
Population 2	14	7	7	9	6

8.

Population	1	2
Sample Size	16	13
Sample Mean	34.6	32.2
Sample Variance	4.8	5.9

Q7. $H_0: \mu_1 = \mu_2$

$H_1: \mu_1 \neq \mu_2$

$\mu_1 = \frac{12+3+8+5}{4} = 7$

$Var_1^2 = 15.33$

$\mu_2 = \frac{14+7+7+9+6}{5} = 8.6$

$Var_2^2 = 10.3$

$\frac{15.33}{10.3} < 3$

$S_p^2 = \frac{(4-1)15.33+(5-1)10.3}{5+4-2} = 12.4$

$df = 5+4-2$

$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{S_p^2(\frac{1}{n_1}+\frac{1}{n_2})}} = \frac{7-8.6}{\sqrt{3.4(\frac{1}{4}+\frac{1}{5})}} = -0.675$

R.R.: two-tailed

$t = \pm 2.365$

rej if $|t| > 2.365$

p-value $p > 0.5$

Since $|-0.675| < 2.365$, fail to rej

Q8 $H_0: \mu_1 = \mu_2$

$H_1: \mu_1 \neq \mu_2$

pooled estimate test: $\frac{5.9}{4.8} = 1.229$

pooled estimate: $\frac{(16-1)4.8+(13-1)5.9}{16+13-2}$

$S_p^2 = 5.2889$

Df: 27

$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{S_p^2(\frac{1}{n_1}+\frac{1}{n_2})}} = \frac{34.6-32.2}{\sqrt{5.2889(\frac{1}{16}+\frac{1}{13})}} = 2.80$

R.R.: two tailed

$t = \pm 2.052$

Rej if $|t| > 2.052$

p value: $t = 2.80$

p value $= 0.009$

Since $2.80 > 2.052$, we rej the null hypothesis

12. Books or iPads? An experiment was conducted to compare the use of iPads versus regular textbooks in teaching algebra to two classes of middle school students.[6] To remove teacher-to-teacher variation, the same teacher taught both classes, and all teaching materials were provided by the same author and publisher. Suppose that after 1 month, 10 students were selected from each class and their scores on an algebra advancement test recorded. The summarized data follow.

	iPad	Textbook
Mean	86.4	79.7
Standard Deviation	8.95	10.7
Sample Size	10	10

a. Use the summary data to test for a significant difference in advancement scores for the two groups using $\alpha = .05$.

b. Find a 95% confidence interval for the difference in mean scores for the two groups.

c. In light of parts a and b, what can we say about using iPads versus traditional textbooks in teaching algebra at the middle school level?

Q12 a) $H_0: \mu_1 = \mu_2$

$H_1: \mu_1 \neq \mu_2$

$\frac{10.7^2}{8.95^2} = 1.4294$

$S_p^2 = \frac{(9)8.95^2 + (9)(10.7^2)}{18} = 97.3$

$t_{18} = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{S_p^2(\frac{1}{n_1} + \frac{1}{n_2})}} = \frac{86.4 - 79.7}{\sqrt{97.3(\frac{1}{10} + \frac{1}{10})}} = 1.52$

$t_{18,.9} = 2.101$

Since $1.52 < 2.101$, we fail to reject the null hypothesis.

b)

95% C.I. $= (86.4 - 79.7) \pm 2.101\sqrt{97.3(\frac{1}{10} + \frac{1}{10})}$

$= [-2.57, 15.97]$

c) There is no difference.