17. Disinfectants An experiment was conducted to study the use of 95% ethanol versus 20% bleach as a disinfectant in removing contamination when culturing plant tissues. The experiment was repeated 15 times with each disinfectant, using cuttings of eggplant tissue.[9] The observation reported was the number of uncontaminated eggplant cuttings after a 4-week storage.

Disinfectant	95% Ethanol	20% Bleach
Mean	3.73	4.80
Variance	2.78095	.17143
n	15	15
	Pooled variance 1.47619	

a. Are you willing to assume that the underlying variances are equal?

b. Using the information from part a, is there evidence of a significant difference in the mean numbers of uncontaminated eggplants for the two disinfectants tested?

a. $\frac{S_1^2}{S_2^2} = \frac{2.78095}{0.17143} \approx 16.22 > 3 \Rightarrow$ unequal variances

b. $H_0: \mu_1 = \mu_2 \quad H_a: \mu_1 \neq \mu_2$

$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{S_p^2(\frac{1}{n_1}+\frac{1}{n_2})}} = \frac{3.73 - 4.8}{\sqrt{1.47619(\frac{1}{15}+\frac{1}{15})}} \approx -2.412$

$df = \frac{(\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2})^2}{\frac{(\frac{S_1^2}{n_1})^2}{n_1-1} + \frac{(\frac{S_2^2}{n_2})^2}{n_2-1}} \approx 15.7 \qquad df_{15, 0.05/2} \approx \pm 2.131$

$-2.412 < -2.131 \Rightarrow$ reject H_0.

20. Dissolved O_2 Content A pollution control inspector suspected that a river community was releasing amounts of semitreated sewage into a river. To check his theory, he drew five randomly selected specimens of river water at a location above the town, and another five below. The dissolved oxygen readings (in parts per million) are as follows:

						$\bar{x}$	s^2
① Above Town	4.8	5.2	5.0	4.9	5.1	5	0.025
② Below Town	5.0	4.7	4.9	4.8	4.9	4.86	0.015

a. Do the data provide sufficient evidence to indicate that the mean oxygen content below the town is less than the mean oxygen content above? Test using $\alpha = .05$.

b. Estimate the difference in the mean dissolved oxygen contents for locations above and below the town. Use a 95% confidence interval.

a. $H_0: \mu_1 \geq \mu_2 \quad H_a: \mu_1 < \mu_2 \qquad \frac{S_1^2}{S_2^2} = 1.67 < 3$

$S_p^2 = \frac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{n_1+n_2-2} = 0.02$

$t = \frac{\bar{x}_1 - \bar{x}_2 - D_0}{S_p\sqrt{\frac{1}{n_1}+\frac{1}{n_2}}} = \frac{4.86 - 5}{0.14\sqrt{\frac{2}{5}}} \approx -1.56$

$t_{5+5-2, 0.05} = -1.86 < -1.56 \Rightarrow$ non reject H_0

b. $(\bar{x}_1 - \bar{x}_2) \pm t_{8, 0.05/2} \cdot S_p\sqrt{\frac{1}{n_1}+\frac{1}{n_2}} = 5 - 4.86 \pm 2.306 \times 0.0895$
$= (-0.066, 0.346)$

Paired-Difference Tests II *Use the information in Exercise 9 to calculate the observed value of the t statistic for testing the difference between the two population means using paired data. Find a rejection region with $\alpha = .05$ to state your conclusions. Construct a 95% confidence interval for $\mu_1 - \mu_2$. Does this confirm the results of your hypothesis test?*

9.

	Pairs				
Population	1	2	3	4	5
1	1.3	1.6	1.1	1.4	1.7
2	1.2	1.5	1.1	1.2	1.8
d	0.1	0.1	0	0.2	-0.1

$\bar{d} = 0.06 \quad S_d \approx 0.114$

$H_0: \mu_d = 0 \quad H_a: \mu_d \neq 0$

$t = \frac{\bar{d} - 0}{S_d/\sqrt{n}} = \frac{0.06}{0.114/\sqrt{5}} \approx 1.176$

$df = 5 - 1 = 4 \qquad t_{4, 0.05/2} = \pm 2.776$

rejection region: $|t| > 2.776$

$|1.176| \not> 2.776 \Rightarrow$ non-reject H_0

$\mu_1 - \mu_2$ 的 95% C.I. $\quad \bar{d} \pm t_{4, 0.05/2} \times \frac{S_d}{\sqrt{n}}$

$= 0.06 \pm 2.776 \times 0.051$

$= (-0.0816, 0.2016)$ contains 0

$\Rightarrow$ non-reject H_0

DATA SET DS1026 **20. Memory Experiments** Twenty students participated in an experiment in which they were first asked to recall a list of 25 words. They were then given a second list, and asked to form images of the words as they were read. The number of words recalled for each student is given in the table.

Student	With Imagery	Without Imagery
1	20	5
2	24	9
3	20	5
4	18	9
5	22	6
6	19	11
7	20	8
8	19	11
9	17	7
10	21	9
11	17	8
12	20	16
13	20	10
14	16	12
15	24	7
16	22	9
17	25	21
18	21	14
19	19	12
20	23	13

Does it appear that the average recall score is higher when imagery is used?

$\bar{d} = 10.25 \quad S_d \approx 4.051 \quad df = 20 - 1 = 19$

$H_0: \mu_d = 0 \quad H_a: \mu_d > 0$

$t = \frac{\bar{d}}{S_d/\sqrt{n}} = \frac{10.25}{4.051/\sqrt{20}} = 11.317$

$t_{19, 0.05/2} = 1.729 < 11.317 \Rightarrow$ reject H_0

21. Music in the Workplace Before contract

DATA SET DS1028

22. Cake Mixes An experiment was conducted to compare the densities (in ounces per cubic inch) of cakes prepared from two different cake mixes. Six cake pans were filled with batter A, and six were filled with batter B. Expecting a variation in oven temperature, the experimenter placed a pan filled with batter A and another with batter B *side by side* at six different locations in the oven. The six paired observations of densities are as follows:

Location	1	2	3	4	5	6
Batter A	.135	.102	.098	.141	.131	.144
Batter B	.129	.120	.112	.152	.135	.163

a. Do the data present sufficient evidence to indicate a difference between the average densities of cakes prepared using the two types of batter?

b. Construct a 95% confidence interval for the difference between the average densities for the two mixes.

a. $\bar{d} = -0.01$ $\quad S_d \approx 0.0095$

$H_0: \mu_d = 0 \quad H_a: \mu_d \neq 0$

$t = \frac{\bar{d}}{S_d/\sqrt{n}} = \frac{-0.01}{0.095/\sqrt{6}} = -2.58$

$t_{5,\, 0.05/2} = \pm 2.571$

$-2.58 < -2.571 \Rightarrow$ reject H_0

b. $\bar{d} \pm t^* \cdot \frac{S_d}{\sqrt{n}} = -0.01 \pm 2.571 \times \frac{0.0095}{\sqrt{6}} = (-0.01998, -0.00002)$

DATA SET DS1041

12. Reaction Times II Refer to Exercise 11. Suppose that the experiment is conducted using eight people as blocks and making a comparison of reaction times within each person; that is, each person is subjected to both stimuli in a random order. The reaction times (in seconds) for the experiment are as follows:

Person	Stimulus 1	Stimulus 2
1	3	4
2	1	2
3	1	3
4	2	1
5	1	2
6	2	3
7	3	3
8	2	3

Do the data present sufficient evidence to indicate a difference in mean reaction times for the two stimuli? Test using $\alpha = .05$.

$H_0: \mu_d = 0 \quad \mu_d \neq 0$

$\bar{d} = -0.75$

$S_d = 0.707$

$t = \frac{-0.75}{0.707/\sqrt{8}} \approx -3$

$t_{7,\, 0.05/2} \approx \pm 2.449 > -3 \Rightarrow$ reject H_0

13. Instrument Precision A precision instrument is guaranteed to read accurately to within 2 units. A sample of four instrument readings on the same object yielded the measurements 353, 351, 351, and 355.

a. Test the null hypothesis that $\sigma = .7$ against the alternative $\sigma > .7$. Use $\alpha = .05$.

b. Find a 90% confidence interval for the population variance.

$H_0: \sigma = 0.7 \quad H_a: \sigma > 0.7$

$\bar{x} = 352.5 \quad s^2 = 3.66$

$\chi^2 = \frac{(n-1)s^2}{\sigma^2} = 22.43$

$df = n-1 = 3 \quad \chi^2_{3,\, 0.05} = 7.815 < 22.45 \Rightarrow$ reject H_0

$\left(\frac{(n-1)s^2}{\chi^2_{\alpha/2,\, df}}, \frac{(n-1)s^2}{\chi^2_{1-\alpha/2,\, df}} \right) = [1.408, 31.25)$