

**HaDy\_MZB**

**Habitat Dynamics MacroZooBenthos**

**A Python toolbox for analyzing patch-scale habitat dynamics**

---

# **User manual**

**Version: 08-2026**

---

Aude Lecrivain, Christine Weber, Nico Bätz

Eawag, Swiss Federal Institute of Aquatic Science and Technology,  
Surface Waters – Research and Management, Kastanienbaum, Switzerland

GitHub repository:

[https://github.com/HaDy-toolbox/HaDy\\_MZB](https://github.com/HaDy-toolbox/HaDy_MZB)

DOI: XXXX

## Impressum

### Authors of the toolbox and user manual

Aude Lecrivain, Christine Weber and Nico Bätz

### Scientific background

This toolbox implements the framework presented in Lecrivain et al. (in Review).

### Technical support

We sincerely thank James Runnalls, Eawag, for technical support and advice.

### Toolbox testing

We sincerely thank the following colleagues for testing the toolbox and providing valuable feedback on its usability, documentation, and practical applicability:

Tobias Hägele, sje Ecohydraulic Engineering GmbH

Fabio Tarena, RiverLy INRAE Lyon

Hervé Capra, RiverLy INRAE Lyon

Hervé Pella, RiverLy INRAE Lyon

Tabea Donauer, Axpo Group

Joël Wittmann, IfU ETHZ

Henri Guanzini, Drosera Ecologie Appliquée SA

Lucie Riquet, géau environnements SA

### Funding

This work was funded by the Swiss Federal Office for the Environment (FOEN).

### How to cite the tool

## Table of contents

|                                                                                              |           |
|----------------------------------------------------------------------------------------------|-----------|
| <b>1. Toolbox presentation .....</b>                                                         | <b>4</b>  |
| Example case study Box 1: Quick Input – Output example (no download or setup required) ..... | 8         |
| <b>2. Technical setup.....</b>                                                               | <b>9</b>  |
| Example case study Box 2: From python repository setup to running the code .....             | 11        |
| <b>3. Input and configuration .....</b>                                                      | <b>13</b> |
| <b>4. Full workflow explanation .....</b>                                                    | <b>21</b> |
| <b>5. Output interpretation.....</b>                                                         | <b>29</b> |
| Example case study Box 3: Outputs from the main.py file execution .....                      | 30        |
| Example case study Box 4: Outputs from the plots.py file execution .....                     | 34        |
| Example case study Box 5: Outputs from the clustering.py file execution .....                | 40        |
| <b>6. Limitations and assumptions .....</b>                                                  | <b>41</b> |
| <b>7. Trouble shooting .....</b>                                                             | <b>42</b> |
| <b>8. Glossary .....</b>                                                                     | <b>43</b> |
| <b>9. QGIS tips and tricks .....</b>                                                         | <b>45</b> |
| <b>10. Appendix.....</b>                                                                     | <b>49</b> |
| <b>11. Literature references .....</b>                                                       | <b>55</b> |

## 1. Toolbox presentation

### 1.1 Purpose of the HaDy\_MZB toolbox

Rivers are dynamic and habitat conditions change over time as discharge varies and interacts with channel morphology. These changes are especially relevant at the patch scale, where organisms experience hydraulic stress, habitat persistence, dewatering, and shifts between suitable and unsuitable habitat conditions.

HaDy\_MZB is a free and open-source Python-based [HaDy-toolbox](#) for quantifying flow-driven habitat dynamics at the patch scale, with a focus on benthic macroinvertebrates—organisms particularly vulnerable to hydraulic stress due to their limited mobility and complex life cycles. However, the toolbox can also be adapted to other organism groups such as fish eggs and larvae, or vegetation, which may likewise be affected by limited mobility or exposure to changing habitat conditions.

HaDy\_MZB links discharge time-series with precomputed hydraulic information for selected discharges to generate habitat time-series. The toolbox is designed to be flexible with respect to both the spatial and temporal scale of the analysis. The study reach can range from a few meters to tens of kilometers in length, while the temporal resolution of the discharge time-series can vary from minutes to hours. Based on these habitat time-series, the toolbox calculates four spatially explicit metrics that quantify habitat availability, habitat stability, and exposure to hydraulic stress over time (see Table 1). A patch represents a fixed spatial unit corresponding to the spatial and temporal perception of macroinvertebrates, typically about 0.25-0.50 m<sup>2</sup>, defined by the computational mesh used in high-resolution hydrodynamic models (see more in Section 8).

The toolbox builds on the patch-scale approach developed by Bätz et al. (2025) and applied in Lecrivain et al. (in Review). In particular, it puts into practice the logic of deriving habitat time-series from discharge scenarios and habitat distributions. Hydropeaking is a key application because frequent sub-daily discharge fluctuations strongly increase habitat dynamics. However, the toolbox can also be used more generally to analyze habitat responses to variable discharge regimes.

Table 1 Metrics derived by the HaDy\_MZB toolbox

| Metric                    | Definition                                                                                                                           | Relevance                                                               |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------|
| M1<br>Habitat probability | Likelihood of different habitat conditions within a patch through time (Bätz et al.2025)                                             | Time-integrated habitat availability; identifies dominant habitat types |
| M2<br>Habitat shifts      | Daily number of times (i.e., frequency) that habitat conditions change to another suitable habitat type in a patch (Bätz et al.2025) | Indicates habitat stability and temporal persistence                    |
| M3<br>Drift risk          | Risk of macroinvertebrate drift based on current velocity and up ramping rate experienced within each patch over time                | Identifies exposure to critical passive drift conditions                |
| M4<br>Desiccation risk    | Risk of egg desiccation linked to hydropeaking-induced water level variations                                                        | Identifies exposure to critical desiccation duration                    |

#### **Who HaDy\_MZB is for:**

HaDy\_MZB is designed for **researchers** and **practitioners** who want to assess discharge-driven habitat dynamics and ecological stressors over time in a spatially explicit way. It supports applications such as hydropeaking impact assessment, comparison of hydrological scenarios, evaluation of morphological complexity, and reach-scale mitigation planning.

The toolbox does not replace hydrodynamic modelling or ecological expert judgement. Instead, it provides a reproducible workflow for translating existing hydraulic or habitat data and discharge time-series into interpretable habitat-dynamics metrics.

## **1.2 Overview of the Workflow**

The execution of the toolbox involves 3 steps (Figure 1):

- **Pre-processing:** the hydrodynamic simulation of the study area is carried out by the user. It provides spatially distributed data on water depth and current velocity for a representative range of discharges (input shapefile.shp). These data are then used to derive habitat types. The discharge time-series is required to describe the temporal evolution of discharge (input discharge.csv). See more in Section 3. **Error! Reference source not found.**
- **Processing:** the toolbox links discharge time-series with hydraulic data to generate habitat time-series and to calculate selected metrics.
- **Post-processing:** the resulting outputs can be visualized, mapped and further analyzed using GIS (see Section 5**Error! Reference source not found.**, and Section 9).




Figure 1 Overview of the execution procedure of the toolbox HaDy\_MZB

The respective roles of the user and the toolbox are summarized in Table 2.

Table 2 Respective roles of the user and the toolbox throughout the workflow

| Step            | Responsibility | Activity                                                                                                                      |
|-----------------|----------------|-------------------------------------------------------------------------------------------------------------------------------|
| Pre-processing  | User           | Selects the discharge time-series and its timestep (time between two discharge values)                                        |
| Pre-processing  | User           | Selects discharges of interest for the hydrodynamic simulation                                                                |
| Pre-processing  | User           | Selects the study area of interest                                                                                            |
| Pre-processing  | User           | Runs the hydrodynamic simulation                                                                                              |
| Pre-processing  | User           | Decides on a habitat classification method and relevant ecological thresholds                                                 |
| Processing      | Toolbox        | Builds <b>habitat time-series</b> from discharge time-series, water depth, current velocity and habitat classification method |
| Processing      | Toolbox        | <b>Calculates the selected patch scale metrics</b>                                                                            |
| Processing      | Toolbox        | Exports results (i.e. metric values) in a shapefile, preview maps, and a csv document                                         |
| Processing      | Toolbox        | Provides graphical plots                                                                                                      |
| Processing      | Toolbox        | Provides additional functions for deeper analysis if needed                                                                   |
| Post-processing | User           | Analyses the output of the toolbox on GIS software                                                                            |
| Post-processing | User           | Adapts the toolbox to include additional plots or analyses if needed                                                          |

### 1.3 Example case study

This manual follows a **running example dataset** provided with the toolbox (Windows operating system). Blue boxes such as the following provide practical guidance and shortcuts to quickly explore, reproduce, and adapt the workflow. Running the example case as an initial analysis is a good way to become familiar with the toolbox.

Example case study boxes are listed below:

- **Example Case Study Box 1 Quick** Input – Output example (no download or setup required)
- **Example Case Study Box 2:** From python repository setup to running the code
- **Example Case Study Box 3:** Outputs from the main.py file execution
- **Example Case Study Box 4 :** Outputs from the plots.py file
- **Example Case Study Box 5 :** Outputs from the clustering.py file

Moreover, videos are accessible [here](#) to guide throughout the process, using the example case study:

- [0 HaDy GitHub general introduction](#)
- [1 HaDy setup and first run](#)
- [2 HaDy example case from inputs to outputs](#)
- [3 HaDy example case QGIS](#)
- [4 HaDy local yaml for tailored analysis](#)
- [5 HaDy output analysis](#)
- [6 HaDy advanced focus on zone](#)

### Example case study Box 1: Quick Input – Output example (no download or setup required)

This box presents where to find the input and output files related to the example case study on GitHub (Figure 2). This does not require any installation, download or setup.

More information in the videos [0 HaDy GitHub general introduction](#) and [2 HaDy example case from inputs to outputs](#).

Example inputs (GitHub [HaDy-toolbox](#) following folders data > input):

- Discharge data (Week\_timestep\_10\_min.csv)
- Input shapefile with water depth and current velocity value (Example\_input\_shapefile.shp)
- Habitat definition (see Section 4.3)

Example outputs (GitHub [HaDy-toolbox](#) following folders data > output > Example\_output):

- Metrics (folder Metric\_files)
- Maps (folder Metric\_files > Preview\_maps for the preview maps, and folder GIS\_project containing the projects and the templates)
- Plots (folder Plots)
- Clustering (folder Clustering)




Figure 2 Quick input-output overview of the example case study on GitHub



## 2. Technical setup

This chapter explains how to install the toolbox and how to use it.

### 2.1 Prerequisites

If the following usual **python core tools** and **plugins** mentioned below are not installed on the laptop, check Section 10.1.

Required tools:

- Python
- Git
- Anaconda/Miniconda
- Visual Studio Code (recommended), **including the python extension**
- Python extension for VS Code

### 2.2 GitHub repository setup

**All steps are explained and illustrated in Example Case Study Box 2 and in the [video 1 HaDy setup and first run](#).**

*Step 1: Create a working directory where the Git repository will be cloned*

Create a folder where the Git repository will be cloned.

The folder can be located anywhere on the system, provided that its path does not contain any spaces. The folder can be given any name, as long as the name does not contain spaces.

*Step 2: Clone the repository*

Open a terminal and go to the created working directory: enter the command “cd” and the absolute path to your directory (**cd path/to/your/directory**).

Once in the working directory, paste the cloning command (Git needs to be installed otherwise this will not work, see Section 2.1):

**git clone https://github.com/HaDy-toolbox/HaDy\_MZB.git**

*Step 3: Setup the virtual environment*

In the terminal (for macOS and Linux), or anaconda prompt (Windows) go to **the cloned folder called HaDy\_MZB** in the working directory, and paste the following commands:

**conda env create -f environment.yml**, and then **conda activate Venv\_HaDy\_MZB**

*Step 4: Open the project on VS code and select the python interpreter*

Open the software VS code previously installed, open the cloned HaDy\_MZB folder.

To select the interpreter: Ctrl+Shift+P (to get the command panel) → type “Python: Select Interpreter”

→ Choose Venv\_HaDy\_MZB.

#### Step 5: Run the code

Go to the python file *main.py* and click the play button (see [Figure 4.5](#)).


The same steps can be done in the *plots.py* or *clustering.py* for further analysis (see Section 4.6).



If the analysis is tailored, make sure that the *local.yaml* file has been adapted **and saved** (more in Section **Error! Reference source not found.****Error! Reference source not found.**) before running the codes. Otherwise, the outputs will not correspond to the specified parameters.

## Example case study Box 2: *From python repository setup to running the code*

### Step 1: Create a working directory where the Git repository will be cloned



### Step 2: Clone the repository

In the terminal, type in “cd” and paste the absolute path to your directory (no spaces allowed)


```
PS C:\Users\lecrivau> cd C:\Users\lecrivau\source\repos\HaDy-toolbox
```

Then type in the cloning command

```
PS C:\Users\lecrivau\source\repos\HaDy-toolbox> git clone https://github.com/HaDy-toolbox/HaDy_MZB.git
```

### Step 3: Setup the virtual environment

On windows, use Anaconda prompt




Go to the cloned HaDy\_MZB and type in the commands

```
C:\Users\lecrivau\source\repos\HaDy-toolbox>cd C:\Users\lecrivau\source\repos\HaDy-toolbox\HaDy_MZB
C:\Users\lecrivau\source\repos\HaDy-toolbox\HaDy_MZB>conda env create -f environment.yml
rs\lecrivau\source\repos\HaDy-toolbox\HaDy_MZB>conda env create -f environment.yml
```

You now see that the environment is activated.

```
(Venv_HaDy_MZB) C:\Users\lecrivau>
```

### Step 4: Open the project on VS code and select the python interpreter




To select the interpreter: Ctrl+Shift+P (to get the command panel) → type “Python: Select

Interpreter” → Choose `Python 3.11.11 (Venv_HaDy_MZB) ~\conda\envs\Venv_HaDy_MZB\python.exe`

### Step 5: Run the code

For the example case study, run the code without modifying the configuration file (config.yaml)



### 2.3 For later use: get the latest version from GitHub

Since the toolbox is in constant evolution, make sure to use the latest version available. To update it, “pull” the changes from the GitHub folder as done on Figure 3.

This can be done directly in the Visual Studio code terminal as follows: command **git pull origin main**




Figure 3 Example of how to pull the last version



ATTENTION

Running this command replaces the contents of the scripts and input\_data/local folders with the latest files from the GitHub repository. Be sure to back up any local changes before executing the command, as they will be overwritten.

### 3. Input and configuration

#### 3.1 Input files

Input files are located in the data > input folder, which contains two subfolders: Discharge\_data and Shapefile. The relevant files should be placed in the appropriate folder to be included in the analysis. The following sections describe the required file contents and minimum requirements.

##### *Discharge time-series input data*

Minimum requirements:

- A CSV file containing discharge variations over time ( $\text{m}^3/\text{s}$ ), as illustrated in Figure 4.
- The file must be located in data > input > Discharge\_data.
- The column containing discharge values must be named Discharge (exactly as written, including the capital letter). The format of the other columns is not relevant and they may be retained or deleted (e.g., Figure 4, right). A date column is included for user convenience but is not read by the code. The timestep between consecutive discharge values is instead specified by the user in the configuration file.
- The timestep specified in the configuration file must match the timestep between consecutive rows in the input file (see Section **Error! Reference source not found.**). The toolbox requires this information, for example, to calculate daily metrics. It is therefore the user's responsibility to provide an input file with the appropriate timestep for the intended analysis.
- Discharge values must be formatted as numbers, with a dot (.) used as the decimal separator.



#### ATTENTION

##### Common issues:

- **Incorrect file location:** The input file must be placed in the specified folder.
- **Incorrect file format:** The input file must be provided in CSV format, not as an Excel file.
- **Incorrect Discharge column name:** The discharge column must be named Discharge exactly, including the capital letter.
- **Incorrect decimal separator:** Decimal values must use a dot (.), not a comma (,).
- **Missing values:** Missing values must be addressed before running the analysis.
- **Inconsistent timesteps:** The timestep specified in the configuration file must match the timestep between consecutive discharge values in the input file. An incorrect timestep can lead to incorrect metric values.

| Date (TU)                | Valeur (en $\text{m}^3/\text{s}$ ) | Statut | Qualification | MÃ©thode | ContinuitÃ© |
|--------------------------|------------------------------------|--------|---------------|----------|-------------|
| 2025-06-24T00:00:00.000Z | 14.4                               | 12     | 20            | 8        | 0           |
| 2025-06-24T00:10:00.000Z | 14.4                               | 12     | 20            | 8        | 0           |
| 2025-06-24T00:20:00.000Z | 14.4                               | 12     | 20            | 8        | 0           |
| 2025-06-24T00:30:00.000Z | 14.4                               | 12     | 20            | 8        | 0           |



| Date (TU)                | Discharge |
|--------------------------|-----------|
| 2025-06-24T00:00:00.000Z | 14.4      |
| 2025-06-24T00:10:00.000Z | 14.4      |
| 2025-06-24T00:20:00.000Z | 14.4      |
| 2025-06-24T00:30:00.000Z | 14.4      |

Figure 4 Example of data downloaded from the HydroPortail platform (left) and of the data retained and renamed for the analysis (right). The Date column is retained here for user convenience but is not read by the code and is therefore not mandatory.

## Shapefile input data

Minimum requirements:

- A polygon shapefile (.shp and associated files) containing water depth and current velocity data for the discharges of interest (e.g., Figure 5). These data should originate from hydrodynamic simulations.
- The shapefile must be located in data > input > Shapefile.
- The prefixes of the relevant column names must be specified in the configuration file (see Section **Error! Reference source not found.**).
- The shapefile must contain the data listed in Table 3.

Table 3 Data that needs to be in the input polygon shapefile

|                                                           |                                                                                                                                                                                                            |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
|-----------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
| Id                                                        | Id of the patch - One row per patch                                                                                                                                                                        |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| x                                                         | Coordinates of the patch center                                                                                                                                                                            |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| y                                                         |                                                                                                                                                                                                            |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Area m2                                                   | Area associated with each patch, used to normalize metrics by area                                                                                                                                         |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Water depth for each discharge (meters)                   | Water depth values for each discharge of interest. Decimals separated by “_”. For example, with the prefix d_, discharges of 31.8 and 6 m <sup>3</sup> /s correspond to d_31_8 and d_6, respectively.      |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |
| Water current velocity for each discharge (meters/second) | Current velocity values for each discharge of interest. Decimals separated by “_”. For example, with the prefix v_, discharges of 31.8 and 6 m <sup>3</sup> /s correspond to v_31_8 and v_6, respectively. |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |  |

| id     | x          | y          | area | z      | d_1  | v_1  | d_2  | v_2  | d_3  | v_3  | d_4_2 | v_4_2 | d_5_5 | v_5_5 | d_7_4 | v_7_4 | d_9  | v_9  | d_10_6 | v_10_6 | d_12 | v_12 |
|--------|------------|------------|------|--------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|------|------|--------|--------|------|------|
| 276139 | 2696154.44 | 1152347.31 | 0.25 | 937.3  | 0.33 | 0.62 | 0.43 | 0.88 | 0.53 | 1.18 | 0.58  | 1.29  | 0.65  | 1.43  | 0.74  | 1.57  | 0.8  | 1.65 | 0.87   | 1.71   | 0.92 | 1.76 |
| 276140 | 2696154.94 | 1152347.31 | 0.25 | 937.3  | 0.34 | 0.6  | 0.45 | 0.87 | 0.55 | 1.17 | 0.59  | 1.28  | 0.66  | 1.42  | 0.75  | 1.56  | 0.82 | 1.65 | 0.88   | 1.71   | 0.93 | 1.76 |
| 276141 | 2696155.44 | 1152347.31 | 0.25 | 937.28 | 0.35 | 0.57 | 0.46 | 0.82 | 0.56 | 1.11 | 0.61  | 1.21  | 0.68  | 1.36  | 0.77  | 1.5   | 0.84 | 1.59 | 0.9    | 1.65   | 0.95 | 1.7  |

Figure 5 Example of data in the polygon shapefile



ATTENTION

### Common issues:

- **Missing files:** If only the .shp file is present in data > input > Shapefile, the toolbox will not run. All associated shapefile components are required (see the files provided in the Example\_input\_shapefile folder).
- **Empty or N/A values:** Empty or N/A values must be replaced with 0 for the toolbox to run the analysis.
- **Data consistency:** If a patch is dry under a given discharge (i.e., water depth = 0), the corresponding current velocity must also be set to 0.
- **Missing data:** Each patch must have both current velocity and water depth data for every discharge included in the analysis.
- **Incorrect formatting:** The water depth and current velocity field names must include the corresponding discharge value. Decimal points in discharge values must be replaced by an underscore (e.g., d\_31\_8 and v\_31\_8 for 31.8 m<sup>3</sup>/s, or

d\_6 and v\_6 for 6 m<sup>3</sup>/s). The prefixes used for water depth columns must be consistent across all water depth fields, and the prefixes used for current velocity columns must be consistent across all current velocity fields. These prefixes must also match those specified in the configuration file (see Section 3.2).

- **Spatial representation:** The input data must be provided as a polygon shapefile, rather than a point shapefile, to ensure a continuous spatial representation of the results.

## 3.2 Configuration

Minimum requirements:

- To run an analysis that differs from the example case study, a local.yaml file must be created in the repository. This file should be created by copying config.yaml and making the necessary modifications to tailor the analysis (see the video [4 HaDy local yaml for tailored analysis](#)).
- The local.yaml file must be located in the root of the repository, in the same location as config.yaml.



### Common issues:

- **Incorrect file name:** The file must be named local.yaml exactly. It is the user's responsibility to create this file when tailoring the analysis parameters.
- **Missing local.yaml file:** If no local.yaml file is present, the toolbox uses config.yaml by default, which corresponds to the example case study.
- **Analysis does not match the specified parameters:** The local.yaml file must be saved after any modifications. Otherwise, the toolbox may run using the previously saved parameters.
- **Previous analysis overwritten:** Each new toolbox run stores its outputs in data > outputs > Toolbox\_run\_output, overwriting any files already present in this folder. Analyses that need to be retained should therefore be copied to another folder or renamed before running the toolbox again.

Table 4 describes the variables that need to be adapted by the user.

*Table 4 Detail of the configuration variables*

#### *Input files*

| <b>Name of the variable</b>         | <b>Signification</b>                                                                                                                                                      | <b>Status</b> |
|-------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| input_shp_filename                  | Name of the input shapefile located in the data > input > Shapefile folder                                                                                                | Mandatory     |
| crs                                 | CRS of the GIS project                                                                                                                                                    | Mandatory     |
| input_discharge_timeseries_filename | Name of the discharge time-series input file located in the data > input > Discharge_data folder                                                                          | Mandatory     |
| time_step_min                       | Timestep of the discharge time-series in minutes (float, e.g., 10.0 for a 10-minute timestep). <b>/!\ Must match the timestep of the discharge time-series input file</b> | Mandatory     |
| shp_id_colname                      | Name of the id column identifying each patch in the input shapefile                                                                                                       | Mandatory     |
| shp_x_colname                       | Name of the column containing the x-coordinate of each patch in the input shapefile.                                                                                      | Mandatory     |
| shp_y_colname                       | Name of the column containing the y-coordinate of each patch in the input shapefile.                                                                                      | Mandatory     |
| shp_area_colname                    | Name of the column containing the area of each patch, used for potential area-based metric normalization.                                                                 | Mandatory     |
| shp_depth_prefix                    | Prefix preceding the discharge values in the input shapefile (e.g., d_ in d_30_2).                                                                                        | Mandatory     |
| shp_vel_prefix                      | Prefix preceding the discharge values in the input shapefile (e.g., v_ in v_30_2).                                                                                        | Mandatory     |

#### *Habitat classification thresholds*

| <b>Name of the variable</b> | <b>Signification</b>                                                                                                                                                              | <b>Status</b> |
|-----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| depth_threshold             | Minimum water depth (m) required for a patch to be considered wet (float, e.g., 0.01 for 1 cm). Patches are considered wet when their water depth is greater than this threshold. | Mandatory     |
| habitat_velocity_thresholds | Habitat classification thresholds based on current velocity and water depth (see Section 4.3 Classify habitats)                                                                   | Mandatory     |
| number_of_habitats          | Number of habitat types defined by the habitat classification. For example, a classification ranging from 0 (dry) to 6 (highest velocity class) contains 7 habitat types.         | Mandatory     |





### Common issues:

- **Incorrect number of habitats:** Ensure that the number of habitat types is counted correctly. For example, habitat classes ranging from 0 to 6 correspond to 7 habitat types. An incorrect number of habitat types will prevent the plots from being generated correctly.
- **Changes to habitat classification:** If the habitat classification method is modified, the attribute\_habitat\_current\_based function in habitat\_classification.py must also be adapted accordingly.

### *Metrics calculation*

| Name of the variable      | Signification                                                                                                                                                                                                                                                                                                                                                                                             | Status    |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| target_habitat            | Target habitat type for which metrics are calculated. Metrics are only calculated for patches that experience this habitat type at least once during the habitat time series. The value must be an integer corresponding to one of the habitat types defined in the habitat classification. To obtain results for different target habitats, the analysis must be run separately for each target habitat. | Mandatory |
| metrics_to_compute        | Selection of the metrics to calculate. Set to true to calculate a metric and false otherwise.                                                                                                                                                                                                                                                                                                             | Mandatory |
| shift_all_daily           | Daily shifts between all habitat types.                                                                                                                                                                                                                                                                                                                                                                   | Mandatory |
| shift_targ_daily          | Daily shifts between the habitat types defined as suitable in suitable_hab_shifts.                                                                                                                                                                                                                                                                                                                        | Mandatory |
| shift_dry_daily           | Daily shifts to dry conditions, defined as water depth below depth_threshold.                                                                                                                                                                                                                                                                                                                             | Mandatory |
| dry_max                   | Maximum consecutive duration of dry conditions, expressed as a number of timesteps.                                                                                                                                                                                                                                                                                                                       | Mandatory |
| desiccation_risk          | Desiccation risk calculated using the thresholds defined in desiccation_thresholds (see Section 4.4)                                                                                                                                                                                                                                                                                                      | Mandatory |
| drift_percentile          | Drift risk based on the specified drift percentile (see Section 4.4 Metrics calculation)                                                                                                                                                                                                                                                                                                                  | Mandatory |
| drift_max                 | Maximum associated drift risk value (see Section 4.4 Metrics calculation)                                                                                                                                                                                                                                                                                                                                 | Mandatory |
| drift_durations           | Number of timesteps spent in each drift risk category.                                                                                                                                                                                                                                                                                                                                                    | Mandatory |
| suitable_hab_shifts       | List of habitat types considered suitable. Shifts between these habitat types are used to calculate shift_targ_daily.                                                                                                                                                                                                                                                                                     | Mandatory |
| start_at_first_occurrence | Set to true to start the desiccation metric calculation after the first occurrence of the target habitat in the habitat time series. Set to false to start the calculation at the beginning of the time series.                                                                                                                                                                                           | Mandatory |

|                            |                                                                                                                                                                                         |                        |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|
| desiccation_thresholds     | Desiccation thresholds based on the consecutive duration spent outside the water, expressed in hours (see Section 4.4).                                                                 | Mandatory              |
| up_ramp                    | Set to true to use drift_thresholds_with_ramp for drift calculations, taking the up-ramping rate into account. Set to false to use drift_thresholds, without considering the ramp rate. | Mandatory              |
| drift_thresholds_with_ramp | Drift thresholds based on current velocity thresholds (m/s), taking the up-ramping rate into account. Used when up_ramp is set to true (see Section 4.4).                               | Conditional on up_ramp |
| drift_thresholds           | Drift thresholds based on current velocity thresholds (m/s), without considering the up-ramping rate. Used when up_ramp is set to false (see Section 4.4 Metrics calculation).          |                        |



#### ATTENTION

The user is responsible for determining desiccation and drift threshold values appropriate for the macroinvertebrate taxa and habitat type considered in the study. The user must also determine whether an up-ramping threshold is relevant for the calculation of drift risk.

#### Optional: advanced parameters for a focus on a specific zone

| Name of the variable     | Signification                                                                                                                                                                                                                             | Status                       |
|--------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|
| focus_on_zone            | Set to true to focus the analysis on a specific zone (e.g., gravel bars for desiccation of eggs) and false otherwise.                                                                                                                     | Mandatory                    |
| zone_interest_filename   | Name of the shapefile defining the zone of interest, including its associated files, located in the data > input > Focus_zone folder.                                                                                                     | Conditional on focus_on_zone |
| id_polygon_zone_interest | Name of the ID column identifying the polygons in the zone of interest (e.g., individual gravel bars), allowing them to be compared. <b>This column must have a different name from the shp_id_colname column in the input shapefile.</b> |                              |
| zone_interest_field      | Name assigned to the zone of interest in the output files. Patches belonging to the zone are assigned a value of 1, while other patches are assigned a value of 0.                                                                        |                              |

#### Optional: files paths for plots and clustering (further analysis)

| Name of the variable | Signification                                                                                | Status   |
|----------------------|----------------------------------------------------------------------------------------------|----------|
| base_output_path     | Output folder located in data > output, containing the data to be used for further analysis. | Optional |

|                         |                                                                                    |          |
|-------------------------|------------------------------------------------------------------------------------|----------|
| final_csv_path          | Relative path to the output CSV file containing the calculated metric values.      | Optional |
| final_shp_path          | Relative path to the output shapefile containing the calculated metric values.     | Optional |
| static_habitat_csv_path | Relative path to the CSV file containing the habitat classification of each patch. | Optional |

### *Tailoring an analysis*

It is recommended to first run the example case study to become familiar with the toolbox and its workflow.

Once the example case study has been successfully run, the analysis can be progressively tailored.



#### **ATTENTION**

Whenever a parameter or code is modified, save the corresponding file before running the code for the changes to be taken into account.

As a first step, the following simple modifications can be tested:

Create a local.yaml file by copying config.yaml (see video [4 HaDy local yaml for tailored analysis](#)) and modify one parameter at a time.

Simple modifications:

- Change the target habitat
- Modify one habitat classification threshold
- Replace the discharge time-series and adapt the timestep accordingly
- Activate or deactivate one metric

More advanced modifications:

- Change desiccation thresholds
- Change drift thresholds
- Run *plots.py* or *clustering.py*
- Change the habitat classification. In this case, adapt both local.yaml and the attribute\_habitat\_current\_based function in habitat\_classification.py accordingly

Changing the spatial focus of the analysis:

- Set focus\_on\_zone to true
- Add a focus zone shapefile and complete the corresponding variables in local.yaml: zone\_interest\_filename, id\_polygon\_zone\_interest, and zone\_interest\_field
- Adapt the habitat classification method to the specific scope of the analysis, if necessary

Running a fully tailored analysis:

Once familiar with the toolbox, the analysis can be adapted to specific study requirements. The following workflow should be followed:

1. Add the input files to the appropriate input folders, as described for the example case study.
2. Create local.yaml by copying config.yaml, then adapt the parameters to the requirements of the analysis and save the file (Ctrl+S). See the video [4 HaDy local yaml for tailored analysis](#).
3. Adapt the habitat classification, if necessary. If the habitat classification method has been modified, update the relevant function in habitat\_classification.py accordingly and save the changes (Ctrl+S).
4. Run main.py to perform the analysis.
5. Adapt plots.py if necessary to account for new habitat types, including habitat names and colors. Save the changes (Ctrl+S) before running the script.
6. Run clustering.py, if clustering is part of the analysis.
7. Explore the results in a GIS software. Import the output shapefile and, when using QGIS, apply the default layouts provided with the toolbox. See the video [3 HaDy example case QGIS](#).

## 4. Full workflow explanation

This chapter explains what happens inside *main.py*, from input data to final metrics.




Figure 6 Full workflow of the *main.py* file

### 4.1 Match discharge time-series to the closest simulated discharges

To run the hydrodynamic simulations, a set of discharges of interest must first be selected. In the example case study, these discharges cover the 1st to 99th percentile range of the discharge time series, following the approach used in Lecrivain et al. (in **review**). However, this is only an example, and other methods can be used to select the discharges of interest.

The selected discharges should be representative of the discharge time series. For each of these discharges, the hydrodynamic simulations must provide water depth and current velocity values for every patch in the study domain, as described for the input shapefile in Section 3.1.

The toolbox then assigns each discharge in the input discharge time series to the closest discharge for which hydrodynamic data are available. The corresponding discharge is recorded in the `Corresponding_known_discharge` column, as illustrated in Figure 7.

**No interpolation is performed for discharges for which hydrodynamic data are not available.** If greater precision is required, or if more extreme discharges need to be considered, the input hydrodynamic data must be adapted accordingly. For example, additional hydrodynamic simulations can be performed for more extreme discharges.

| Datetime      | Discharge   | Corresponding_known_discharge |
|---------------|-------------|-------------------------------|
| 1/6/2025 0:00 | 1.597203229 | 2                             |
| 1/6/2025 0:10 | 2.260468285 | 2                             |
| 1/6/2025 0:20 | 11.27516493 | 10.6                          |
| 1/6/2025 0:30 | 13.44403064 | 13.4                          |
| 1/6/2025 0:40 | 11.49950344 | 12                            |
| 1/6/2025 0:50 | 9.056749663 | 9                             |

Figure 7 Example of discharge time-series where each discharge is associated to the closest known one from the input shapefile provided by the user

## 4.2 Select relevant discharges and crop the input shapefile

Based on the `Corresponding_known_discharge` values, the toolbox determines the list of discharges to retain for the study. To reduce computational time, the input data are then cropped to retain:

- Only data corresponding to discharges within the scope of the study.
- Only patches that are wet under the maximum discharge considered.

## 4.3 Classify habitats

The user is responsible for defining the habitat classification method according to the scope of the analysis. **Error! Reference source not found.** presents the water depth- and current velocity-based classification used in the example case study. Similar approaches can be implemented with different numbers of habitat types and different threshold values. These parameters must be defined and adapted in both the configuration file (`local.yaml`) and the `habitat_classification.py` file.

| Color      | Current Velocity Class                                   |
|------------|----------------------------------------------------------|
| Yellow     | Dry                                                      |
| Green      | Stagnant to slow current<br>$v_{40} < 5$ cm/s            |
| Purple     | Slow current<br>$5 \leq v_{40} < 25$ cm/s                |
| Orange     | Slow to fast current<br>$25 \leq v_{40} < 75$ cm/s       |
| Light Blue | Fast to very fast current<br>$75 \leq v_{40} < 150$ cm/s |
| Dark Blue  | Very fast current<br>$150 \leq v_{40} < 250$ cm/s        |
| Red        | Violent current<br>$v_{40} \geq 250$ cm/s                |

Figure 8 Example of macroinvertebrate habitat classification at community-level (adapted from Tonolla 2023)

Another example focuses on a specific zone, with `focus_on_zone` set to true. This approach is implemented in the `attribute_habitat_types_zone_only` function in the `habitat_classification.py` script and targets egg deposition on gravel banks. Habitat type -1 indicates that a patch is located outside the zone of interest (gravel banks in this example), while the other habitat types are illustrated in Figure 9.




Figure 9 Example of habitat classification focusing on egg desiccation on gravel banks



#### ATTENTION

When modifying code in the repository (e.g., `habitat_classification.py`), make sure to save the changes before running the code for them to be taken into account. In the file editor, this can be done using `Ctrl+S`.

As previously mentioned, modified files should also be saved in a safe location before pulling the latest version of the Git repository. Pulling the latest version may overwrite local changes.

## 4.4 Metrics calculation

### Scope of the metrics calculation





The simulated domain includes all patches that are wet under the maximum discharge of the discharge time series. Habitat probabilities are calculated for all these wetted patches.

Metrics are calculated only for patches that experience the target habitat type at least once during the habitat time series. The metrics to be calculated can be selected by the user in the configuration file.

### Description of the metrics

The four patch-scale macroinvertebrate habitat metrics presented in Table 5 Overview of the four patch-scale metrics used to characterize macroinvertebrate habitat dynamics, including definitions, calculation are derived from the habitat time series and quantify complementary ecological aspects of habitat dynamics. Habitat probability (M1) and habitat shifts (M2) describe the availability and temporal persistence of suitable conditions and are therefore generally related to habitat quality. Drift (M3) and desiccation (M4) describe exposure to potentially harmful hydraulic events and their associated ecological risks.

Table 5 Overview of the four patch-scale metrics used to characterize macroinvertebrate habitat dynamics, including definitions, calculations, ranges, and ecological relevance

| Metric                     | Definition                                                                                                                                  | Calculation               | Range                                                                                 | Relevance                                                               |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|---------------------------|---------------------------------------------------------------------------------------|-------------------------------------------------------------------------|
| M1)<br>Habitat probability | Likelihood of experiencing different habitat conditions within a patch over time (Bätz et al. 2025).                                        | $M1 = \frac{\sum t_H}{T}$ |  | Time-integrated habitat availability; identifies dominant habitat types |
| M2)<br>Habitat shifts      | Daily number of habitat changes to another suitable habitat type within a patch (Bätz et al. 2025).                                         | $M2 = \frac{\sum S}{T}$   |  | Indicates habitat stability and temporal persistence                    |
| M3) Drift risk             | Risk of macroinvertebrate drift based on current velocity and, where relevant, the up-ramping rate experienced within each patch over time. | M3 = P90 (Rdrift)         |  | Identifies exposure to critical passive drift conditions                |
| M4)<br>Desiccation risk    | Risk of egg desiccation associated with hydropeaking-induced water-level variations.                                                        | M4 = max (Rdes)           |  | Identifies exposure to critical desiccation duration                    |

tH = duration of occurrence of a specific habitat type over the time-series

T = total duration of the time-series

S = habitat-shift between two consecutive time steps

P90 = 90th percentile value over the time-series



max = maximum value over the time-series  
Rdrift = drift risk as defined in Table 6  
Rdes = desiccation risk as defined in Table 7

Drift and desiccation risks are based on threshold values that depend not only on the taxa considered, but also on the available data and the habitat type targeted by the analysis.

**Error! Reference source not found.** and **Error! Reference source not found.** present the threshold values used in the example case study.

Table 6 focuses on drift risk for two habitat types: "Slow" (habitat type 2 in the toolbox) and "Slow to intermediate" (habitat type 3 in the toolbox). For the "Slow" habitat type, the analysis considers the up-ramping rate (up\_ramp set to true in the configuration file, with a corresponding value specified in the drift ramp thresholds). The "Slow to intermediate" habitat type does not consider the up-ramping rate.

The parameters specified in the configuration file determine which drift calculation function is used by the metric calculation scripts. It is the user's responsibility to determine whether the up-ramping rate should be considered, based on the available data and the scope of the study.

*Table 6 Macroinvertebrate drift risk thresholds and corresponding metric values for the two habitat types considered in the example case study.*

| Metric value<br>( $R_{\text{drift}}$ ) | Drift risk<br>class | Habitat type                         |                         |                                      |
|----------------------------------------|---------------------|--------------------------------------|-------------------------|--------------------------------------|
|                                        |                     | Slow <sup>a</sup>                    |                         | Slow to<br>intermediate <sup>b</sup> |
|                                        |                     | Current velocity<br>threshold        | Up-ramping rate         | Current velocity<br>threshold        |
| 1                                      | Low                 | $v_{40} \leq 0.4 \text{ m/s}$        | Not relevant            | $v_{40} \leq 1.2 \text{ m/s}$        |
| 2                                      | Moderate            | $0.4 < v_{40} < 0.7 \text{ m/s}$     | Not relevant            | $1.2 < v_{40} < 2.1 \text{ m/s}$     |
|                                        |                     | $0.7 \leq v_{40} \leq 1 \text{ m/s}$ | $< 1 \text{ cm/min}$    |                                      |
| 3                                      | High                | $0.7 \leq v_{40} \leq 1 \text{ m/s}$ | $\geq 1 \text{ cm/min}$ | $2.1 \leq v_{40} \leq 3 \text{ m/s}$ |
| 4                                      | Very high           | $v_{40} > 1 \text{ m/s}$             | Not relevant            | $v_{40} > 3 \text{ m/s}$             |

<sup>a</sup>: thresholds taken from the flume experiments in Tonolla et al. (2019) and incorporating findings from Frieze et al. (2025).

<sup>b</sup>: thresholds mathematically extrapolated from the "slow" current habitat, by proportionally adjusting the upper limit of the typical current velocity range. Up-ramping rate was not included to avoid introducing additional assumptions.

**Error! Reference source not found.** presents the example of egg desiccation risk threshold values implemented in the toolbox.

*Table 7 Macroinvertebrate desiccation risk thresholds and metric values based on Kennedy et al. (2016) and Jaulin (2025)*

| Metric value ( $R_{\text{des}}$ ) | Desiccation risk class | Max. dry duration     |
|-----------------------------------|------------------------|-----------------------|
| 1                                 | Low                    | $\leq 1 \text{ hour}$ |
| 2                                 | Moderate               | 1 - 5 hours           |

|   |           |             |
|---|-----------|-------------|
| 3 | High      | 5 - 9 hours |
| 4 | Very high | > 9 hours   |

The `start_at_first_occurrence` parameter can affect the calculated desiccation risk. If set to true, desiccation risk is calculated only after the first occurrence of the target habitat type. If set to false, the calculation starts at the beginning of the discharge time series.

For example, for an analysis focusing on egg desiccation on gravel banks, setting `start_at_first_occurrence` to true means that the desiccation risk calculation starts after the first occurrence of the target habitat type considered suitable for egg deposition.

#### *Advanced statistics when focusing on a certain zone*

When `focus_on_zone` is set to true, additional statistics, such as median and quartile values, are calculated for the target habitat type and dry conditions. These statistics provide a more detailed understanding of habitat probability distributions (see Table 11 in Section 5.1 and the corresponding part of Section 5.2).

#### 4.5 Estimation of the computational time

Computational time can vary substantially depending on the spatial and temporal extent of the analysis. Examples of estimated computational times are provided in Table 8 Figure 7.

*Table 8 Estimation of computational times based on the number of patches and of timesteps included in the analysis*

| Total number of patches | Number of patches for metrics calculation | Number of timesteps | Estimated computational time |
|-------------------------|-------------------------------------------|---------------------|------------------------------|
| 30 446                  | 25 769                                    | 1008                | ~ 5 minutes                  |
| 365 000                 | 27 350                                    | 1008                | ~ 50 minutes                 |
| 664 000                 | 76 060                                    | 4393                | ~ 4 hours                    |

#### 4.6 Additional functions

*Table 9 Association between the Python script to run and the corresponding outputs*

| Output to get                    | Python file to run   |
|----------------------------------|----------------------|
| Metrics results and preview maps | <i>main.py</i>       |
| Plots                            | <i>plots.py</i>      |
| Clustering                       | <i>clustering.py</i> |

Before running the additional scripts, the paths to the files to be analyzed (`base_output_path`, `final_csv_path`, `final_shp_path`, and `static_habitat_csv_path`) must be specified in the configuration file

(see Section **Error! Reference source not found.**). The configuration file must then be saved before running the scripts.

The metrics used by the plotting and clustering functions are determined from the configuration file. The selected metrics must therefore match the metrics available in the files being analyzed. For example, if `desiccation_risk` is set to `true` in the configuration file, the corresponding `desiccation_risk` field must also be present in the input file, as the functions will search for this field.

### *Plots*

**Output location:** data > outputs > Toolbox\_run\_output > Plots

The plots focus on two geographical extents:

- **The entire simulated domain:** habitat probabilities and the most probable habitat types are assessed for all simulated patches. This provides an overview of the spatial distribution of habitats and helps explain why some patches do not experience the target habitat type during the habitat time series.
- **The metric calculation domain:** only patches that experience the target habitat type at least once during the habitat time series are included.

If the habitat classification is modified, the habitat information used by the plotting functions must also be adapted accordingly. In particular, the habitat labels and colors defined in `plots.py` need to correspond to the habitat codes used in the analysis. The appropriate label dictionary also needs to be adapted depending on whether `focus_on_zone` is set to `true` or `false`. If the habitat codes in the analysis do not match the codes defined in these dictionaries, the plotting functions may fail to generate the plots correctly.

Additional or more targeted plots can be created by modifying existing functions or adding new ones. As previously mentioned, pulling the latest version of the repository from GitHub can overwrite local modifications. Any customized code should therefore be saved in a separate, safe location before updating the repository.

Examples of outputs are presented in Section 5.2.

### *Clustering*

**Output location:** data > outputs > Toolbox\_run\_output > Clustering

The `clustering.py` script performs a k-means clustering analysis to synthesize the information contained in the four habitat metrics. The objective is to group patches with similar multimetric profiles, thereby summarizing complex combinations of macroinvertebrate habitat quality and ecological risk into a smaller number of interpretable groups.

Because each metric represents a different dimension of macroinvertebrate habitat dynamics, clustering can reveal patterns that may not be apparent when the metrics are examined individually.

The optimal number of clusters is determined using the silhouette score. The maximum number of clusters considered is set to 9 by default and can be modified in the code if needed.

The clustering functions provide information on cluster statistics and Principal Component Analysis (PCA) loadings. The cluster number assigned to each patch is also added to the CSV and shapefile containing the metric results. Further information on the output files is provided in Section 5.3**Error! Reference source not found..**

## 5. Output interpretation

**Output location:** data > output.

Relevant subfolders are used to organize the different types of outputs. The video [5 HaDy output analysis](#) provides additional guidance on how to explore and interpret the outputs.

### 5.1 Outputs from the main.py file execution

Running *main.py* generates the outputs presented in Table 10.

Table 10 Outputs generated by main.py

| Output name                                           | Output location             | Explanation                                                                                                                                                                                                                                                                                       |
|-------------------------------------------------------|-----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| discharge_with_match.csv                              | Discharge_with_match        | Associates each discharge in the input discharge time series with the closest discharge for which hydrodynamic simulation data are available.                                                                                                                                                     |
| cropped_discharge_and_wetted_area (csv and shapefile) | Cropped_shapefiles          | Contains the input data cropped to the discharges considered in the analysis and to the spatial area of interest, corresponding to patches that are wet under the maximum discharge considered. This reduces the study domain before metric calculation and therefore reduces computational time. |
| habitat_classification.csv                            | Habitat_classification      | Provides a static representation of the habitat classification for each patch and each discharge considered, before temporal integration. This file is used by the plotting functions to represent habitat availability at individual discharges.                                                 |
| metrics (csv and shapefile)                           | Metric_files                | Contains the final metric results for patches that experience the target habitat type at least once during the habitat time series.                                                                                                                                                               |
| metrics_essentials.shp                                | Metric_files                | Contains a selection of essential fields: patch ID, x-coordinate, y-coordinate, area, and the selected metrics.                                                                                                                                                                                   |
| Preview maps                                          | Metric_files > Preview_maps | Provides a visual overview of the results before opening the GIS project. A preview map is generated for each selected metric.                                                                                                                                                                    |

### Example case study Box 3: Outputs from the main.py file execution

The outputs from the example case study are located in data > output > Example\_output. The folder contains the files listed in Table 10, organized into their respective subfolders.

The extent of the zoomed area shown in the preview maps can be adapted by modifying the corresponding settings in crop\_metric\_and\_map\_preview.py. This allows the preview maps to be customized to focus on a specific area of interest.




Figure 10 Preview maps from the example case study. Panel a) shows the full reach, panel b) identifies the zoomed area, and panel c) shows the corresponding zoomed area

For the example case study, both QGIS and ArcGIS projects are provided to facilitate exploration of the metric outputs. These are located in data > output > Example\_output > QGIS\_project.

For QGIS users, an example map template (example\_case\_maps\_template.qpt) and default symbologies for the four metrics (.qml files) are available in the same folder.

For further guidance on exploring the results in QGIS, see the support video [3 HaDy example case QGIS](#).



Shapefile field names are limited to 10 characters. As a result, some field names in the output shapefile are shortened. Table 11 provides explanation.

In Table 11, “h” stands for habitat, and is followed by the corresponding habitat type number. As previously mentioned, metrics are calculated only for patches that experience the target habitat type at least once during the habitat time series.

*Table 11 Description of the fields contained in the metric output files (CSV and shapefile)*

| Shapefile column name      | Associated explanation                                                                                                                                                                                                                                                                                                                        |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Id                         | Patch ID, identical to the corresponding field in the input shapefile.                                                                                                                                                                                                                                                                        |
| x                          | X-coordinate of the patch, identical to the corresponding field in the input shapefile.                                                                                                                                                                                                                                                       |
| y                          | Y-coordinate of the patch, identical to the corresponding field in the input shapefile.                                                                                                                                                                                                                                                       |
| area                       | Patch area, identical to the corresponding field in the input shapefile.                                                                                                                                                                                                                                                                      |
| d_discharge                | Water depth for each discharge considered in the analysis, identical to the corresponding fields in the input shapefile.                                                                                                                                                                                                                      |
| v_discharge                | Current velocity for each discharge considered in the analysis, identical to the corresponding fields in the input shapefile.                                                                                                                                                                                                                 |
| focus_zone                 | Only generated when focus_on_zone is true. Indicates whether the patch is located within the zone of interest: 0 if outside and 1 if inside.                                                                                                                                                                                                  |
| id_focus                   | Only generated when focus_on_zone is true. Indicates the ID of the corresponding polygon in the zone-of-interest shapefile (e.g., the ID of an individual gravel bank).                                                                                                                                                                       |
| <b>Habitat probability</b> |                                                                                                                                                                                                                                                                                                                                               |
| dur_h                      | Duration of each habitat type during the habitat time series, expressed as a number of timesteps. The number of dur_h fields depends on the habitat classification. For example, the egg desiccation classification contains habitat types from -1 to 3, while the example community-level classification contains habitat types from 0 to 6. |
| prob_h                     | Probability of experiencing the corresponding habitat type, calculated as the duration of that habitat type divided by the total duration of the habitat time series.                                                                                                                                                                         |
| mostProb                   | Most probable habitat type for each patch over the habitat time series, corresponding to the habitat type with the highest probability. This is calculated for all patches in the simulated domain, not only those experiencing the target habitat type (see <b>Error! Reference source not found.</b> )                                      |

|                                                                                                   |                                                                                                                                                                                                                                                                                                                         |
|---------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| h_first                                                                                           | Timestep of the first occurrence of the corresponding habitat type. For example, h2_first gives the timestep of the first occurrence of habitat type 2 for a given patch.                                                                                                                                               |
| <b>Statistics on the target habitat type – advanced analysis focusing on a specific zone only</b> |                                                                                                                                                                                                                                                                                                                         |
| h_nb_seq                                                                                          | Number of separate sequences during which the target habitat type occurs in the patch over the habitat time series.                                                                                                                                                                                                     |
| h_dur_max                                                                                         | Maximum consecutive duration of the target habitat type in the patch, expressed in hours.                                                                                                                                                                                                                               |
| h_dur_med                                                                                         | Median consecutive duration of the target habitat type in the patch, expressed in hours.                                                                                                                                                                                                                                |
| h_dur_q1                                                                                          | First quartile of the distribution of consecutive durations of the target habitat type, expressed in hours.                                                                                                                                                                                                             |
| h_dur_q3                                                                                          | Third quartile of the distribution of consecutive durations of the target habitat type, expressed in hours.                                                                                                                                                                                                             |
| <b>Habitat shifts</b>                                                                             |                                                                                                                                                                                                                                                                                                                         |
| h_sh_all                                                                                          | Total number of daily habitat shifts experienced by the patch over the habitat time series.                                                                                                                                                                                                                             |
| h_sh_suit                                                                                         | If not focusing on a zone: total number of daily shifts to the suitable habitat types specified in the configuration file. If focusing on a zone: total number of daily shifts to the target habitat type.                                                                                                              |
| h_sh_dry                                                                                          | Total number of daily shifts to dry conditions over the habitat time series.                                                                                                                                                                                                                                            |
| <b>Desiccation</b>                                                                                |                                                                                                                                                                                                                                                                                                                         |
| h_dryMax                                                                                          | Maximum consecutive duration of dry conditions, expressed as a number of timesteps. The starting point of the calculation depends on start_at_first_occurrence: if false, the calculation starts at the beginning of the habitat time series; if true, it starts after the first occurrence of the target habitat type. |
| h_desicR                                                                                          | Associated desiccation risk value.                                                                                                                                                                                                                                                                                      |
| <b>Desiccation additional statistics – advanced analyses focusing on a specific zone only</b>     |                                                                                                                                                                                                                                                                                                                         |
| h_dry_seq                                                                                         | Number of separate sequences of dry conditions in the patch over the habitat time series.                                                                                                                                                                                                                               |
| h_dry_max                                                                                         | Maximum consecutive duration of dry conditions in the patch, expressed in hours.                                                                                                                                                                                                                                        |
| h_dry_med                                                                                         | Median consecutive duration of dry conditions in the patch, expressed in hours.                                                                                                                                                                                                                                         |
| h_dry_q1                                                                                          | First quartile of the distribution of consecutive dry durations, expressed in hours.                                                                                                                                                                                                                                    |
| h_dry_q3                                                                                          | Third quartile of the distribution of consecutive dry durations, expressed in hours.                                                                                                                                                                                                                                    |
| h_dryMedR                                                                                         | Risk value associated with the median consecutive duration of dry conditions.                                                                                                                                                                                                                                           |



|              |                                                                                                 |
|--------------|-------------------------------------------------------------------------------------------------|
| h_dry_q1R    | Risk value associated with the first quartile of the distribution of consecutive dry durations. |
| h_dry_q3R    | Risk value associated with the third quartile of the distribution of consecutive dry durations. |
| <b>Drift</b> |                                                                                                 |
| h_dd_Risk    | Duration spent in each drift risk category, expressed as a number of timesteps.                 |
| h_driftP     | Drift risk percentile.                                                                          |
| h_driftM     | Maximum drift risk value.                                                                       |

## 5.2 Outputs from the plots.py execution

The *plots.py* script generates different outputs depending on the parameters selected in the configuration file (see Section 4.6, Plots).

**Output location:** data > output > Toolbox\_run\_output > Plots

Additional guidance on interpreting the outputs is provided in the video [5 HaDy output analysis](#).

## Example case study Box 4: Outputs from the plots.py file execution

### Static representation of habitat distribution per discharge

These figures show how habitat types are distributed across the study area at each simulated discharge.


The **x-axis** represents the different simulated discharges. For each discharge, the figures show the spatial distribution of habitat conditions within the selected spatial coverage.

**Rows:** Because patches do not all have the same area, habitat distributions are presented both:

- By **number of patches**, and
- By **total area** occupied by each habitat type.

**Columns:** Two spatial coverages are considered:

- **Entire simulated domain:** all patches that are wet under the maximum simulated discharge.
- **Target-habitat patches only:** only patches that experience the selected target habitat at least once during the habitat time series.



### Most probable habitat type over the habitat time-series

For each patch, the **most probable habitat type** is the habitat type that occurs most frequently throughout the habitat time series.

The probability of each habitat type is calculated as:

Time spent in that habitat type ÷ Total duration of the habitat time series

These figures summarize the dominant habitat conditions by integrating habitat variability over both **space** and **time**.

**Columns:** The two spatial coverages are identical to those described above:

- The entire simulated domain, and
- Only patches that experience the target habitat during the habitat time series.

|                                                                                                                                                                                                                                                        | Entire simulated domain                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Target-habitat patches only                                                                                                                                                                                                                                                                                                                                      |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Most probable habitat type distribution (number of patches)                                                                                                                                                                                            | <p>This figure shows which habitat type is the dominant one for each patch over the habitat time series.</p> <p>For example, the habitat type "<b>fast to very fast</b>" is the most probable habitat type for <b>62.8%</b> of the patches in the simulated study area, indicating that this habitat condition dominates most locations over time.</p>                                                                                                                                                                                                                                                                                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                                  |
| Probability associated with the most probable habitat type                                                                                                                                                                                             | <p>Knowing the dominant habitat type alone does not indicate how strongly that habitat dominates a patch. This figure therefore shows the probability associated with each patch's most probable habitat type.</p> <p>For example, although "<b>fast to very fast</b>" is the dominant habitat for many patches, its associated median probability is approximately <b>0.75</b>. This means that these patches spend about <b>75% of the habitat time series</b> under "fast to very fast" conditions.</p> <p>In this example, such high probabilities are consistent with the hydropeaking discharge regime, indicating that a large proportion of the riverbed experiences fast-flowing conditions for most of the simulated period.</p> |                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Metrics representations</b> <p>The following figures are computed only for patches that experience the target habitat at least once during the habitat time series (target-habitat patches). See Section 4.4 details on the metrics calculation</p> |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                                  |
|                                                                                                                                                                                                                                                        | Target-habitat patches only                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | Explanation                                                                                                                                                                                                                                                                                                                                                      |
| Habitat probability of the target habitat type                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | <p>This shows the probability that each patch is in the target habitat over the habitat time-series.</p> <p>In this example, the target habitat is "<b>slow to fast</b>." Although the selected patches all experience this habitat at least once, most spend only a limited amount of time in it: <b>87.8%</b> of the patches spend less than <b>50%</b> of</p> |

|                                   |                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|-----------------------------------|--------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                   |                                                                                                                                | the habitat time series under target habitat conditions.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Habitat probability distributions | <p>Distribution of habitat probabilities (patches experiencing habitat target 3 at least once during the flow time-series)</p> | <p>This presents the probability distributions of <b>all habitat types</b> for the target-habitat patches.</p> <p>It complements the previous figure by showing how the remaining time is distributed among the other habitat types.</p> <p>In this example, the selected patches spend most of their time under "<b>fast to very fast</b>" conditions, confirming the dominance of this habitat observed in the previous analyses.</p>                                                                                                                                    |
| Daily shifts to suitable habitats | <p>Percentage of patches</p> <p>Habitat shifts</p>                                                                             | <p>This shows the number of <b>daily transitions into suitable habitat conditions</b> over the habitat time series.</p> <p>Suitable habitat types are specified by the user in the configuration file. In this example, the suitable habitats are:</p> <ul style="list-style-type: none"> <li>• Stagnant</li> <li>• Slow</li> <li>• Slow to fast</li> <li>• Fast to very fast</li> <li>• Very fast</li> </ul> <p>A shift is counted each time a patch changes from an <b>unsuitable habitat</b> (dry or violent current) to any of the defined suitable habitat types.</p> |
| Desiccation risk                  | <p>Percentage of patches</p> <p>Desiccation risk</p>                                                                           | <p>See Section 4.4 for details on the risk calculation.</p> <p>This figure shows the proportion of patches falling into each desiccation risk category, providing an overview of the spatial distribution of desiccation risk within the selected patches.</p>                                                                                                                                                                                                                                                                                                             |
| Drift risk                        | <p>Percentage of patches</p> <p>Drift risk</p>                                                                                 | <p>See Section 4.4 for details on the risk calculation.</p> <p>This figure shows the proportion of patches falling into each drift risk category, providing an overview of the spatial distribution of desiccation risk within the selected patches.</p>                                                                                                                                                                                                                                                                                                                   |

The available plotting functions are described in Section 4.6. Users can freely modify the plotting script to adapt the figures to their specific needs, for example by customizing existing plots or adding or removing figures.

To prevent these modifications from being overwritten when updating the toolbox from GitHub, any customized version of plots.py should be saved under a different filename or in a separate location.



The generated figures depend on the parameters defined in the local.yaml configuration file. When patches have different sizes, some plots can be normalized by patch area instead of by number of patches, depending on the selected settings.







**Advanced plots when focusing on a specific zone**





For more advanced analyses, additional plots are available when the focus\_on\_zone Boolean parameter is set to True. These outputs are presented in Table 12.

For example, when comparing different gravel banks, each patch must be assigned a zone ID indicating the gravel bank to which it belongs. The toolbox then generates plots separately for each zone, allowing direct comparison of habitat dynamics and associated metrics between gravel banks. Further information in the video [6 HaDy advanced focus on zone](#).

Table 12 Advanced plots generated by plot.py when focusing on a specific zone

| Target habitat: additional statistics                                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                            |                       |        |        |       |       |        |         |         |        |
|-----------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------|-----------------------|--------|--------|-------|-------|--------|---------|---------|--------|
| Number of individual sequences during which the target habitat occurs over the habitat time series. |  <table border="1"> <caption>Data for Figure 1: Percentage of patches by number of habitat 2 windows</caption> <thead> <tr> <th>Number of habitat 2 window</th> <th>Percentage of patches</th> </tr> </thead> <tbody> <tr> <td>0-35</td> <td>5.0%</td> </tr> <tr> <td>36-72</td> <td>58.6%</td> </tr> <tr> <td>73-108</td> <td>35.8%</td> </tr> <tr> <td>109-145</td> <td>~35.8%</td> </tr> </tbody> </table> | Number of habitat 2 window | Percentage of patches | 0-35   | 5.0%   | 36-72 | 58.6% | 73-108 | 35.8%   | 109-145 | ~35.8% |
| Number of habitat 2 window                                                                          | Percentage of patches                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                            |                       |        |        |       |       |        |         |         |        |
| 0-35                                                                                                | 5.0%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                            |                       |        |        |       |       |        |         |         |        |
| 36-72                                                                                               | 58.6%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                            |                       |        |        |       |       |        |         |         |        |
| 73-108                                                                                              | 35.8%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                            |                       |        |        |       |       |        |         |         |        |
| 109-145                                                                                             | ~35.8%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                            |                       |        |        |       |       |        |         |         |        |
| Distribution of the durations of target-habitat sequences, expressed in hours.                      |  <table border="1"> <caption>Data for Figure 2: Distribution of habitat 2 window durations</caption> <thead> <tr> <th>Statistic</th> <th>Approximate Range (h)</th> </tr> </thead> <tbody> <tr> <td>Median</td> <td>5 - 10</td> </tr> <tr> <td>Q1</td> <td>2 - 5</td> </tr> <tr> <td>Q3</td> <td>10 - 22</td> </tr> </tbody> </table>                                                                         | Statistic                  | Approximate Range (h) | Median | 5 - 10 | Q1    | 2 - 5 | Q3     | 10 - 22 |         |        |
| Statistic                                                                                           | Approximate Range (h)                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                            |                       |        |        |       |       |        |         |         |        |
| Median                                                                                              | 5 - 10                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                            |                       |        |        |       |       |        |         |         |        |
| Q1                                                                                                  | 2 - 5                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                            |                       |        |        |       |       |        |         |         |        |
| Q3                                                                                                  | 10 - 22                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                            |                       |        |        |       |       |        |         |         |        |
| Desiccation risk and dry durations                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                            |                       |        |        |       |       |        |         |         |        |
| Desiccation risk                                                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                            |                       |        |        |       |       |        |         |         |        |

|                                                                                   |                                                                                                                                                                                                             |
|-----------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>Distribution of desiccation risk values across all zones.</p>                  | <p>Desiccation risk distribution across patches (habitat 2, derived from Max / Median / Q1 / Q3 dry window duration)</p>  |
| <p>Median desiccation risk calculated for each gravel bank.</p>                   | <p>Desiccation risk (median window) per gravel bank</p>                                                                   |
| <p>Desiccation risk values calculated for each gravel bank.</p>                   | <p>Desiccation risk (DryMax) per gravel bank</p>                                                                          |
| <p>First quartile of desiccation risk values calculated for each gravel bank.</p> | <p>Desiccation risk (Q1 window) per gravel bank</p>                                                                      |
| <p>Third quartile of desiccation risk values calculated for each gravel bank.</p> | <p>Desiccation risk (Q3 window) per gravel bank</p>                                                                     |
| <p><b>Dry durations</b></p>                                                       |                                                                                                                                                                                                             |
| <p>Distribution of dry window durations across all zones.</p>                     | <p>Distribution of dry window durations across patches (habitat 2)</p>                                                   |

| Number of separate dry periods identified across all zones.               |  <p>Percentage of patches</p> <p>Number of dry windows</p> <table border="1"> <thead> <tr> <th>Number of dry windows</th> <th>Percentage of patches</th> </tr> </thead> <tbody> <tr> <td>0-16</td> <td>3.1%</td> </tr> <tr> <td>17-33</td> <td>59.4%</td> </tr> <tr> <td>34-50</td> <td>37.5%</td> </tr> <tr> <td>51-67</td> <td>0%</td> </tr> </tbody> </table> | Number of dry windows | Percentage of patches | 0-16 | 3.1% | 17-33 | 59.4% | 34-50 | 37.5% | 51-67 | 0% |
|---------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|-----------------------|------|------|-------|-------|-------|-------|-------|----|
| Number of dry windows                                                     | Percentage of patches                                                                                                                                                                                                                                                                                                                                                                                                                             |                       |                       |      |      |       |       |       |       |       |    |
| 0-16                                                                      | 3.1%                                                                                                                                                                                                                                                                                                                                                                                                                                              |                       |                       |      |      |       |       |       |       |       |    |
| 17-33                                                                     | 59.4%                                                                                                                                                                                                                                                                                                                                                                                                                                             |                       |                       |      |      |       |       |       |       |       |    |
| 34-50                                                                     | 37.5%                                                                                                                                                                                                                                                                                                                                                                                                                                             |                       |                       |      |      |       |       |       |       |       |    |
| 51-67                                                                     | 0%                                                                                                                                                                                                                                                                                                                                                                                                                                                |                       |                       |      |      |       |       |       |       |       |    |
| Median duration of dry periods for each gravel bank, expressed in hours.  |  <p>Median dry window duration (h) per gravel bank</p> <p>Median dry window duration (h)</p> <p>Gravel bank</p>                                                                                                                                                                                                                                                 |                       |                       |      |      |       |       |       |       |       |    |
| Maximum duration of dry periods for each gravel bank, expressed in hours. |  <p>Max dry window duration (h) per gravel bank</p> <p>Max dry window duration (h)</p> <p>Gravel bank</p>                                                                                                                                                                                                                                                       |                       |                       |      |      |       |       |       |       |       |    |
| Number of separate dry periods identified for each gravel bank.           |  <p>Number of dry windows per gravel bank</p> <p>Number of dry windows</p> <p>Gravel bank</p>                                                                                                                                                                                                                                                                  |                       |                       |      |      |       |       |       |       |       |    |


### 5.3 Outputs from the clustering.py execution

The *clustering.py* script generates different outputs depending on the parameters selected in the configuration file (see Section 4.6, Clustering).

**Output location:** data > output > Toolbox\_run\_output > Clustering

Additional guidance on interpreting the outputs is provided in the video [5 HaDy output analysis](#).

#### Example case study Box 5: Outputs from the clustering.py file execution

| Output                                       | Description                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
|----------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|-------|---|------|---|------|---|------|---|------|---|------|---|------|---|------|---|------|
| Hab_silhouette.png                           | <p>Plot showing the silhouette score for different numbers of clusters. The number of clusters corresponding to the highest silhouette score is selected as the optimal number of clusters.</p>  <table border="1"> <caption>Silhouette Score Data</caption> <thead> <tr> <th>Number of clusters</th> <th>Score</th> </tr> </thead> <tbody> <tr><td>2</td><td>0.48</td></tr> <tr><td>3</td><td>0.49</td></tr> <tr><td>4</td><td>0.37</td></tr> <tr><td>5</td><td>0.39</td></tr> <tr><td>6</td><td>0.41</td></tr> <tr><td>7</td><td>0.39</td></tr> <tr><td>8</td><td>0.41</td></tr> <tr><td>9</td><td>0.36</td></tr> </tbody> </table> | Number of clusters | Score | 2 | 0.48 | 3 | 0.49 | 4 | 0.37 | 5 | 0.39 | 6 | 0.41 | 7 | 0.39 | 8 | 0.41 | 9 | 0.36 |
| Number of clusters                           | Score                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 2                                            | 0.48                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 3                                            | 0.49                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 4                                            | 0.37                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 5                                            | 0.39                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 6                                            | 0.41                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 7                                            | 0.39                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 8                                            | 0.41                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| 9                                            | 0.36                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| Hab_PCA_clusters.png                         | <p>Plot showing the position of the identified clusters in the PC1–PC2 space.</p>  <p>PC1–PC2 space.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| Hab_PCA_loadings.csv                         | CSV file containing the PC1 and PC2 loadings for each metric.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |
| metrics_with_clusters<br>(csv and shapefile) | <p>Metric output files supplemented with clustering results. Additional fields include Cluster_hab, indicating the cluster assigned to each patch, and PC1_hab and PC2_hab, containing the coordinates of each patch in the first two principal component dimensions.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                |                    |       |   |      |   |      |   |      |   |      |   |      |   |      |   |      |   |      |



## 6. Limitations and assumptions

The following limitations and assumptions should be considered when interpreting the toolbox outputs:

- **Input hydrodynamic data:** The toolbox assumes that the user has identified the discharges relevant to the study and performed the corresponding hydrodynamic simulations. The resulting water depth and current velocity data must be provided in the required input shapefile format.
- **Habitat classification and metric thresholds:** The toolbox provides example habitat classification thresholds and desiccation and drift risk thresholds. These values are not universally applicable and must be adapted to the objectives of the study, the habitat type considered, and the macroinvertebrate taxa of interest.
- **GIS analysis:** The toolbox provides output shapefiles and preview maps, but the detailed spatial exploration and interpretation of the results are performed by the user in a GIS software.
- **Discharge representation:** The toolbox assigns each discharge in the input time series to the closest discharge for which hydrodynamic simulation data are available. No interpolation is performed between simulated discharges. The representativeness and resolution of the selected simulated discharges therefore directly affect the resulting habitat time series and metrics.
- **Temporal resolution:** Metric calculations depend on the timestep of the input discharge time series. The timestep must accurately represent the temporal resolution required for the study.
- **Threshold-based risk assessment:** Drift and desiccation risks are estimated from predefined threshold values. These metrics should therefore be interpreted as relative or threshold-based indicators of exposure rather than direct measurements of demographic or population-level effects.

## 7. Trouble shooting

The following recommendations can help become familiar with the toolbox and avoid common issues:

- **Start small, in space and time:** Begin with a small spatial study area and a short discharge time series. This reduces computational time and makes it easier to check whether the results are consistent and plausible. To crop the original shapefile to a smaller study area, the QGIS procedure is described in Section 9.2
- **Save scripts and analysis files separately:** Pulling the latest version of the GitHub repository may overwrite locally modified scripts and files. Save any customized scripts, configuration files, and analysis results in a separate, safe location before updating the repository.
- **Follow the input formatting requirements carefully:** The format of the input data, particularly the column names in the input shapefile, is important because the toolbox extracts information from these names. For example, a column representing a discharge of 18.3 m<sup>3</sup>/s must follow the required format, such as depth\_18\_3, rather than depth\_18.3. If the required format is not respected, the toolbox may fail to run and no outputs will be generated. Column names can be modified using standard GIS software such as QGIS.
- **Check the local.yaml configuration file:** If no local.yaml file is created, the toolbox uses config.yaml by default, which contains the parameters of the example case study. For a tailored analysis, create and adapt local.yaml as described in Section 3.2, and save the file before running the toolbox.
- **Check that the configuration matches the input data:** Ensure that the parameters entered in local.yaml correspond to the actual input files. In particular, check the input filenames, column names and prefixes, discharge timestep, habitat classification, number of habitats, and selected metrics.
- **Test each modification progressively:** When adapting the toolbox, changing one parameter or function at a time makes it easier to identify the source of an issue. After each modification, save the relevant file before running the toolbox and check whether the resulting outputs are as expected.

## 8. Glossary

*Patch*: A spatially fixed unit within a river reach, typically corresponding to areas on the order of 0.25 to 100 m<sup>2</sup> (micro- to meso-scale). The appropriate patch size depends on the species and life stage considered, as well as their mobility. While the location of a patch remains constant over time, the habitat conditions within it may vary. In this toolbox, patches usually correspond to the mesh cells of the hydrodynamic simulation for which water depth and current velocity data are available.

*Habitat*: A set of environmental conditions characterized by dominant physical and chemical properties (e.g., current velocity, water depth, and substrate). These conditions can be described at different spatial scales, including the micro- to meso-scale within a river reach. Different species and life stages rely on specific habitat conditions to fulfil essential functions during their life cycle, such as feeding, reproduction, resting, or refuge seeking. The spatial distribution of habitats within a river reach can change over time.

*Habitat shift (M2)*: A transition from one habitat type to another resulting from changes in environmental conditions over time and/or space. In this toolbox, habitat shifts are assessed at the patch scale, capturing temporal changes in habitat conditions within a fixed spatial unit (i.e., the patch).

*Discharge time series*: The temporal evolution of river discharge (m<sup>3</sup>/s), represented at a defined timestep (e.g., minutes or hours between two consecutive discharge values).

*Habitat time series*: The temporal sequence of habitat conditions within each patch, obtained by combining the spatial habitat classification with the discharge time series.

*Target habitat*: The habitat type that is the primary focus of the analysis. Habitat types are defined according to a chosen habitat classification scheme, with each type assigned a numerical code for processing within the toolbox. Metrics are calculated specifically for patches that experience the target habitat at least once during the habitat time series.

*Suitable habitats*: Habitat types considered favorable for the macroinvertebrate taxa under study, based on the criteria defined in the selected habitat classification scheme.

*Habitat probability (M1)*: A patch-scale metric representing the proportion of the habitat time series during which a given habitat type occurs within a patch. It quantifies the temporal availability of habitat conditions within the patch and can be used to identify dominant habitat types.

*Habitat shifts (M2)*: A patch-scale metric representing the number of daily transitions between habitat conditions within a patch. Depending on the selected configuration, shifts can be calculated between all habitat types, between suitable habitat types, or towards dry conditions. The metric provides an indication of habitat stability and temporal persistence.

*Drift risk (M3):* A patch-scale metric representing exposure to hydraulically induced drift. It is calculated from the drift-risk values obtained for the individual timesteps of the habitat time series. The default metric is the 90th percentile of these values, thereby representing frequent exposure to elevated drift conditions while reducing the influence of isolated extreme events.

*Desiccation risk (M4):* A patch-scale metric representing exposure to drying conditions over time. It is derived from the habitat time series and reflects the occurrence and persistence of conditions in which the patch becomes dry due to insufficient water depth, indicating potential ecological stress for macroinvertebrates.

## 9. QGIS tips and tricks

### 9.1 Analyzing a shapefile value field on QGIS

The output shapefiles contain several attribute fields, corresponding to patch characteristics and calculated habitat metrics. An attribute field is a column in the shapefile attribute table containing a specific variable or metric for each patch.

To visualize a specific metric spatially, the corresponding attribute field can be selected as the basis for the layer symbology. Figure 11 provides an example of how to do this in QGIS.

For the example case study, a QGIS project and a map template are provided. The QGIS project can be opened directly to explore the example outputs with the predefined layers and symbologies.




Figure 11 Example of how to visualize an output metric in QGIS

Default symbologies for the four metrics are also provided in: data > output > Example\_output > GIS\_project

The .qml files can be loaded in QGIS to apply the predefined symbology to the corresponding metric as follows:

1. Open the output shapefile in QGIS.
2. In the Layers panel, right-click the layer and select Properties.
3. Open the Symbology tab.

4. Select Style → Load Style.
5. Select the relevant .qml file from data > output > Example\_output > GIS\_project.
6. Click Apply and OK.

The example map template (.qpt) can be used to reproduce the layout of the example maps with other output layers. The template can be opened from QGIS using Project → New Print Layout from Template and then adapted to the desired study area, layers, and metrics.

## 9.2 Crop a shapefile to a specific area in QGIS

Cropping the input shapefile allows the creation of a reduced shapefile containing only the data within the area of interest. This can be useful when testing the toolbox on a smaller spatial extent, thereby reducing computational time.

### Step 1. Prepare the layers

Two layers are required:

- the input shapefile to be cropped;
- a polygon mask defining the area of interest.
- 

### Step 2. Use the “Clip” tool

1. Open the Processing Toolbox from Processing → Toolbox.
2. Search for Clip.
3. Select Vector overlay → Clip.
4. In the dialog:
  - Input layer: select the shapefile to be cropped;
  - Overlay layer: select the polygon defining the area of interest;
  - Clipped: choose the location and name of the new shapefile.
5. Click Run.

The resulting shapefile contains only the features located within the mask polygon.

### Step 3. Features with missing attribute values

The Clip tool can process features even when some attribute fields contain missing values. Missing attribute values therefore do not prevent the cropping operation.

However, missing values may affect subsequent analyses. If the resulting shapefile is subsequently used for statistical analyses, joins, or toolbox processing, the relevant fields should be checked and missing values handled according to the requirements described in Section 3.1.

#### **Step 4. Optional: use “Extract by location”**

If only features fully contained within the mask polygon should be retained, Extract by location can be used instead:

1. Open Vector → Research Tools → Extract by Location.
2. Select the input shapefile as the input layer.
3. Select the polygon mask as the comparison layer.
4. Select the within spatial relationship.
5. Run the tool.

This avoids retaining features that only partially overlap the area of interest.

### **9.3 Joining layers on QGIS**

This procedure can be used when two input shapefiles need to be combined into a single shapefile. For example, one shapefile may contain the mesh cells with water depth and current velocity attributes, while another contains only the polygon geometry of the patches.

The desired output is a single polygon shapefile containing both the patch geometry and the associated attributes.

The two shapefiles should have a common identifier that can be used to establish the correspondence between features. If no common identifier is available, the layers can instead be joined based on their spatial location, provided that the features occupy corresponding locations.

#### **Joining based on a common identifier**

If both shapefiles contain the same unique patch identifier:

1. Open both shapefiles in QGIS.
2. Open the Processing Toolbox.
3. Search for Join attributes by field value.
4. Select the shapefile containing the patch geometry as the target layer.
5. Select the corresponding shapefile as the layer to join.
6. Select the common identifier field in both layers.
7. Select the output location and run the tool.

#### **Joining based on spatial location**

If the two shapefiles contain corresponding features at the same locations:

1. Select both layers in QGIS.

2. Create a spatial index for each layer if required:
3. Layer → Source → Geometry → Create Spatial Index.
4. Open the Processing Toolbox.
5. Search for Join attributes by location.
6. Select the polygon shapefile as the target layer.
7. Select the layer containing the attributes to be joined.
8. Select the appropriate spatial relationship.
9. Run the tool.

The resulting polygon shapefile contains the geometry of the target layer together with the attributes from the second shapefile.



## 10. Appendix

### 10.1 Technical setup support: prerequisites for using the toolbox

#### *Installation of the core tools*

The following tools are only recommendations. If you are used to working with python, conda virtual environments and GitHub, skip this part and use your usual tools.

#### Step 1: Download and install python

Get the latest version (minimum Python 3) from the official website: [Download Python | Python.org](https://www.python.org/downloads/), and follow the instructions.

| Windows                                                                                                                                                    | macOS                                                                                          | Linux (Ubuntu/Debian)                                                                                                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <a href="https://www.python.org/downloads/windows/">Python Releases for Windows   Python.org</a><br>Download install manager<br>Check “Add Python to PATH” | <a href="https://www.python.org/downloads/mac-osx/">Python Releases for macOS   Python.org</a> | <a href="https://www.python.org/downloads/source/">Python Source Releases   Python.org</a><br><br><a href="https://www.geeksforgeeks.org/how-to-install-python-on-linux/">How to Install Python on Linux - GeeksforGeeks</a> |

#### Step 2: Install Git

From the official website: [Git](https://git-scm.com/), and follow the instructions.

| Windows                                                                                                                                             | macOS                                                                  | Linux (Ubuntu/Debian)                                                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------|--------------------------------------------------------------------------|
| <a href="https://git-scm.com/download/win">Git - Install for Windows</a><br><a href="https://git-scm.com/download/win">Keep all default options</a> | <a href="https://git-scm.com/download/mac">Git - Install for macOS</a> | <a href="https://git-scm.com/download/linux">Git - Install for Linux</a> |


#### Step 3: Install anaconda navigator

From the official software: [Installing Navigator - Anaconda](https://anaconda.com/anaconda-navigator/), and follow the instructions.

#### Step 4: Install Visual Studio code and its python extension

From the official website: [Download Visual Studio Code - Mac, Linux, Windows](https://code.visualstudio.com/download)


For Microsoft users only, install the python extension: open VS Code and follow instructions.



### Basic commands: accessing a folder on a terminal

To open a terminal, open the search bar, type in "terminal", and open it (see below for Windows).

Then type in the command `cd` and then your folder absolute path.



## 10.2 Description of files and functions

### Main.py

The **main.py** file collects functions from other python files and variables entered by the user in the configuration file and executes the code.

| Description of the workflow (in chronological order)        | Name of the function | Where the function is located | Name of the output |
|-------------------------------------------------------------|----------------------|-------------------------------|--------------------|
| Getting the known discharge values from the column names of | get_discharge_values | support_functions.py          | discharge_values   |

|                                                                                                                                                      |                                                  |                                             |                                       |
|------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------|---------------------------------------------|---------------------------------------|
| the input polygon shapefile                                                                                                                          |                                                  |                                             |                                       |
| Match the discharge of the discharge time-series to the closest known one in discharge_values                                                        | match_closest_discharge                          | <i>discharge_time_series.py</i>             | discharge_with_match.csv              |
| Select relevant discharges for the analysis: the list of corresponding known discharges                                                              | get_study_discharges                             | <i>discharge_time_series.py</i>             | relevant_discharges                   |
| Crop the shapefile: to the corresponding relevant discharges; and keeping only the wetted mask (patches wet for the max considered discharge)        | prepare_wetted_shapefile_for_relevant_discharges | <i>support_functions.py</i>                 | cropped_discharge_and_wetted_area.shp |
| Export the cropped shapefile to a csv                                                                                                                | prepare_csv                                      | <i>habitat_classification.py</i>            | cropped_discharge_and_wetted_area.csv |
| If focus_on_zone is true                                                                                                                             |                                                  |                                             |                                       |
| Attribute habitat only for patches in the focus zone (otherwise -1 by default). This habitat classification function needs to be adapted by the user | attribute_habitat_types_zone_only                | <i>habitat_attribution.py</i>               | habitat_classification.csv            |
| Metrics calculation                                                                                                                                  | process_mesh_data_focus_on_zone                  | <i>metrics_calculation_focus_on_zone.py</i> | metrics.csv                           |
| Save the metrics to a shapefile                                                                                                                      | join_mesh_with_CSV_data                          | <i>support_functions.py</i>                 | metrics.shp                           |
| If focus_on_zone is false                                                                                                                            |                                                  |                                             |                                       |
| Attribute habitats based on the example habitat classification                                                                                       | attribute_habitat_current_based                  | <i>habitat_classification.py</i>            | habitat_classification.csv            |
| Metrics calculation                                                                                                                                  | process_mesh_data                                | <i>metrics_calculation.py</i>               | metrics.csv                           |
| Save the metrics to a shapefile                                                                                                                      | join_mesh_with_CSV_data                          | <i>support_functions.py</i>                 | metrics.shp                           |

### *Crop\_metric\_and\_map\_preview.py*

The *crop\_metric\_and\_map\_preview.py* script provides a quick visual overview of the metric results by cropping the metric shapefile to a user-defined area and generating preview maps. It is intended to facilitate the initial exploration of the spatial distribution of the calculated metrics before opening the complete dataset in a GIS software.

The zoomed area can be defined manually by entering the coordinates of the desired extent directly in the relevant function in the script. This allows the user to select a specific area of interest independently of the automatically generated preview extent. The coordinates can be adapted according to the study area and the spatial feature to be examined.

The script can be modified by the user to change the selected area, the metrics displayed, or the appearance of the preview maps.

| Description of the workflow (in chronological order)     | Name of the function         | Name of the output                   |
|----------------------------------------------------------|------------------------------|--------------------------------------|
| Crop the metric shapefile to a specific area of interest | <i>crop_metric_shapefile</i> | Cropped metric shapefile             |
| Generate preview maps for the selected metrics           | <i>map_preview</i>           | Preview maps of the selected metrics |

### *Plot.py*

The ***plot.py*** contains the function for the different plots. The user can always add and/or modify these functions (more information in Section 4.6 and Section 5.2 **Error! Reference source not found.**).

| Description of the workflow (in chronological order)                                     | Name of the function                         | Name of the output                                         |
|------------------------------------------------------------------------------------------|----------------------------------------------|------------------------------------------------------------|
| ALL PATCHES IN SIMULATED DOMAIN<br>Plot most probable habitat type and related intensity | <i>plot_most_prob_intensity</i>              | hist_most_prob_and_associated_intensity_simulated_zone.png |
| TARGET HABITAT<br>Plot most probable habitat type and related intensity                  | <i>plot_most_prob_intensity_target_hab</i>   | hist_most_prob_and_associated_intensity_target_habitat.png |
| ALL PATCHES IN SIMULATED DOMAIN<br>Percentage of most probable habitat types             | <i>plot_most_prob_percentages</i>            | most_prob_percentages_simulated_zone.png                   |
| TARGET HABITAT<br>Percentage of most probable habitat types                              | <i>plot_most_prob_percentages_target_hab</i> | most_prob_percentages_target_hab.png                       |

|                                                                                                                                                                                |                                                                |                                                                                                                                                                             |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ALL PATCHES IN SIMULATED DOMAIN<br>Horizontal stacked bar of dominant habitat types                                                                                            | <i>plot_most_prob_horizontal</i>                               | most_prob_horizontal_simulated_zone.png                                                                                                                                     |
| TARGET HABITAT<br>Horizontal stacked bar of dominant habitat types                                                                                                             | <i>plot_most_prob_horizontal_target_hab</i>                    | most_prob_horizontal_target_hab.png                                                                                                                                         |
| ALL PATCHES IN SIMULATED DOMAIN<br>IN PERCENTAGE OF PATCHES -<br>Habitat availability per discharge (stacked %)                                                                | <i>plot_habitat_availability_per_discharge</i>                 | habitat_availability_per_discharge_simulated_domain.png                                                                                                                     |
| TARGET HABITAT<br>IN PERCENTAGE OF PATCHES -<br>Habitat availability per discharge (stacked %)                                                                                 | <i>plot_habitat_availability_per_discharge_target_hab</i>      | habitat_availability_per_discharge_target_hab.png                                                                                                                           |
| FUNCTION: ALL PATCHES IN SIMULATED DOMAIN<br>IN AREA –<br>Habitat availability per discharge (stacked %)                                                                       | <i>plot_habitat_availability_per_discharge_area</i>            | habitat_availability_per_discharge_simulated_domain_area.png                                                                                                                |
| TARGET HABITAT<br>IN AREA –<br>Habitat availability per discharge (stacked %)                                                                                                  | <i>plot_habitat_availability_per_discharge_area_target_hab</i> | habitat_availability_per_discharge_simulated_domain_target_hab_area.png                                                                                                     |
| TARGET HABITAT<br>Habitat probability for each habitat type<br>(Among all patches that experience the target habitat, what is the distribution of time spent in each habitat?) | <i>plot_probability_distribution_target_hab</i>                | habitat_probabilities_target_hab.png                                                                                                                                        |
| TARGET HABITAT<br>Plot percentage histograms for selected metrics of a target habitat                                                                                          | <i>plot_histograms_target_habitat</i>                          | histogram_prob_percentage.png<br>histogram_shifts_percentages.png<br>histogram_desicc_percentages.png<br>histogram_drift_percentages.png<br>boxplot_dry_window_duration.png |

|                                                                                                                                                       |                                    |                                                                                                                                                                                                                                                                                |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                                                                       |                                    | boxplot_targ_window_duration.png                                                                                                                                                                                                                                               |
| TARGET HABITAT<br>Compare desiccation metrics across gravel banks (id_focus) using Kruskal-Wallis and Dunn post-hoc tests, with boxplots per polygon. | <i>plot_desiccation_by_polygon</i> | desicc_max_h_by_polygon.png<br>desicc_median_h_by_polygon.png<br>desicc_risk_by_polygon.png<br>desicc_risk_median_by_polygon.png<br>desicc_risk_q1_by_polygon.png<br>desicc_risk_q3_by_polygon.png<br>desicc_window_count_by_polygon.png<br>desiccation_summary_statistics.csv |

### Clustering.py

The **clustering.py** contains the function regarding the clustering presented in Section 4.6 and Section 5.3  
**Error! Reference source not found.** The user can always add and/or modify this file.

| Description of the workflow (in chronological order)                                                | Name of the function                     | Name of the output                                                                  |
|-----------------------------------------------------------------------------------------------------|------------------------------------------|-------------------------------------------------------------------------------------|
| Performing the clustering:<br>Finding the best number of clusters, PCA analysis, cluster statistics | <i>perform_clustering_target_habitat</i> | metrics_with_clusters.csv<br>PCA_clusters.png<br>PCA_loadings.csv<br>silhouette.png |
| Join a GeoDataFrame (patch geometry) with a results DataFrame and save to a new shapefile           | <i>join_mesh_with_CSV_data</i>           | metrics_with_clusters.shp                                                           |

## 11. Literature references

- Bätz N, Judes C, Vanzo D, et al (2025) Patch-scale habitat dynamics: three metrics to assess ecological impacts of frequent hydropeaking. *Journal of Ecohydraulics* 10:79 – 106. <https://doi.org/10.1080/24705357.2024.2426790>
- Friese N, Mathers KL, Weber C, et al (2025) Habitat-specific response of macroinvertebrate drift to flow pulses: Implications for hydropeaking management. *Freshwater Biology* 70:e70103. 774 <https://doi.org/10.1111/fwb.70103>
- Jaulin E (2025) Looking at the reproduction of stream insects to improve our understanding of the ecological effects of artificial flow variations. Université Claude Bernard Lyon 1
- Kennedy TA, Muehlbauer JD, Yackulic CB, et al (2016) Flow management for hydropower extirpates aquatic insects, undermining river food webs. *BioScience* 66:561 – 575. 802 <https://doi.org/10.1093/biosci/biw059>
- Lecrivain A, De Cesare G, Weber C, & Bätz N (in Review). Macroinvertebrate habitat dynamics under frequent hydropower-induced discharge fluctuations: patch-scale metrics to quantify effects of hydropeaking and morphological complexity.
- Tonolla D, Schülting L, Dossi F, et al (2019) Assessment of hydropeaking impacts on the drift of aquatic macroinvertebrates – Flume and field experiments. Final Report
- Tonolla D (Ed.) (2023) Élaboration et évaluation de mesures d’assainissement des éclusées – Nouvelles connaissances issues de la recherche et de la pratique. Rapport sur mandat de l’OFEV.