

Finding Roots of Equations

Fixed-Point Iteration

Finding Roots of Equations

Fixed-Point Iteration

Looking for solutions of equations of the form

$$f(x) = x,$$

called fixed points of f .

Finding Roots of Equations

Fixed-Point Iteration

Looking for solutions of equations of the form

$$f(x) = x,$$

called fixed points of f .

Starting from a number x_0 , we generate a sequence x_1, x_2, \dots of numbers by the rule

$$x_{n+1} = f(x_n).$$

Finding Roots of Equations

Fixed-Point Iteration

Looking for solutions of equations of the form

$$f(x) = x,$$

called fixed points of f .

Starting from a number x_0 , we generate a sequence x_1, x_2, \dots of numbers by the rule

$$x_{n+1} = f(x_n).$$

In certain circumstances, $\lim x_n$ exists and converges to a fixed point.

Finding Roots of Equations

Fixed-Point Iteration

THEOREM A fixed-point theorem

Suppose that f is defined on an interval $I = [a, b]$ and satisfies the following two conditions:

- (i) $f(x)$ belongs to I whenever x belongs to I and
- (ii) there exists a constant K with $0 < K < 1$ such that for every u and v in I ,

$$|f(u) - f(v)| \leq K|u - v|.$$


Then f has a unique fixed point r in I , that is, $f(r) = r$, and starting with any number x_0 in I , the iterates

$$x_1 = f(x_0), \quad x_2 = f(x_1), \quad \dots$$

converge to r .

Finding Roots of Equations

Fixed-Point Iteration



Finding Roots of Equations

Fixed-Point Iteration

EXAMPLE

Find a root of the equation $\cos x = 5x$.

Finding Roots of Equations

Fixed-Point Iteration

EXAMPLE Find a root of the equation $\cos x = 5x$.

Solution This equation is of the form $f(x) = x$, where $f(x) = \frac{1}{5} \cos x$. Since $\cos x$ is close to 1 for x near 0, we see that $\frac{1}{5} \cos x$ will be close to $\frac{1}{5}$ when $x = \frac{1}{5}$. This suggests that a reasonable first guess at the fixed point is $x_0 = \frac{1}{5} = 0.2$.

Finding Roots of Equations

Fixed-Point Iteration

EXAMPLE Find a root of the equation $\cos x = 5x$.

Solution This equation is of the form $f(x) = x$, where $f(x) = \frac{1}{5} \cos x$. Since $\cos x$ is close to 1 for x near 0, we see that $\frac{1}{5} \cos x$ will be close to $\frac{1}{5}$ when $x = \frac{1}{5}$. This suggests that a reasonable first guess at the fixed point is $x_0 = \frac{1}{5} = 0.2$.


$$x_1 = \frac{1}{5} \cos x_0, \quad x_2 = \frac{1}{5} \cos x_1, \quad x_3 = \frac{1}{5} \cos x_2, \dots$$

The root is 0.196 164 28 to 8 decimal places.

n	x_n
0	0.2
1	0.196 013 32
2	0.196 170 16
3	0.196 164 05
4	0.196 164 29
5	0.196 164 28
6	0.196 164 28


Finding Roots of Equations

Newton's Method




Finding Roots of Equations

Newton's Method




Finding Roots of Equations

Newton's Method




Finding Roots of Equations

Newton's Method



Finding Roots of Equations

Newton's Method



$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Looking for a
fixed point of
the function
 $g(x) = x - \frac{f(x)}{f'(x)}$.

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$y = f(x_0) + f'(x_0)(x - x_0)$$

Finding Roots of Equations

Newton's Method

EXAMPLE

Use Newton's Method to find the only real root of the equation $x^3 - x - 1 = 0$ correct to 10 decimal places.

Finding Roots of Equations

Newton's Method

EXAMPLE

Use Newton's Method to find the only real root of the equation $x^3 - x - 1 = 0$ correct to 10 decimal places.

Solution We have $f(x) = x^3 - x - 1$ and $f'(x) = 3x^2 - 1$. Since f is continuous and since $f(1) = -1$ and $f(2) = 5$, the equation has a root in the interval $[1, 2]$.

$$x_{n+1} = x_n - \frac{x_n^3 - x_n - 1}{3x_n^2 - 1} = \frac{2x_n^3 + 1}{3x_n^2 - 1},$$

n	x_n	$f(x_n)$
0	1.5	0.875 000 000 000 ...
1	1.347 826 086 96 ...	0.100 682 173 091 ...
2	1.325 200 398 95 ...	0.002 058 361 917 ...
3	1.324 718 174 00 ...	0.000 000 924 378 ...
4	1.324 717 957 24 ...	0.000 000 000 000 ...
5	1.324 717 957 24 ...	

Indeterminate Forms

Type	Example
$[0/0]$	$\lim_{x \rightarrow 0} \frac{\sin x}{x}$
$[\infty/\infty]$	$\lim_{x \rightarrow 0} \frac{\ln(1/x^2)}{\cot(x^2)}$
$[0 \cdot \infty]$	$\lim_{x \rightarrow 0+} x \ln \frac{1}{x}$
$[\infty - \infty]$	$\lim_{x \rightarrow (\pi/2)-} \left(\tan x - \frac{1}{\pi - 2x} \right)$
$[0^0]$	$\lim_{x \rightarrow 0+} x^x$
$[\infty^0]$	$\lim_{x \rightarrow (\pi/2)-} (\tan x)^{\cos x}$
$[1^\infty]$	$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x} \right)^x$

Indeterminate Forms

THEOREM **The first l'Hôpital Rule**

Suppose the functions f and g are differentiable on the interval (a, b) , and $g'(x) \neq 0$ there. Suppose also that

- (i) $\lim_{x \rightarrow a+} f(x) = \lim_{x \rightarrow a+} g(x) = 0$ and
- (ii) $\lim_{x \rightarrow a+} \frac{f'(x)}{g'(x)} = L$ (where L is finite or ∞ or $-\infty$).

Then

$$\lim_{x \rightarrow a+} \frac{f(x)}{g(x)} = L.$$

Similar results hold if every occurrence of $\lim_{x \rightarrow a+}$ is replaced by $\lim_{x \rightarrow b-}$ or even $\lim_{x \rightarrow c}$ where $a < c < b$. The cases $a = -\infty$ and $b = \infty$ are also allowed.

Indeterminate Forms

EXAMPLE

Evaluate $\lim_{x \rightarrow 1} \frac{\ln x}{x^2 - 1}$.

Solution We have $\lim_{x \rightarrow 1} \frac{\ln x}{x^2 - 1} \left[\frac{0}{0} \right]$

$$= \lim_{x \rightarrow 1} \frac{1/x}{2x} = \lim_{x \rightarrow 1} \frac{1}{2x^2} = \frac{1}{2}.$$

Indeterminate Forms

EXAMPLE

Evaluate $\lim_{x \rightarrow 0} \frac{2 \sin x - \sin(2x)}{2e^x - 2 - 2x - x^2}$.

Solution We have (using l'Hôpital's Rule three times)

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{2 \sin x - \sin(2x)}{2e^x - 2 - 2x - x^2} & \quad \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ &= \lim_{x \rightarrow 0} \frac{2 \cos x - 2 \cos(2x)}{2e^x - 2 - 2x} \quad \text{cancel the 2s} \\ &= \lim_{x \rightarrow 0} \frac{\cos x - \cos(2x)}{e^x - 1 - x} \quad \text{still } \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ &= \lim_{x \rightarrow 0} \frac{-\sin x + 2 \sin(2x)}{e^x - 1} \quad \text{still } \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ &= \lim_{x \rightarrow 0} \frac{-\cos x + 4 \cos(2x)}{e^x} = \frac{-1 + 4}{1} = 3. \end{aligned}$$

Indeterminate Forms

EXAMPLE

Evaluate (a) $\lim_{x \rightarrow (\pi/2)^-} \frac{2x - \pi}{\cos^2 x}$ and (b) $\lim_{x \rightarrow 1^+} \frac{x}{\ln x}$.

Indeterminate Forms

EXAMPLE

Evaluate (a) $\lim_{x \rightarrow (\pi/2)^-} \frac{2x - \pi}{\cos^2 x}$ and (b) $\lim_{x \rightarrow 1^+} \frac{x}{\ln x}$.

Solution

$$\begin{aligned} \text{(a)} \quad & \lim_{x \rightarrow (\pi/2)^-} \frac{2x - \pi}{\cos^2 x} \quad \left[\begin{array}{c} 0 \\ 0 \end{array} \right] \\ &= \lim_{x \rightarrow (\pi/2)^-} \frac{2}{-2 \sin x \cos x} = -\infty \end{aligned}$$

Indeterminate Forms

EXAMPLE

Evaluate (a) $\lim_{x \rightarrow (\pi/2)^-} \frac{2x - \pi}{\cos^2 x}$ and (b) $\lim_{x \rightarrow 1^+} \frac{x}{\ln x}$.

Solution

$$\begin{aligned} \text{(a)} \quad & \lim_{x \rightarrow (\pi/2)^-} \frac{2x - \pi}{\cos^2 x} \quad \left[\begin{array}{c} 0 \\ 0 \end{array} \right] \\ &= \lim_{x \rightarrow (\pi/2)^-} \frac{2}{-2 \sin x \cos x} = -\infty \end{aligned}$$

(b) l'Hôpital's Rule cannot be used to evaluate $\lim_{x \rightarrow 1^+} x/(\ln x)$ because this is not an indeterminate form. The denominator approaches 0 as $x \rightarrow 1^+$, but the numerator does not approach 0. Since $\ln x > 0$ for $x > 1$, we have, directly,

$$\lim_{x \rightarrow 1^+} \frac{x}{\ln x} = \infty.$$

(Had we tried to apply l'Hôpital's Rule, we would have been led to the erroneous answer $\lim_{x \rightarrow 1^+} (1/(1/x)) = 1$.)

Indeterminate Forms

EXAMPLE

Evaluate $\lim_{x \rightarrow 0+} \left(\frac{1}{x} - \frac{1}{\sin x} \right)$.

Solution The indeterminate form here is of type $[\infty - \infty]$ to which l'Hôpital's Rule cannot be applied. However, it becomes $[0/0]$ after we combine the fractions into one fraction.

$$\begin{aligned} & \lim_{x \rightarrow 0+} \left(\frac{1}{x} - \frac{1}{\sin x} \right) \quad [\infty - \infty] \\ &= \lim_{x \rightarrow 0+} \frac{\sin x - x}{x \sin x} \quad \left[\frac{0}{0} \right] \\ &= \lim_{x \rightarrow 0+} \frac{\cos x - 1}{\sin x + x \cos x} \quad \left[\frac{0}{0} \right] \\ &= \lim_{x \rightarrow 0+} \frac{-\sin x}{2 \cos x - x \sin x} = \frac{-0}{2} = 0. \end{aligned}$$

Indeterminate Forms

THEOREM **The second l'Hôpital Rule**

Suppose that f and g are differentiable on the interval (a, b) and that $g'(x) \neq 0$ there. Suppose also that

(i) $\lim_{x \rightarrow a+} g(x) = \pm\infty$ and

(ii) $\lim_{x \rightarrow a+} \frac{f'(x)}{g'(x)} = L$ (where L is finite, or ∞ or $-\infty$).

Then

$$\lim_{x \rightarrow a+} \frac{f(x)}{g(x)} = L.$$

Again, similar results hold for $\lim_{x \rightarrow b-}$ and for $\lim_{x \rightarrow c}$, and the cases $a = -\infty$ and $b = \infty$ are allowed.

Indeterminate Forms

EXAMPLE

Evaluate (a) $\lim_{x \rightarrow \infty} \frac{x^2}{e^x}$ and (b) $\lim_{x \rightarrow 0^+} x^a \ln x$, where $a > 0$.

Indeterminate Forms

EXAMPLE

Evaluate (a) $\lim_{x \rightarrow \infty} \frac{x^2}{e^x}$ and (b) $\lim_{x \rightarrow 0+} x^a \ln x$, where $a > 0$.

Solution

$$\begin{aligned} \text{(a)} \quad & \lim_{x \rightarrow \infty} \frac{x^2}{e^x} \quad \left[\frac{\infty}{\infty} \right] \\ &= \lim_{x \rightarrow \infty} \frac{2x}{e^x} \quad \text{still } \left[\frac{\infty}{\infty} \right] \\ &= \lim_{x \rightarrow \infty} \frac{2}{e^x} = 0. \end{aligned}$$

Similarly, one can show that $\lim_{x \rightarrow \infty} x^n / e^x = 0$ for any positive integer n by repeated applications of l'Hôpital's Rule.

Indeterminate Forms

EXAMPLE

Evaluate (a) $\lim_{x \rightarrow \infty} \frac{x^2}{e^x}$ and (b) $\lim_{x \rightarrow 0+} x^a \ln x$, where $a > 0$.

Solution

$$\begin{aligned} \text{(a)} \quad & \lim_{x \rightarrow \infty} \frac{x^2}{e^x} \quad \left[\frac{\infty}{\infty} \right] \\ &= \lim_{x \rightarrow \infty} \frac{2x}{e^x} \quad \text{still } \left[\frac{\infty}{\infty} \right] \\ &= \lim_{x \rightarrow \infty} \frac{2}{e^x} = 0. \end{aligned}$$

Similarly, one can show that $\lim_{x \rightarrow \infty} x^n / e^x = 0$ for any positive integer n by repeated applications of l'Hôpital's Rule.

$$\begin{aligned} \text{(b)} \quad & \lim_{x \rightarrow 0+} x^a \ln x \quad (a > 0) \quad [0 \cdot (-\infty)] \\ &= \lim_{x \rightarrow 0+} \frac{\ln x}{x^{-a}} \quad \left[\frac{-\infty}{\infty} \right] \\ &= \lim_{x \rightarrow 0+} \frac{1/x}{-ax^{-a-1}} = \lim_{x \rightarrow 0+} \frac{x^a}{-a} = 0. \end{aligned}$$

Indeterminate Forms

EXAMPLE Evaluate $\lim_{x \rightarrow 0+} x^x$.

Solution This indeterminate form is of type $[0^0]$. Let $y = x^x$. Then

$$\lim_{x \rightarrow 0+} \ln y = \lim_{x \rightarrow 0+} x \ln x = 0,$$

$$\text{Hence } \lim_{x \rightarrow 0} x^x = \lim_{x \rightarrow 0+} y = e^0 = 1.$$

$$\lim y = e^{\ln(\lim y)} = e^{\lim(\ln y)}$$

Indeterminate Forms

EXAMPLE

Evaluate $\lim_{x \rightarrow \infty} \left(1 + \sin \frac{3}{x}\right)^x$.

Solution This indeterminate form is of type 1^∞ . Let $y = \left(1 + \sin \frac{3}{x}\right)^x$. Then, taking \ln of both sides,

$$\begin{aligned}\lim_{x \rightarrow \infty} \ln y &= \lim_{x \rightarrow \infty} x \ln \left(1 + \sin \frac{3}{x}\right) \quad [\infty \cdot 0] \\&= \lim_{x \rightarrow \infty} \frac{\ln \left(1 + \sin \frac{3}{x}\right)}{\frac{1}{x}} \quad \left[\frac{0}{0}\right] \\&= \lim_{x \rightarrow \infty} \frac{\frac{1}{1 + \sin \frac{3}{x}} \left(\cos \frac{3}{x}\right) \left(-\frac{3}{x^2}\right)}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{3 \cos \frac{3}{x}}{1 + \sin \frac{3}{x}} = 3.\end{aligned}$$

Hence $\lim_{x \rightarrow \infty} \left(1 + \sin \frac{3}{x}\right)^x = e^3$.

Extreme Values

Absolute extreme values

Function f has an **absolute maximum value** $f(x_0)$ at the point x_0 in its domain if $f(x) \leq f(x_0)$ holds for every x in the domain of f .

Similarly, f has an **absolute minimum value** $f(x_1)$ at the point x_1 in its domain if $f(x) \geq f(x_1)$ holds for every x in the domain of f .

Extreme Values

Absolute extreme values

Function f has an **absolute maximum value** $f(x_0)$ at the point x_0 in its domain if $f(x) \leq f(x_0)$ holds for every x in the domain of f .


Similarly, f has an **absolute minimum value** $f(x_1)$ at the point x_1 in its domain if $f(x) \geq f(x_1)$ holds for every x in the domain of f .

THEOREM

Existence of extreme values

If the domain of the function f is a *closed, finite interval* or a union of finitely many such intervals, and if f is *continuous* on that domain, then f must have an absolute maximum value and an absolute minimum value.

Extreme Values



Extreme Values

Local extreme values

Function f has a **local maximum value (loc max)** $f(x_0)$ at the point x_0 in its domain provided there exists a number $h > 0$ such that $f(x) \leq f(x_0)$ whenever x is in the domain of f and $|x - x_0| < h$.

Similarly, f has a **local minimum value (loc min)** $f(x_1)$ at the point x_1 in its domain provided there exists a number $h > 0$ such that $f(x) \geq f(x_1)$ whenever x is in the domain of f and $|x - x_1| < h$.

Extreme Values

Critical Points, Singular Points, and Endpoints

Figure 4.17 suggests that a function $f(x)$ can have local extreme values only at points x of three special types:

- (i) **critical points** of f (points x in $\mathcal{D}(f)$ where $f'(x) = 0$),
- (ii) **singular points** of f (points x in $\mathcal{D}(f)$ where $f'(x)$ is not defined), and
- (iii) **endpoints** of the domain of f (points in $\mathcal{D}(f)$ that do not belong to any open interval contained in $\mathcal{D}(f)$).

Extreme Values

Critical Points, Singular Points, and Endpoints

Figure 4.17 suggests that a function $f(x)$ can have local extreme values only at points x of three special types:


- (i) **critical points** of f (points x in $\mathcal{D}(f)$ where $f'(x) = 0$),
- (ii) **singular points** of f (points x in $\mathcal{D}(f)$ where $f'(x)$ is not defined), and
- (iii) **endpoints** of the domain of f (points in $\mathcal{D}(f)$ that do not belong to any open interval contained in $\mathcal{D}(f)$).

THEOREM

Locating extreme values

If the function f is defined on an interval I and has a local maximum (or local minimum) value at point $x = x_0$ in I , then x_0 must be either a critical point of f , a singular point of f , or an endpoint of I .

Extreme Values



A function need not have extreme values
at a critical point or a singular point

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Solution Since g is a polynomial, it can have no singular points. For critical points, we calculate

$$\begin{aligned} g'(x) &= 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) \\ &= 3(x + 1)(x - 3) \\ &= 0 \quad \text{if } x = -1 \text{ or } x = 3. \end{aligned}$$

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Solution Since g is a polynomial, it can have no singular points. For critical points, we calculate

$$\begin{aligned} g'(x) &= 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) \\ &= 3(x + 1)(x - 3) \\ &= 0 \quad \text{if } x = -1 \text{ or } x = 3. \end{aligned}$$

not in the domain!

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Solution Since g is a polynomial, it can have no singular points. For critical points, we calculate

$$\begin{aligned} g'(x) &= 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) \\ &= 3(x + 1)(x - 3) \\ &= 0 \quad \text{if } x = -1 \text{ or } x = 3. \end{aligned}$$

not in the domain!

$$g(-2) = 0, \quad g(-1) = 7, \quad g(2) = -20.$$

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Solution Since g is a polynomial, it can have no singular points. For critical points, we calculate

$$\begin{aligned} g'(x) &= 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) \\ &= 3(x + 1)(x - 3) \\ &= 0 \quad \text{if } x = -1 \text{ or } x = 3. \end{aligned}$$

not in the domain!

$$g(-2) = 0, \quad g(-1) = 7, \quad g(2) = -20.$$

maximum
value

minimum
value

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of the function $g(x) = x^3 - 3x^2 - 9x + 2$ on the interval $-2 \leq x \leq 2$.

Solution Since g is a polynomial, it can have no singular points. For critical points, we calculate


$$\begin{aligned} g'(x) &= 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) \\ &= 3(x + 1)(x - 3) \\ &= 0 \quad \text{if } x = -1 \text{ or } x = 3. \end{aligned}$$

not in the domain!

$$g(-2) = 0, \quad g(-1) = 7, \quad g(2) = -20.$$

maximum
value

minimum
value



Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of $h(x) = 3x^{2/3} - 2x$ on the interval $[-1, 1]$.

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of $h(x) = 3x^{2/3} - 2x$ on the interval $[-1, 1]$.

Solution The derivative of h is

$$h'(x) = 3 \left(\frac{2}{3} \right) x^{-1/3} - 2 = 2(x^{-1/3} - 1).$$

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of $h(x) = 3x^{2/3} - 2x$ on the interval $[-1, 1]$.

Solution The derivative of h is

$$h'(x) = 3 \left(\frac{2}{3} \right) x^{-1/3} - 2 = 2(x^{-1/3} - 1).$$

$x = 0$ is a singular point of h . Also, h has a critical point at $x = 1$

$$h(-1) = 5, \quad h(0) = 0, \quad h(1) = 1.$$

Extreme Values

Finding Absolute Extreme Values

EXAMPLE

Find the maximum and minimum values of $h(x) = 3x^{2/3} - 2x$ on the interval $[-1, 1]$.

Solution The derivative of h is

$$h'(x) = 3 \left(\frac{2}{3} \right) x^{-1/3} - 2 = 2(x^{-1/3} - 1).$$

$x = 0$ is a singular point of h . Also, h has a critical point at $x = 1$

$$\boxed{h(-1) = 5}, \quad \boxed{h(0) = 0}, \quad h(1) = 1.$$

max.

min.

Extreme Values

THEOREM

The First Derivative Test

PART I. Testing interior critical points and singular points.

Suppose that f is continuous at x_0 , and x_0 is not an endpoint of the domain of f .

- (a) If there exists an open interval (a, b) containing x_0 such that $f'(x) > 0$ on (a, x_0) and $f'(x) < 0$ on (x_0, b) , then f has a local maximum value at x_0 .
- (b) If there exists an open interval (a, b) containing x_0 such that $f'(x) < 0$ on (a, x_0) and $f'(x) > 0$ on (x_0, b) , then f has a local minimum value at x_0 .

PART II. Testing endpoints of the domain.

Suppose a is a left endpoint of the domain of f and f is right continuous at a .

- (c) If $f'(x) > 0$ on some interval (a, b) , then f has a local minimum value at a .
- (d) If $f'(x) < 0$ on some interval (a, b) , then f has a local maximum value at a .

Suppose b is a right endpoint of the domain of f and f is left continuous at b .

- (e) If $f'(x) > 0$ on some interval (a, b) , then f has a local maximum value at b .
- (f) If $f'(x) < 0$ on some interval (a, b) , then f has a local minimum value at b .

Extreme Values

EXAMPLE

Find the local and absolute extreme values of $f(x) = x^4 - 2x^2 - 3$ on the interval $[-2, 2]$. Sketch the graph of f .