

# Sketching the Graph of a Function

---

## Asymptotes

The graph of  $y = f(x)$  has a **vertical asymptote** at  $x = a$  if

either  $\lim_{x \rightarrow a-} f(x) = \pm\infty$     or     $\lim_{x \rightarrow a+} f(x) = \pm\infty$ ,    or both.

# Sketching the Graph of a Function

---

## Asymptotes

### EXAMPLE

Find the vertical asymptotes of  $f(x) = \frac{1}{x^2 - x}$ . How does the graph approach these asymptotes?

# Sketching the Graph of a Function

---

## Asymptotes

### EXAMPLE

Find the vertical asymptotes of  $f(x) = \frac{1}{x^2 - x}$ . How does the graph approach these asymptotes?

**Solution** The denominator  $x^2 - x = x(x - 1)$  approaches 0 as  $x$  approaches 0 or 1, so  $f$  has vertical asymptotes at  $x = 0$  and  $x = 1$ .

$$\lim_{x \rightarrow 0^-} \frac{1}{x^2 - x} = \infty,$$

$$\lim_{x \rightarrow 1^-} \frac{1}{x^2 - x} = -\infty,$$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^2 - x} = -\infty,$$

$$\lim_{x \rightarrow 1^+} \frac{1}{x^2 - x} = \infty.$$

# Sketching the Graph of a Function

## Asymptotes

### EXAMPLE

Find the vertical asymptotes of  $f(x) = \frac{1}{x^2 - x}$ . How does the graph approach these asymptotes?


**Solution** The denominator  $x^2 - x = x(x - 1)$  approaches 0 as  $x$  approaches 0 or 1, so  $f$  has vertical asymptotes at  $x = 0$  and  $x = 1$ .

$$\lim_{x \rightarrow 0^-} \frac{1}{x^2 - x} = \infty,$$

$$\lim_{x \rightarrow 1^-} \frac{1}{x^2 - x} = -\infty,$$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^2 - x} = -\infty,$$

$$\lim_{x \rightarrow 1^+} \frac{1}{x^2 - x} = \infty.$$



## Sketching the Graph of a Function

### **Asymptotes**

The graph  $y=f(x)$  has a horizontal asymptote at  $y=b$  if either

$$\lim_{x \rightarrow \infty} f(x) = b \text{ or } \lim_{x \rightarrow -\infty} f(x) = b.$$

# Sketching the Graph of a Function

---

## Asymptotes

---

### EXAMPLE

---

Find the horizontal asymptotes of

(a)  $f(x) = \frac{1}{x^2 - x}$     and    (b)  $g(x) = \frac{x^4 + x^2}{x^4 + 1}$ .

# Sketching the Graph of a Function

---

## Asymptotes

### EXAMPLE

Find the horizontal asymptotes of

$$(a) f(x) = \frac{1}{x^2 - x} \quad \text{and} \quad (b) g(x) = \frac{x^4 + x^2}{x^4 + 1}.$$

### *Solution*

(a) The function  $f$  has horizontal asymptote  $y = 0$  since

$$\lim_{x \rightarrow \pm\infty} \frac{1}{x^2 - x} = \lim_{x \rightarrow \pm\infty} \frac{1/x^2}{1 - (1/x)} = \frac{0}{1} = 0.$$

# Sketching the Graph of a Function

---

## Asymptotes

### EXAMPLE

Find the horizontal asymptotes of

$$(a) f(x) = \frac{1}{x^2 - x} \quad \text{and} \quad (b) g(x) = \frac{x^4 + x^2}{x^4 + 1}.$$

### *Solution*

(a) The function  $f$  has horizontal asymptote  $y = 0$  since

$$\lim_{x \rightarrow \pm\infty} \frac{1}{x^2 - x} = \lim_{x \rightarrow \pm\infty} \frac{1/x^2}{1 - (1/x)} = \frac{0}{1} = 0.$$

(b) The function  $g$  has horizontal asymptote  $y = 1$  since

$$\lim_{x \rightarrow \pm\infty} \frac{x^4 + x^2}{x^4 + 1} = \lim_{x \rightarrow \pm\infty} \frac{1 + (1/x^2)}{1 + (1/x^4)} = \frac{1}{1} = 1.$$



# Sketching the Graph of a Function

## Asymptotes

### EXAMPLE

Find the horizontal asymptotes of

$$(a) f(x) = \frac{1}{x^2 - x} \quad \text{and} \quad (b) g(x) = \frac{x^4 + x^2}{x^4 + 1}.$$

### Solution

(a) The function  $f$  has horizontal asymptote  $y = 0$  since

$$\lim_{x \rightarrow \pm\infty} \frac{1}{x^2 - x} = \lim_{x \rightarrow \pm\infty} \frac{1/x^2}{1 - (1/x)} = \frac{0}{1} = 0.$$

(b) The function  $g$  has horizontal asymptote  $y = 1$  since

$$\lim_{x \rightarrow \pm\infty} \frac{x^4 + x^2}{x^4 + 1} = \lim_{x \rightarrow \pm\infty} \frac{1 + (1/x^2)}{1 + (1/x^4)} = \frac{1}{1} = 1.$$

Observe that the graph of  $g$  crosses its asymptote twice.

# Sketching the Graph of a Function

---

## Asymptotes

The straight line  $y = ax + b$  (where  $a \neq 0$ ) is an **oblique asymptote** of the graph of  $y = f(x)$  if

$$\text{either } \lim_{x \rightarrow -\infty} (f(x) - (ax + b)) = 0 \quad \text{or} \quad \lim_{x \rightarrow \infty} (f(x) - (ax + b)) = 0,$$

or both.


# Sketching the Graph of a Function

## Asymptotes

### EXAMPLE

Consider the function  $f(x) = \frac{x^2 + 1}{x} = x + \frac{1}{x}$ , whose graph is shown. The straight line  $y = x$  is a *two-sided* oblique asymptote of the graph of  $f$  because

$$\lim_{x \rightarrow \pm\infty} (f(x) - x) = \lim_{x \rightarrow \pm\infty} \frac{1}{x} = 0.$$




# Sketching the Graph of a Function

## Asymptotes

### EXAMPLE

The graph of  $y = \frac{x e^x}{1 + e^x}$  has a horizontal asymptote  $y = 0$  at the left and an oblique asymptote  $y = x$  at the right:

$$\lim_{x \rightarrow -\infty} \frac{x e^x}{1 + e^x} = \frac{0}{1} = 0 \quad \text{and}$$
$$\lim_{x \rightarrow \infty} \left( \frac{x e^x}{1 + e^x} - x \right) = \lim_{x \rightarrow \infty} \frac{x(e^x - 1 - e^x)}{1 + e^x} = \lim_{x \rightarrow \infty} \frac{-x}{1 + e^x} = 0.$$



# Sketching the Graph of a Function

---

## Asymptotes

### Asymptotes of a rational function

Suppose that  $f(x) = \frac{P_m(x)}{Q_n(x)}$ , where  $P_m$  and  $Q_n$  are polynomials of degree  $m$  and  $n$ , respectively. Suppose also that  $P_m$  and  $Q_n$  have no common linear factors. Then

- (a) The graph of  $f$  has a vertical asymptote at every position  $x$  such that  $Q_n(x) = 0$ .
- (b) The graph of  $f$  has a two-sided horizontal asymptote  $y = 0$  if  $m < n$ .
- (c) The graph of  $f$  has a two-sided horizontal asymptote  $y = L$ , ( $L \neq 0$ ) if  $m = n$ .  $L$  is the quotient of the coefficients of the highest degree terms in  $P_m$  and  $Q_n$ .
- (d) The graph of  $f$  has a two-sided oblique asymptote if  $m = n + 1$ . This asymptote can be found by dividing  $Q_n$  into  $P_m$  to obtain a linear quotient,  $ax + b$ , and remainder,  $R$ , a polynomial of degree at most  $n - 1$ . That is,

$$f(x) = ax + b + \frac{R(x)}{Q_n(x)}.$$

The oblique asymptote is  $y = ax + b$ .

- (e) The graph of  $f$  has no horizontal or oblique asymptotes if  $m > n + 1$ .

# Sketching the Graph of a Function

---

## Asymptotes

### EXAMPLE

Find the oblique asymptote of  $y = \frac{x^3}{x^2 + x + 1}$ .

# Sketching the Graph of a Function

---

## Asymptotes

### EXAMPLE

Find the oblique asymptote of  $y = \frac{x^3}{x^2 + x + 1}$ .

**Solution** We can either obtain the quotient by long division:

$$\begin{array}{r|l} x^3 & x^2 + x + 1 \\ \hline x^3 + x^2 + x & x - 1 \\ \hline -x^2 - x & \\ -x^2 - x - 1 & \\ \hline 1 & \end{array}$$

$$\frac{x^3}{x^2 + x + 1} = x - 1 + \frac{1}{x^2 + x + 1}$$

# Sketching the Graph of a Function

## Asymptotes

### EXAMPLE

Find the oblique asymptote of  $y = \frac{x^3}{x^2 + x + 1}$ .

**Solution** We can either obtain the quotient by long division:

$$\begin{array}{r|l} x^3 & x^2 + x + 1 \\ \hline x^3 + x^2 + x & x - 1 \\ \hline -x^2 - x & \\ -x^2 - x - 1 & \\ \hline 1 & \end{array}$$

$$\frac{x^3}{x^2 + x + 1} = \boxed{x - 1} + \frac{1}{x^2 + x + 1}$$

$y = x - 1$  is the oblique asymptote.



# Sketching the Graph of a Function

---

## **Checklist for curve sketching**

# Sketching the Graph of a Function

## **Checklist for curve sketching**

1. Calculate  $f'(x)$  and  $f''(x)$ , and express the results in factored form.

# Sketching the Graph of a Function

## **Checklist for curve sketching**

1. Calculate  $f'(x)$  and  $f''(x)$ , and express the results in factored form.
2. Examine  $f(x)$  to determine its domain and the following items:
  - (a) Any vertical asymptotes. (Look for zeros of denominators.)
  - (b) Any horizontal or oblique asymptotes. (Consider  $\lim_{x \rightarrow \pm\infty} f(x)$ .)
  - (c) Any obvious symmetry. (Is  $f$  even or odd?)
  - (d) Any easily calculated intercepts (points with coordinates  $(x, 0)$  or  $(0, y)$ ) or endpoints or other “obvious” points.

# Sketching the Graph of a Function

## **Checklist for curve sketching**

3. Examine  $f'(x)$  for the following:
  - (a) Any critical points.
  - (b) Any points where  $f'$  is not defined. (These will include singular points, endpoints of the domain of  $f$ , and vertical asymptotes.)
  - (c) Intervals on which  $f'$  is positive or negative. It's a good idea to convey this information in the form of a chart such as those used in the examples. Conclusions about where  $f$  is increasing and decreasing and classification of some critical and singular points as local maxima and minima can also be indicated on the chart.

# Sketching the Graph of a Function

## **Checklist for curve sketching**

4. Examine  $f''(x)$  for the following:
  - (a) Points where  $f''(x) = 0$ .
  - (b) Points where  $f''(x)$  is undefined. (These will include singular points, endpoints, vertical asymptotes, and possibly other points as well, where  $f'$  is defined but  $f''$  isn't.)
  - (c) Intervals where  $f''$  is positive or negative and where  $f$  is therefore concave up or down. Use a chart.
  - (d) Any inflection points.

## Sketching the Graph of a Function

**EXAMPLE**

Sketch the graph of  $y = \frac{x^2 + 2x + 4}{2x}$ .

## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $y = \frac{x^2 + 2x + 4}{2x}$ .

### ***Solution***

Domain: all  $x$  except 0. Vertical asymptote:  $x = 0$ ,

## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $y = \frac{x^2 + 2x + 4}{2x}$ .

### ***Solution***

Domain: all  $x$  except 0. Vertical asymptote:  $x = 0$ ,

$$y = \frac{x}{2} + 1 + \frac{2}{x},$$

$y = (x/2) + 1$  is an oblique asymptote.



## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $y = \frac{x^2 + 2x + 4}{2x}$ .

### ***Solution***

Domain: all  $x$  except 0. Vertical asymptote:  $x = 0$ ,

$$y = \frac{x}{2} + 1 + \frac{2}{x},$$

$y = (x/2) + 1$  is an oblique asymptote.

$$y' = \frac{1}{2} - \frac{2}{x^2} = \frac{x^2 - 4}{2x^2},$$

$$y'' = \frac{4}{x^3}.$$

# Sketching the Graph of a Function

## EXAMPLE

Sketch the graph of  $y = \frac{x^2 + 2x + 4}{2x}$ .

## Solution

Domain: all  $x$  except 0. Vertical asymptote:  $x = 0$ ,

$$y = \frac{x}{2} + 1 + \frac{2}{x},$$

$y = (x/2) + 1$  is an oblique asymptote.

$$y' = \frac{1}{2} - \frac{2}{x^2} = \frac{x^2 - 4}{2x^2},$$

$$y'' = \frac{4}{x^3}.$$

$\overline{x^3}$	CP			ASY		CP	
$x$	-2			0		2	
$y'$	+	0	-	undef	-	0	+
$y''$	-		-	undef	+		+
$y$	↗	max	↘	undef	↘	min	↗
	(		(		)		)

# Sketching the Graph of a Function

## EXAMPLE

Sketch the graph of  $y = \frac{x^2 + 2x + 4}{2x}$ .

## Solution

Domain: all  $x$  except 0. Vertical asymptote:  $x = 0$ ,


$$y = \frac{x}{2} + 1 + \frac{2}{x},$$

$y = (x/2) + 1$  is an oblique asymptote.

$$y' = \frac{1}{2} - \frac{2}{x^2} = \frac{x^2 - 4}{2x^2},$$

$$y'' = \frac{4}{x^3}.$$

$\overline{x^3}$	CP		ASY		CP		
$x$	-2		0		2		
$y'$	+	0	-	undef	-	0	+
$y''$	-		-	undef	+		+
$y$	↗	max	↘	undef	↘	min	↗
	(		(		)		)



## Sketching the Graph of a Function

### EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

## Sketching the Graph of a Function

### EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

### ***Solution***

Domain: all  $x$  except  $\pm 2$ .

## Sketching the Graph of a Function

### EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

### ***Solution***

Domain: all  $x$  except  $\pm 2$ .    Vertical asymptotes:  $x = -2$  and  $x = 2$ .

## Sketching the Graph of a Function

### EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

### ***Solution***

Domain: all  $x$  except  $\pm 2$ . Vertical asymptotes:  $x = -2$  and  $x = 2$ .

Horizontal asymptote:  $y = 1$  (as  $x \rightarrow \pm\infty$ ).

## Sketching the Graph of a Function

### EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

### ***Solution***

Domain: all  $x$  except  $\pm 2$ . Vertical asymptotes:  $x = -2$  and  $x = 2$ .

Horizontal asymptote:  $y = 1$  (as  $x \rightarrow \pm\infty$ ).

Symmetry: about the  $y$ -axis ( $y$  is even).



## Sketching the Graph of a Function

### EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

### ***Solution***

Domain: all  $x$  except  $\pm 2$ . Vertical asymptotes:  $x = -2$  and  $x = 2$ .

Horizontal asymptote:  $y = 1$  (as  $x \rightarrow \pm\infty$ ).

Symmetry: about the  $y$ -axis ( $y$  is even).

Intercepts:  $(0, 1/4)$ ,  $(-1, 0)$ , and  $(1, 0)$ .

# Sketching the Graph of a Function

---

## **EXAMPLE**

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

### ***Solution***

Domain: all  $x$  except  $\pm 2$ . Vertical asymptotes:  $x = -2$  and  $x = 2$ .

Horizontal asymptote:  $y = 1$  (as  $x \rightarrow \pm\infty$ ).

Symmetry: about the  $y$ -axis ( $y$  is even).

Intercepts:  $(0, 1/4)$ ,  $(-1, 0)$ , and  $(1, 0)$ .

$$f'(x) = \frac{-6x}{(x^2 - 4)^2}, \quad f''(x) = \frac{6(3x^2 + 4)}{(x^2 - 4)^3}.$$

# Sketching the Graph of a Function

## EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

## Solution

Domain: all  $x$  except  $\pm 2$ . Vertical asymptotes:  $x = -2$  and  $x = 2$ .

Horizontal asymptote:  $y = 1$  (as  $x \rightarrow \pm\infty$ ).

Symmetry: about the  $y$ -axis ( $y$  is even).

Intercepts:  $(0, 1/4)$ ,  $(-1, 0)$ , and  $(1, 0)$ .

$$f'(x) = \frac{-6x}{(x^2 - 4)^2}, \quad f''(x) = \frac{6(3x^2 + 4)}{(x^2 - 4)^3}.$$

	ASY			CP	ASY		
$x$	$-2$			$0$	$2$		
$f'$	+	undef	+	$0$	-	undef	-
$f''$	+	undef	-		-	undef	+
$f$	$\nearrow$	undef	$\nearrow$	max	$\searrow$	undef	$\searrow$
	$\cup$		$\cap$		$\cap$		$\cup$

# Sketching the Graph of a Function

## EXAMPLE

Sketch the graph of  $f(x) = \frac{x^2 - 1}{x^2 - 4}$ .

## Solution

Domain: all  $x$  except  $\pm 2$ . Vertical asymptotes:  $x = -2$  and  $x = 2$ .


Horizontal asymptote:  $y = 1$  (as  $x \rightarrow \pm\infty$ ).

Symmetry: about the  $y$ -axis ( $y$  is even).

Intercepts:  $(0, 1/4)$ ,  $(-1, 0)$ , and  $(1, 0)$ .

$$f'(x) = \frac{-6x}{(x^2 - 4)^2}, \quad f''(x) = \frac{6(3x^2 + 4)}{(x^2 - 4)^3}.$$

	ASY		CP		ASY	
$x$	$-2$		$0$		$2$	
$f'$	+	undef	+	$0$	-	undef
$f''$	+	undef	-		-	undef
$f$	$\nearrow$	undef	$\nearrow$	max	$\searrow$	undef
	$\cup$		$\cap$		$\cap$	$\cup$



## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

## Sketching the Graph of a Function

---

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .

# Sketching the Graph of a Function

---

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .

Domain: all  $x$ .

# Sketching the Graph of a Function

---

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .

Domain: all  $x$ .

Horizontal asymptote:  $y = 0$ . Note that if  $t = x^2/2$ , then

$|xe^{-x^2/2}| = \sqrt{2t}e^{-t} \rightarrow 0$  as  $t \rightarrow \infty$  (hence as  $x \rightarrow \pm\infty$ ).



# Sketching the Graph of a Function

---

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .

Domain: all  $x$ .

Horizontal asymptote:  $y = 0$ . Note that if  $t = x^2/2$ , then

$|xe^{-x^2/2}| = \sqrt{2t}e^{-t} \rightarrow 0$  as  $t \rightarrow \infty$  (hence as  $x \rightarrow \pm\infty$ ).

Symmetry: about the origin ( $y$  is odd). Intercepts:  $(0, 0)$ .

# Sketching the Graph of a Function

---

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .

Domain: all  $x$ .

Horizontal asymptote:  $y = 0$ . Note that if  $t = x^2/2$ , then

$|xe^{-x^2/2}| = \sqrt{2t}e^{-t} \rightarrow 0$  as  $t \rightarrow \infty$  (hence as  $x \rightarrow \pm\infty$ ).

Symmetry: about the origin ( $y$  is odd). Intercepts:  $(0, 0)$ .

Critical points:  $x = \pm 1$ ; points  $(\pm 1, \pm 1/\sqrt{e}) \approx (\pm 1, \pm 0.61)$ .

$y'' = 0$  at  $x = 0$  and  $x = \pm\sqrt{3}$ ;

points  $(0, 0)$ ,  $(\pm\sqrt{3}, \pm\sqrt{3}e^{-3/2}) \approx (\pm 1.73, \pm 0.39)$ .

# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .

Domain: all  $x$ .

Horizontal asymptote:  $y = 0$ . Note that if  $t = x^2/2$ , then  $|xe^{-x^2/2}| = \sqrt{2t}e^{-t} \rightarrow 0$  as  $t \rightarrow \infty$  (hence as  $x \rightarrow \pm\infty$ ).

Symmetry: about the origin ( $y$  is odd). Intercepts:  $(0, 0)$ .

Critical points:  $x = \pm 1$ ; points  $(\pm 1, \pm 1/\sqrt{e}) \approx (\pm 1, \pm 0.61)$ .

$y'' = 0$  at  $x = 0$  and  $x = \pm\sqrt{3}$ ;

points  $(0, 0), (\pm\sqrt{3}, \pm\sqrt{3}e^{-3/2}) \approx (\pm 1.73, \pm 0.39)$ .

$x$	$-\sqrt{3}$	CP	$-1$	$0$	CP	$1$	$\sqrt{3}$	
$y'$	$-$	$-$	$0$	$+$	$+$	$0$	$-$	$-$
$y''$	$-$	$0$	$+$	$+$	$0$	$-$	$-$	$0$
$y$	$\searrow$	$\searrow$	min	$\nearrow$	$\nearrow$	max	$\searrow$	$\searrow$
	$\cap$	infl	$\cup$	$\cup$	infl	$\cap$	$\cap$	infl

# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $y = xe^{-x^2/2}$ .

**Solution** We have  $y' = (1 - x^2)e^{-x^2/2}$ ,  $y'' = x(x^2 - 3)e^{-x^2/2}$ .


Domain: all  $x$ .

Horizontal asymptote:  $y = 0$ . Note that if  $t = x^2/2$ , then  $|xe^{-x^2/2}| = \sqrt{2t}e^{-t} \rightarrow 0$  as  $t \rightarrow \infty$  (hence as  $x \rightarrow \pm\infty$ ).

Symmetry: about the origin ( $y$  is odd). Intercepts:  $(0, 0)$ .

Critical points:  $x = \pm 1$ ; points  $(\pm 1, \pm 1/\sqrt{e}) \approx (\pm 1, \pm 0.61)$ .

$y'' = 0$  at  $x = 0$  and  $x = \pm\sqrt{3}$ ;  
points  $(0, 0), (\pm\sqrt{3}, \pm\sqrt{3}e^{-3/2}) \approx (\pm 1.73, \pm 0.39)$ .



			CP				CP			
$x$		$-\sqrt{3}$		$-1$		$0$		$1$		$\sqrt{3}$
$y'$	$-$	$-$	$0$	$+$		$+$	$0$	$-$		$-$
$y''$	$-$	$0$	$+$	$+$	$0$	$-$		$-$	$0$	$+$
$y$	$\searrow$	$\searrow$	min	$\nearrow$		$\nearrow$	max	$\searrow$		$\searrow$
	$\curvearrowright$	infl	$\curvearrowright$	$\curvearrowright$	infl	$\curvearrowright$		$\curvearrowright$	infl	$\curvearrowright$

## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

Domain: all  $x$ .

## Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

Domain: all  $x$ .

Asymptotes: none. ( $f(x)$  grows like  $x^{4/3}$  as  $x \rightarrow \pm\infty$ .)



# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

Domain: all  $x$ .

Asymptotes: none. ( $f(x)$  grows like  $x^{4/3}$  as  $x \rightarrow \pm\infty$ .)

Symmetry: about the  $y$ -axis ( $f$  is an even function).

# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

Domain: all  $x$ .

Asymptotes: none. ( $f(x)$  grows like  $x^{4/3}$  as  $x \rightarrow \pm\infty$ .)

Symmetry: about the  $y$ -axis ( $f$  is an even function).

Intercepts:  $(\pm 1, 0), (0, 1)$ .

# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

Domain: all  $x$ .

Asymptotes: none. ( $f(x)$  grows like  $x^{4/3}$  as  $x \rightarrow \pm\infty$ .)

Symmetry: about the  $y$ -axis ( $f$  is an even function).

Intercepts:  $(\pm 1, 0)$ ,  $(0, 1)$ .

Critical points:  $x = 0$ ; singular points:  $x = \pm 1$ .

$f''(x) = 0$  at  $x = \pm\sqrt{3}$ ; points  $(\pm\sqrt{3}, 2^{2/3}) \approx (\pm 1.73, 1.59)$ ;

$f''(x)$  not defined at  $x = \pm 1$ .

# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}.$

Domain: all  $x$ .

Asymptotes: none. ( $f(x)$  grows like  $x^{4/3}$  as  $x \rightarrow \pm\infty$ .)

Symmetry: about the  $y$ -axis ( $f$  is an even function).

Intercepts:  $(\pm 1, 0), (0, 1)$ .

Critical points:  $x = 0$ ; singular points:  $x = \pm 1$ .

$f''(x) = 0$  at  $x = \pm\sqrt{3}$ ; points  $(\pm\sqrt{3}, 2^{2/3}) \approx (\pm 1.73, 1.59)$ ;

$f''(x)$  not defined at  $x = \pm 1$ .

$x$				SP		CP		SP			
		$-\sqrt{3}$		$-1$		$0$		$1$		$\sqrt{3}$	
$f'$	$-$	$-$	undef	$+$	$0$	$-$	undef	$+$	$+$	$+$	
$f''$	$+$	$0$	$-$	undef	$-$	$-$	undef	$-$	$0$	$+$	
$f$	$\searrow$	$\searrow$	min	$\nearrow$	max	$\searrow$	min	$\nearrow$	$\nearrow$	$\nearrow$	
	$\smile$	infl	$\frown$	$\frown$	$\frown$	$\frown$	$\frown$	$\frown$	infl	$\smile$	

# Sketching the Graph of a Function

**EXAMPLE** Sketch the graph of  $f(x) = (x^2 - 1)^{2/3}$ .

**Solution**  $f'(x) = \frac{4}{3} \frac{x}{(x^2 - 1)^{1/3}}, \quad f''(x) = \frac{4}{9} \frac{x^2 - 3}{(x^2 - 1)^{4/3}}$

Domain: all  $x$ .

Asymptotes: none. ( $f(x)$  grows like  $x^{4/3}$  as  $x \rightarrow \pm\infty$ .)


Symmetry: about the  $y$ -axis ( $f$  is an even function).

Intercepts:  $(\pm 1, 0)$ ,  $(0, 1)$ .

Critical points:  $x = 0$ ; singular points:  $x = \pm 1$ .

$f''(x) = 0$  at  $x = \pm\sqrt{3}$ ; points  $(\pm\sqrt{3}, 2^{2/3}) \approx (\pm 1.73, 1.59)$ ;

$f''(x)$  not defined at  $x = \pm 1$ .



$x$		$-\sqrt{3}$		SP		CP		SP		$\sqrt{3}$	
$f'$	-	-	undef	+	0	-	undef	+	+	+	
$f''$	+	0	-	undef	-	-	undef	-	0	+	
$f$	$\searrow$	$\searrow$	min	$\nearrow$	max	$\searrow$	min	$\nearrow$	$\nearrow$	$\nearrow$	
	$\smile$	infl	$\smile$	$\smile$		$\smile$		$\smile$	infl	$\smile$	


## Extreme-Value Problems

### EXAMPLE

A rectangular animal enclosure is to be constructed having one side along an existing long wall and the other three sides fenced. If 100 m of fence are available, what is the largest possible area for the enclosure?


## Extreme-Value Problems

**EXAMPLE** A rectangular animal enclosure is to be constructed having one side along an existing long wall and the other three sides fenced. If 100 m of fence are available, what is the largest possible area for the enclosure?



## Extreme-Value Problems

**EXAMPLE** A rectangular animal enclosure is to be constructed having one side along an existing long wall and the other three sides fenced. If 100 m of fence are available, what is the largest possible area for the enclosure?




$$x + 2y = 100 \longrightarrow x = 100 - 2y,$$

$$A = A(y) = (100 - 2y)y = 100y - 2y^2.$$



## Extreme-Value Problems

**EXAMPLE** A rectangular animal enclosure is to be constructed having one side along an existing long wall and the other three sides fenced. If 100 m of fence are available, what is the largest possible area for the enclosure?



$$x + 2y = 100 \longrightarrow x = 100 - 2y,$$

$$A = A(y) = (100 - 2y)y = 100y - 2y^2.$$


$0 < y < 50$  (so that area makes sense)

The problem turns into finding the absolute maximum of  $A = 100y - 2y^2$  on  $[0, 50]$ .

## Extreme-Value Problems

### EXAMPLE

A rectangular animal enclosure is to be constructed having one side along an existing long wall and the other three sides fenced. If 100 m of fence are available, what is the largest possible area for the enclosure?



$$A' = 100 - 4y = 0 \Leftrightarrow y = 25$$

The only critical point of  $A$  is  $y = 25$ .  $\Rightarrow$  Largest area  $= A(25)$   
 $= 1250 \text{ m}^2$ .

$$x + 2y = 100 \longrightarrow x = 100 - 2y,$$

$$A = A(y) = (100 - 2y)y = 100y - 2y^2.$$

$0 < y < 50$  (so that area makes sense)


The problem turns into finding the absolute maximum of  $A = 100y - 2y^2$  on  $[0, 50]$ .

## Extreme-Value Problems

### EXAMPLE

A lighthouse  $L$  is located on a small island 5 km north of a point  $A$  on a straight east-west shoreline. A cable is to be laid from  $L$  to point  $B$  on the shoreline 10 km east of  $A$ . The cable will be laid through the water in a straight line from  $L$  to a point  $C$  on the shoreline between  $A$  and  $B$ , and from there to  $B$  along the shoreline. The part of the cable lying in the water costs \$5,000/km, and the part along the shoreline costs \$3,000/km.

Where should  $C$  be chosen to minimize the total cost of the cable?




## Extreme-Value Problems

### EXAMPLE

A lighthouse  $L$  is located on a small island 5 km north of a point  $A$  on a straight east-west shoreline. A cable is to be laid from  $L$  to point  $B$  on the shoreline 10 km east of  $A$ . The cable will be laid through the water in a straight line from  $L$  to a point  $C$  on the shoreline between  $A$  and  $B$ , and from there to  $B$  along the shoreline. The part of the cable lying in the water costs \$5,000/km, and the part along the shoreline costs \$3,000/km.

Where should  $C$  be chosen to minimize the total cost of the cable?






## Extreme-Value Problems

### EXAMPLE

A lighthouse  $L$  is located on a small island 5 km north of a point  $A$  on a straight east-west shoreline. A cable is to be laid from  $L$  to point  $B$  on the shoreline 10 km east of  $A$ . The cable will be laid through the water in a straight line from  $L$  to a point  $C$  on the shoreline between  $A$  and  $B$ , and from there to  $B$  along the shoreline. The part of the cable lying in the water costs \$5,000/km, and the part along the shoreline costs \$3,000/km.

Where should  $C$  be chosen to minimize the total cost of the cable?




$$T = T(x) = 5,000\sqrt{25 - x^2} + 3,000(10 - x), \quad (0 \leq x \leq 10).$$

## Extreme-Value Problems

### EXAMPLE

A lighthouse  $L$  is located on a small island 5 km north of a point  $A$  on a straight east-west shoreline. A cable is to be laid from  $L$  to point  $B$  on the shoreline 10 km east of  $A$ . The cable will be laid through the water in a straight line from  $L$  to a point  $C$  on the shoreline between  $A$  and  $B$ , and from there to  $B$  along the shoreline. The part of the cable lying in the water costs \$5,000/km, and the part along the shoreline costs \$3,000/km.

Where should  $C$  be chosen to minimize the total cost of the cable?



$$0 = \frac{dT}{dx} = \frac{5,000x}{\sqrt{25 + x^2}} - 3,000.$$

$$5,000x = 3,000\sqrt{25 + x^2}$$

$$25x^2 = 9(25 + x^2)$$

$$16x^2 = 225$$

$$x^2 = \frac{225}{16} = \frac{15^2}{4^2} \Rightarrow x = \pm \frac{15}{4}$$


$$T = T(x) = 5,000\sqrt{25 + x^2} + 3,000(10 - x), \quad (0 \leq x \leq 10).$$

## Extreme-Value Problems

### EXAMPLE

A lighthouse  $L$  is located on a small island 5 km north of a point  $A$  on a straight east-west shoreline. A cable is to be laid from  $L$  to point  $B$  on the shoreline 10 km east of  $A$ . The cable will be laid through the water in a straight line from  $L$  to a point  $C$  on the shoreline between  $A$  and  $B$ , and from there to  $B$  along the shoreline. The part of the cable lying in the water costs \$5,000/km, and the part along the shoreline costs \$3,000/km.

Where should  $C$  be chosen to minimize the total cost of the cable?



$$0 = \frac{dT}{dx} = \frac{5,000x}{\sqrt{25 + x^2}} - 3,000.$$

$$5,000x = 3,000\sqrt{25 + x^2}$$

$$25x^2 = 9(25 + x^2)$$

$$16x^2 = 225$$

$$x^2 = \frac{225}{16} = \frac{15^2}{4^2} \Rightarrow x = \pm \frac{15}{4}$$

$$T = T(x) = 5,000\sqrt{25 + x^2} + 3,000(10 - x), \quad (0 \leq x \leq 10).$$

$$T(0) = 55,000, \quad T\left(\frac{15}{4}\right) = 50,000, \quad T(10) \approx 55,902.$$




## Extreme-Value Problems

### EXAMPLE

A lighthouse  $L$  is located on a small island 5 km north of a point  $A$  on a straight east-west shoreline. A cable is to be laid from  $L$  to point  $B$  on the shoreline 10 km east of  $A$ . The cable will be laid through the water in a straight line from  $L$  to a point  $C$  on the shoreline between  $A$  and  $B$ , and from there to  $B$  along the shoreline. The part of the cable lying in the water costs \$5,000/km, and the part along the shoreline costs \$3,000/km.

Where should  $C$  be chosen to minimize the total cost of the cable?



$$0 = \frac{dT}{dx} = \frac{5,000x}{\sqrt{25 + x^2}} - 3,000.$$

$$5,000x = 3,000\sqrt{25 + x^2}$$

$$25x^2 = 9(25 + x^2)$$

$$16x^2 = 225$$

$$x^2 = \frac{225}{16} = \frac{15^2}{4^2} \Rightarrow x = \pm \frac{15}{4}$$

$$T = T(x) = 5,000\sqrt{25 + x^2} + 3,000(10 - x), \quad (0 \leq x \leq 10).$$


$$T(0) = 55,000, \quad T\left(\frac{15}{4}\right) = 50,000, \quad T(10) \approx 55,902.$$



## Extreme-Value Problems

### EXAMPLE


Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



# Extreme-Value Problems

## EXAMPLE

Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



the length  $L$  of the ladder


$$L = L(\theta) = \frac{1}{\cos \theta} + \frac{2}{\sin \theta},$$

where  $0 < \theta < \pi/2$ .

# Extreme-Value Problems

## EXAMPLE

Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



$$\lim_{\theta \rightarrow (\pi/2)^-} L(\theta) = \infty$$

$$\lim_{\theta \rightarrow 0^+} L(\theta) = \infty$$

the length  $L$  of the ladder


$$L = L(\theta) = \frac{1}{\cos \theta} + \frac{2}{\sin \theta},$$

where  $0 < \theta < \pi/2$ .

# Extreme-Value Problems

## EXAMPLE

Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



$$\left[ \begin{array}{l} \lim_{\theta \rightarrow (\pi/2)^-} L(\theta) = \infty \\ \lim_{\theta \rightarrow 0^+} L(\theta) = \infty \end{array} \right] \rightarrow$$

$L(\theta)$  must have a minimum value on  $(0, \pi/2)$ , occurring at a critical point.

the length  $L$  of the ladder


$$L = L(\theta) = \frac{1}{\cos \theta} + \frac{2}{\sin \theta},$$

where  $0 < \theta < \pi/2$ .

# Extreme-Value Problems

## EXAMPLE

Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



$$\left[ \begin{array}{l} \lim_{\theta \rightarrow (\pi/2)^-} L(\theta) = \infty \\ \lim_{\theta \rightarrow 0^+} L(\theta) = \infty \end{array} \right] \rightarrow$$

$L(\theta)$  must have a minimum value on  $(0, \pi/2)$ , occurring at a critical point.

$$0 = L'(\theta) = \frac{\sin \theta}{\cos^2 \theta} - \frac{2 \cos \theta}{\sin^2 \theta} = \frac{\sin^3 \theta - 2 \cos^3 \theta}{\cos^2 \theta \sin^2 \theta}$$

$$\downarrow$$
$$\sin^3 \theta = 2 \cos^3 \theta \longrightarrow \tan^3 \theta = 2$$

the length  $L$  of the ladder


$$L = L(\theta) = \frac{1}{\cos \theta} + \frac{2}{\sin \theta},$$

where  $0 < \theta < \pi/2$ .

# Extreme-Value Problems

## EXAMPLE

Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



$$\left[ \begin{array}{l} \lim_{\theta \rightarrow (\pi/2)^-} L(\theta) = \infty \\ \lim_{\theta \rightarrow 0^+} L(\theta) = \infty \end{array} \right] \rightarrow$$

$L(\theta)$  must have a minimum value on  $(0, \pi/2)$ , occurring at a critical point.

$$0 = L'(\theta) = \frac{\sin \theta}{\cos^2 \theta} - \frac{2 \cos \theta}{\sin^2 \theta} = \frac{\sin^3 \theta - 2 \cos^3 \theta}{\cos^2 \theta \sin^2 \theta}$$

$$\downarrow$$
$$\sin^3 \theta = 2 \cos^3 \theta \longrightarrow \tan^3 \theta = 2$$

the length  $L$  of the ladder

$$L = L(\theta) = \frac{1}{\cos \theta} + \frac{2}{\sin \theta},$$

where  $0 < \theta < \pi/2$ .

$$\sec^2 \theta = 1 + \tan^2 \theta = 1 + 2^{2/3}$$


$$\cos \theta = \frac{1}{(1 + 2^{2/3})^{1/2}}$$

$$\sin \theta = \tan \theta \cos \theta = \frac{2^{1/3}}{(1 + 2^{2/3})^{1/2}}.$$

# Extreme-Value Problems

## EXAMPLE

Find the length of the shortest ladder that can extend from a vertical wall, over a fence 2 m high located 1 m away from the wall, to a point on the ground outside the fence.



$$\left[ \begin{array}{l} \lim_{\theta \rightarrow (\pi/2)^-} L(\theta) = \infty \\ \lim_{\theta \rightarrow 0^+} L(\theta) = \infty \end{array} \right] \rightarrow$$

$L(\theta)$  must have a minimum value on  $(0, \pi/2)$ , occurring at a critical point.

$$0 = L'(\theta) = \frac{\sin \theta}{\cos^2 \theta} - \frac{2 \cos \theta}{\sin^2 \theta} = \frac{\sin^3 \theta - 2 \cos^3 \theta}{\cos^2 \theta \sin^2 \theta}$$

$$\downarrow$$
$$\sin^3 \theta = 2 \cos^3 \theta \longrightarrow \tan^3 \theta = 2$$

the length  $L$  of the ladder

$$L = L(\theta) = \frac{1}{\cos \theta} + \frac{2}{\sin \theta},$$

where  $0 < \theta < \pi/2$ .

$$\sec^2 \theta = 1 + \tan^2 \theta = 1 + 2^{2/3}$$

$$\cos \theta = \frac{1}{(1 + 2^{2/3})^{1/2}}$$

$$\sin \theta = \tan \theta \cos \theta = \frac{2^{1/3}}{(1 + 2^{2/3})^{1/2}}.$$

Therefore the minimal value of  $L(\theta)$  is

$$\begin{aligned} \frac{1}{\cos \theta} + \frac{2}{\sin \theta} &= (1 + 2^{2/3})^{1/2} + 2 \frac{(1 + 2^{2/3})^{1/2}}{2^{1/3}} \\ &= (1 + 2^{2/3})^{3/2} \approx 4.16. \end{aligned}$$