ACTG175: Shrinkage versus non−shrinkage

procedure	successful_repetitions	mean_interval_width
Non−shrinkage	10000	59.6110
Shrinkage	10000	59.9349

ACTG175: Paired comparison (Monte Carlo uncertainty)

metric	estimate	monte_carlo_se	mc95_lower	mc95_upper
Width reduction: non−shrinkage minus shrinkage	−0.3239	2e−04	−0.3243	−0.3235
Mean percentage width reduction	−0.5442	5e−04	−0.5451	−0.5433
Repetitions in which shrinkage is narrower (%)	0.0000	0e+00	0.0000	0.0000

ACTG175: Results by treatment versus control

variance_method	contrast	repetitions	mean_estimate	mean_width
Non−shrinkage	1 vs 0	10000	63.2316	59.9975
Non−shrinkage	2 vs 0	10000	71.2288	59.7348
Non−shrinkage	3 vs 0	10000	49.6884	59.1006
Shrinkage	1 vs 0	10000	63.2316	60.4121
Shrinkage	2 vs 0	10000	71.2288	60.1542
Shrinkage	3 vs 0	10000	49.6884	59.2383

ACTG175: 10000 repetitions; 40 regression imputations each; identical imputations for both procedures.

How wide were the intervals?



Shrinkage was 0.54% wider on average across repetitions.

Same patients and natural missing values in every repetition; only imputation draws change.
Width comparisons assess imputation stability and precision; recovery of unknown outcomes cannot be measured.