ACTG175: results in the layout of thesis Table 4.2

Table 1. Descriptive averages of simultaneous inference results over 10,000 repetitions.

Treatment–control comparison	Mean point estimate	Mean lower / upper 95% interval limits*	Mean interval length
Proposed procedure without shrinkage			
1 vs 0	63.232	(33.233, 93.230)	59.998
2 vs 0	71.229	(41.361, 101.096)	59.735
3 vs 0	49.688	(20.138, 79.239)	59.101
Proposed procedure with shrinkage			
1 vs 0	63.232	(33.026, 93.438)	60.412
2 vs 0	71.229	(41.152, 101.306)	60.154
3 vs 0	49.688	(20.069, 79.308)	59.238

* The limits above are averages of the 10,000 separately computed interval endpoints. They are descriptive summaries, not a single 95% confidence interval.

Each repetition used 40 regression imputations shared by the shrinkage and non-shrinkage procedures. Natural missing values retained; repetitions vary only regression-imputation draws.

The table layout follows thesis Table 4.2; only the two procedures requested by the user are shown. No equal-variance or imputation-method comparison is added.

ACTG175: one actual 40-imputation application analysis

Table 2. Simultaneous statistical inferences from the first prespecified repetition.

Treatment–control comparison	Point estimate	95% simultaneous confidence interval	Interval length
Proposed procedure without shrinkage			
1 vs 0	63.329	(34.587, 92.072)	57.485
2 vs 0	71.330	(43.176, 99.483)	56.307
3 vs 0	48.162	(18.603, 77.721)	59.118
Proposed procedure with shrinkage			
1 vs 0	63.329	(34.332, 92.327)	57.995
2 vs 0	71.330	(42.961, 99.698)	56.736
3 vs 0	48.162	(18.549, 77.775)	59.226

This page follows the application meaning of thesis Table 4.2: each displayed interval is an actual simultaneous 95% interval from one 40-imputation analysis. It uses repetition 1, selected by its prespecified seed—not by its results.

Table 2 illustrates the application table; it does not replace the full 10,000-repetition results on page 1. These two types of results must not be interpreted interchangeably.