

MY
Sem-P™
// (Semaglutide)
DIARY

SUPPORTING YOUR
Sem-P™
/(Semaglutide)
JOURNEY

Table of Contents

- Know your Sem-P™
 - Weekly Sem-P™ Dosing Card
 - Sem-P™ Dose Calculation
 - How to Use Sem-P™ Pen
 - Sem-P™ Pen Storage Conditions
 - Sem-P™ Rotation Guide for Injection Site
 - Sem-P™ Common Side Effects
 - Managing Side Effects with Sem-P™
 - Sem-P™ Meal Plan
 - Glucose Monitoring with Sem-P™
- 

Know Your **Sem P**TM //(Semaglutide)

Sem PTM injection 2 mg or 4 mg is an injectable prescription medicine used along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes. To reduce the risk of major cardiovascular events such as heart attack, stroke or death in adults with type 2 diabetes and with known heart disease. It is not known if **Sem P**TM can be used in people who have had pancreatitis. **Sem P**TM is not for use in people with type 1 diabetes. It is not known if **Sem P**TM is safe and effective for use in children under 18 years of age.

Weekly Dosing Card

Choose Your Sem-P™ Day

1st Dose	2nd Dose	3rd Dose	4th Dose

	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday		Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
Dose Day	Sem-P INJECTOR DAY							Dose Day	Sem-P INJECTOR DAY							Sem-P INJECTOR DAY
	✓								✗	Inject your dose						
If Missed								If Missed		Inject your dose					Skip your dose	✓

If you missed your dose on given day, you can take next dose within five days. Otherwise skip the dose and take on selected dose day.

Dose Calculation

Week 1-4

Initiate
(0.25 mg)
Each Week



Week 5-8

Escalate & maintain
(0.5 mg)
Each Week



Week 9+

Maintain (1 mg)
Each Week




Dose in gram	Frequency	Duration
Initiate with 0.25 mg	Once Weekly	Continue for 4 weeks
Escalate & maintain to 0.5 mg	Once Weekly	Continue for 4 weeks
Maintain at 1.0 mg	Once Weekly	Continue



If you need more information
contact us **021-111-455-455**

How to Use


Sem-PTM Pen



Wash Your Hands With Soap And Water.


1

Pull off the pen cap.




2

Check that Sem-P injection in your pen is clear and colorless. If Sem-P injection looks cloudy or contains particles, do not use the pen.




3

Take a new needle, and tear off the paper tab. Keep the needle straight and screw it onto the pen until fix. Do not over tightened.



4

Pull off the needle cap and keep it for later use.




5

Make sure that there is no any air bubble into the Sem-P injection.


First Time Use for Each New Pen: Check the Sem-P flow

- Turn the dose selector until the dose counter shows the flow check symbol (↔).
- Hold the pen with the needle pointing up.
- Press and hold the button until the dose selector shows 0. A drop of Sem-P injection will appear at the needle tip.
- If no drop appears, repeat Step (a),(b) & (c) up to 6 times. If there is still no drop change the needle and repeat Step (a),(b) & (c), 1 more time.
- Do not use the pen if a drop of Sem-P injection still does not appear.




6

Choose your injection site. Sem-P Solution for Injection can be injected under the skin (subcutaneously) of your stomach area (abdomen) or thighs, and with the help of someone else, in the back of the upper arm.




7

To inject, press the injection button completely in, until you hear or feel a click. Leave the needle under the skin for at least 10 seconds. This ensures that the selected Sem-P dose is delivered. Withdraw the needle. You have completed your injection and the selected Sem-P dose has been delivered. The dose selector display will show '0'.




8

After you have withdrawn the needle, a few droplets of Sem-P injection may appear at the needle tip. This is normal and has no effect on the dose delivered.



9


Select the dose you need to inject by turning the dose selector until your required dose (0.25mg or 0.5mg or 1 mg) lines up with the dose pointer.



10


If you select a different dose than you need, simply turn back the dose selector until the correct dose lines up with the dose pointer.

Only doses of 0.25mg & 0.5mg can be selected with the dose selector for the Sem-P 2mg/1.5mL pen & 1mg can be selected for 4mg/3mL pen.




11

Carefully replace the needle cap onto the needle.




12

Hold the Pen holder and unscrew the needle. Dispose off the needle properly as instructed by your healthcare professional.



13

Always firmly replace the pen cap after use.



Storage Conditions

Storing Sem-P[™] Pen

- ~ Store all opened and unopened **Pen** in the refrigerator at (2°C to 8°C).
- ~ Do not freeze. Do not use if Sem-P[™] **Pen** has been frozen.
- ~ Store the **Pen** in the original carton to protect from light.
- ~ Keep Sem-P[™] **Pen**, needles, and all medicines out of the reach of children.

Rotation Guide for Injection Site



Choose
injection site



Divide area
into four sections




Choose one
section each week




One finger width
from last site

Common Side Effects



Decreased
Appetite



Stomach
Pain



Nausea &
Vomiting



Diarrhea



Constipation



If you need more information
contact us **021-111-455-455**

Managing Side Effects

Eating smaller, more frequent meals

Stop to avoid overeating when full



Choosing light foods & avoiding spicy & fatty foods.

Staying hydrated with plenty of water throughout the day & green tea once or twice a day.

Mild foods includes:

Plain rice, pasta, or oatmeal or boiled potatoes without seasoning. Skinless chicken breast or other lean meats or non-citrus fruits like bananas. Low-fat dairy products like yogurt or cottage cheese (if tolerated)



Fresh Vegetables

Half your plate should be filled with non starchy vegetables such as cucumber, broccoli, Spanish, pumpkin, ridge gourd (turai).



Nutritious Carbohydrate

Fill one-quarter with foods like brown rice, oats, brown bread or corn

Sem P™
/// (Semaglutide) ///

Friendly Meal Plan

Fresh Fruit Healthy Fats


You should also try to eat a piece of fruit with each meal or few teaspoons of a healthy fat



Protein

One-quarter should be filled with protein, such as chicken, fish, lean meat or eggs





Sem.PTM
 (Semaglutide)

Friendly Diet

Foods diabetic patients can eat

Pumpkin 	Bitter Gourd 	Okra 
Turnip 	Salad Leaves 	Eggplant 
Cucumber 	Lemon, GreenChilli 	Spring Onion 
Broth 	Salad 	Egg White 

Foods diabetic patients can eat as advised

Rice 	Roti 	Bread 
Fish 	Noodles 	Pulses 
Peach 	Sweet Lime Orange 	Chicken 
Apple 	Papaya 	Watermelon 
Diet Toffee 	Carrot 	Banana 
Margarine 	Skimmed Milk 	Beans Peas 
Yogurt 	Paneer 	Egg 
Porridge 	Diet Toffee 	
Cooking Oil 		

Foods diabetic patients should avoid

Pudding 	Ice Cream 	Cake Pastry 
Dried Fruits 	Honey 	Jam Jelly 
Dates 	Mithai 	FruitJuice / ColdDrinks 
NanKhatai / BakeryBiscuit 	Chewing Gum Chocolate 	
	Condensed Milk 	

Glucose Screening

Days	Week 1		Week 2		Week 3		Week 4	
	FBS	RBS	FBS	RBS	FBS	RBS	FBS	RBS
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

***FBS:** Fasting Blood Sugar (taken after 12 hrs of fasting)

***RBS:** Random Blood Sugar (taken after 2 hrs of meal)

Glucose Screening

Days	Week 1		Week 2		Week 3		Week 4	
	FBS	RBS	FBS	RBS	FBS	RBS	FBS	RBS
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

***FBS:** Fasting Blood Sugar (taken after 12 hrs of fasting)

***RBS:** Random Blood Sugar (taken after 2 hrs of meal)

Glucose Screening

Days	Week 1		Week 2		Week 3		Week 4	
	FBS	RBS	FBS	RBS	FBS	RBS	FBS	RBS
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

***FBS:** Fasting Blood Sugar (taken after 12 hrs of fasting)

***RBS:** Random Blood Sugar (taken after 2 hrs of meal)

Glucose Screening

Days	Week 1		Week 2		Week 3		Week 4	
	FBS	RBS	FBS	RBS	FBS	RBS	FBS	RBS
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

***FBS:** Fasting Blood Sugar (taken after 12 hrs of fasting)

***RBS:** Random Blood Sugar (taken after 2 hrs of meal)

Glucose Screening

Days	Week 1		Week 2		Week 3		Week 4	
	FBS	RBS	FBS	RBS	FBS	RBS	FBS	RBS
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

***FBS:** Fasting Blood Sugar (taken after 12 hrs of fasting)

***RBS:** Random Blood Sugar (taken after 2 hrs of meal)

Glucose Screening

Days	Week 1		Week 2		Week 3		Week 4	
	FBS	RBS	FBS	RBS	FBS	RBS	FBS	RBS
Sunday								
Monday								
Tuesday								
Wednesday								
Thursday								
Friday								
Saturday								

***FBS:** Fasting Blood Sugar (taken after 12 hrs of fasting)
***RBS:** Random Blood Sugar (taken after 2 hrs of meal)