


# Rings

August 12, 2024



## Product of Ideals

### Product of Ideals (without finite sums)

Given two ideals  $I$  and  $J$  in a ring  $R$ , the product  $IJ$  without finite sums is defined as:

$$IJ \triangleq \{ab : a \in I, b \in J\}.$$

[Example in  $\mathbb{Z}$ ] Consider  $I = 2\mathbb{Z}$  and  $J = 3\mathbb{Z}$  in  $\mathbb{Z}$ :

$$2\mathbb{Z} = \{2n : n \in \mathbb{Z}\},$$

$$3\mathbb{Z} = \{3m : m \in \mathbb{Z}\}.$$

By definition,

$$IJ = \{2n \cdot 3m : n, m \in \mathbb{Z}\} = \{6nm : n, m \in \mathbb{Z}\}.$$

Consider two general elements  $6n_1m_1$  and  $6n_2m_2$  in  $IJ$ . Their sum is

$$6n_1m_2 + 6n_2m_2 = 6(n_1m_1 + n_2m_2) = 6 \sum_{k=1}^2 n_k m_k \stackrel{?}{\in} IJ = \{6nm : n, m \in \mathbb{Z}\}.$$

Without finite sums, we only have products  $6ab$ .

### Product of Ideals (with finite sums)

$$IJ \triangleq \left\{ \sum_{k=1}^t a_k b_k : a_k \in I, b_k \in J, t \in \mathbb{N} \right\}.$$

Consider  $x = \sum_{k=1}^s 6a_k b_k$  and  $y = \sum_{k=1}^t 6a_k b_k$ . Then

$$\begin{aligned} x + y &= \sum_{k=1}^s 6a_k b_k + \sum_{l=1}^t 6a_l b_l = 6 \left( \sum_{k=1}^s a_k b_k + \sum_{k=1}^t a_k b_k \right) \\ &= 6 \sum_{k=1}^{s+t} a_k b_k \end{aligned}$$

[Example in  $\mathbb{C}[x]$ ]

### Product of Ideals (without finite sums)

Consider  $I = (x)$  and  $J = (x^2)$  in  $\mathbb{C}[x]$ :

$$\begin{aligned}(x) &= \{x \cdot p(x) : p(x) \in \mathbb{C}[x]\}, \\(x^2) &= \{x^2 \cdot q(x) : q \in \mathbb{C}[x]\}.\end{aligned}$$

By definition,

$$IJ = \{x \cdot p(x) \cdot x^2 \cdot q(x) : p(x), q(x) \in \mathbb{C}[x]\} = \{x^3 p(x) q(x) : p(x), q(x) \in \mathbb{C}[x]\}.$$

Consider two general elements  $x^3 p_1(x) q_1(x)$  and  $x^3 p_2(x) q_2(x)$  in  $IJ$ . Their sum is

$$x^3 p_1(x) q_1(x) + x^3 p_2(x) q_2(x) = x^3 (p_1(x) q_1(x) + p_2(x) q_2(x)) = x^3 \sum_{i=1}^2 p_i(x) q_i(x).$$

### Product of Ideals (with finite sums)

$$IJ \triangleq \left\{ \sum_{k=1}^t a_k b_k : a_k \in I, b_k \in J, t \in \mathbb{N} \right\}.$$

Consider  $x = \sum_{k=1}^s 6a_k b_k$  and  $y = \sum_{k=1}^t 6a_k b_k$ . Then

$$\begin{aligned}x + y &= \sum_{k=1}^s 6a_k b_k + \sum_{l=1}^t 6a_l b_l = 6 \left( \sum_{k=1}^s a_k b_k + \sum_{k=1}^t a_k b_k \right) \\&= 6 \sum_{k=1}^{s+t} a_k b_k\end{aligned}$$

[Example in  $\mathbb{Z}[x]$ ] Consider  $I = (2, x)$  and  $J = (3, x^2)$  in  $\mathbb{Z}[x]$ :

$$(2, x) = \{2f(x) + xg(x) : f, g \in \mathbb{Z}[x]\},$$

$$(3, x^2) = \{3h(x) + x^2k(x) : h, k \in \mathbb{C}[x]\}.$$

Let

$$IJ \triangleq \{(2f + xg)(3h + x^2k) : f, g, h, k \in \mathbb{Z}[x]\}.$$

Consider


$$x_1 = (2 + x)(3 + x^2) \in IJ,$$


$$x_2 = (2x + x^3)(3 + x^2) \in IJ.$$


Then


$$x_1 = (2 + x)(3 + x^2) = 6 + 2x^2 + 3x + x^3,$$

$$x_2 = 6x + 2x^3 + 3x^3 + x^5 = 6x + 5x^3 + x^5.$$











$$IJ = \left\{ \sum_{k=1}^t (a_k b_k) \mid a_k, b_k \in \mathbb{Z}, t \in \mathbb{N} \right\}$$



$$IJ = \left\{ \sum_{k=1}^t (a_k b_k) \mid a_k, b_k \in \mathbb{Z}, t \in \mathbb{N} \right\}$$

$$\mathbb{Z}\mathbb{Z} = \left\{ \sum_{k=1}^t (a_k)(b_k) : a_k, b_k \in \mathbb{Z}, t \in \mathbb{N} \right\}$$





Ring (Set)	Addition Operation	Multiplication Operation	Additive Identity	Multiplicative Identity	Form of Element	Commutative
Integers $\mathbb{Z}$	Standard addition +	Standard multiplication $\cdot$	0	1	Integers	Yes
Real Numbers $\mathbb{R}$	Standard addition +	Standard multiplication $\cdot$	0	1	Real numbers	Yes
Complex Numbers $\mathbb{C}$	Standard addition +	Standard multiplication $\cdot$	0	1	$a + bi$ where $a, b \in \mathbb{R}$	Yes
Polynomials with Real Coefficients $\mathbb{R}[x]$	Polynomial addition	Polynomial multiplication	0 (zero polynomial)	1 (constant polynomial)	$a_n x^n + \dots + a_1 x + a_0$	Yes
Matrices $M_n(\mathbb{R})$	Matrix addition	Matrix multiplication	Zero matrix	Identity matrix	$n \times n$ matrices	No
Integers Modulo $n$ $\mathbb{Z}/n\mathbb{Z}$	Addition modulo $n$	Multiplication modulo $n$	0	1	$\{0, 1, \dots, n-1\}$	Yes

Group (Set)	Operation	Identity	Form of Element	Normal Subgroups	Abelian
Symmetric Group $S_3$	Composition of permutations	Identity permutation $e$	Permutations of 3 elements	$\{e, (123), (132)\},$ $\{e, (12), (13), (23), (123), (132)\}$	No
Dihedral Group $D_4$	Composition of symmetries	Identity symmetry $e$	Rotations $r^k$ and reflections $sr^k$ , $k \in \{0, 1, 2, 3\}$	$\{e, r, r^2, r^3\},$ $\{e, r^2, s, sr^2\}$	No
Quaternion Group $Q_8$	Quaternion multiplication	1	Quaternions $\pm 1, \pm i, \pm j, \pm k$	$\{1, -1\}, \{1, -1, i, -i\},$ $\{1, -1, j, -j\},$ $\{1, -1, k, -k\}$	No
Integers Modulo $n$ $\mathbb{Z}/n\mathbb{Z}$	Addition modulo $n$	0	$\{0, 1, \dots, n-1\}$	All subgroups are normal	Yes

Ring (Set)	Addition Operation	Multiplication Operation	Additive Identity	Multiplicative Identity	Form of Element
Matrices $M_n(\mathbb{R})$	Matrix addition	Matrix multiplication	Zero matrix	Identity matrix	$n \times n$ matrices
Quaternions $\mathbb{H}$	Quaternion addition	Quaternion multiplication	0	1	$a + bi + cj + dk$ where $a, b, c, d \in \mathbb{R}$
Differential Operators	Operator addition	Operator composition	Zero operator	Identity operator	$\sum a_i \frac{d^i}{dx^i}$
Group Rings $\mathbb{R}[G]$	Group ring addition	Group ring multiplication	Zero element	Identity element	$\sum a_g g$ where $a_g \in \mathbb{R}$ and $g \in G$
Endomorphism Rings $\text{End}(V)$	Function addition	Function composition	Zero map	Identity map	Linear transformations on vector space $V$
Octonions $\mathbb{O}$	Octonion addition	Octonion multiplication	0	1	$a + be_1 + ce_2 + de_3 + ee_4 + fe_5 + ge_6 + he_7$ where $a, b, c, d, e, f, g, h \in \mathbb{R}$

Type	Ideal	Principal Ideal	Prime Ideal	Maximal Ideal
<b>Definition</b>	A subset closed under addition and multiplication by any ring element	An ideal generated by a single element	An ideal where if $ab \in I$ , then $a \in I$ or $b \in I$	An ideal such that there are no larger ideals except the ring itself
Is an Ideal	O	O	O	O
Can be Principal	O	O	O	O
Is Prime	X	X	O	O
Is Maximal	X	X	X	O

Ring (Set)	Principal Ideals	Prime Ideals	Maximal Ideals
$\mathbb{Z}$	$(2), (3), (5)$	$(2), (3), (5)$	$(2), (3), (5)$
$\mathbb{R}[x]$	$(x), (x-1), (x^2+1)$	$(x)$	$(x-1)$
$\mathbb{C}[x]$	$(x), (x-i), (x+i)$	$(x-i), (x+i)$	$(x-i), (x+i)$
$\mathbb{Z}/6\mathbb{Z}$	$(2), (3)$	None	$(2), (3)$
$\mathbb{Z}[i]$	$(1+i), (2)$	$(1+i)$	$(1+i)$
$\mathbb{Z}/p\mathbb{Z}$ where $p$ is prime	$(0), (1)$	$(0)$	$(0)$
$\mathbb{R}$	$(0)$	None	$(0)$
$\mathbb{C}$	$(0)$	None	$(0)$
$\mathbb{Z}[x]$	$(2), (3), (x)$	$(x), (2)$	None
$M_n(\mathbb{R})$	$(E_{11})$	None	None
$\mathbb{H}$ (Quaternions)	$(1+i)$	None	None

Ring (Set)	Principal Ideals	Prime Ideals	Maximal Ideals
$\mathbb{Z}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$
$\mathbb{R}[x]$	$(x) = \{xf(x) \mid f(x) \in \mathbb{R}[x]\}$ $(x-1) = \{(x-1)f(x) \mid f(x) \in \mathbb{R}[x]\}$ $(x^2+1) = \{(x^2+1)f(x) \mid f(x) \in \mathbb{R}[x]\}$	$(x) = \{xf(x) \mid f(x) \in \mathbb{R}[x]\}$	$(x-1) = \{(x-1)f(x) \mid f(x) \in \mathbb{R}[x]\}$
$\mathbb{C}[x]$	$(x) = \{xf(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x-i) = \{(x-i)f(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x+i) = \{(x+i)f(x) \mid f(x) \in \mathbb{C}[x]\}$	$(x-i) = \{(x-i)f(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x+i) = \{(x+i)f(x) \mid f(x) \in \mathbb{C}[x]\}$	$(x-i) = \{(x-i)f(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x+i) = \{(x+i)f(x) \mid f(x) \in \mathbb{C}[x]\}$
$\mathbb{Z}/6\mathbb{Z}$	$(2) = \{0, 2, 4\}$ $(3) = \{0, 3\}$	None	$(2) = \{0, 2, 4\}$ $(3) = \{0, 3\}$
$\mathbb{Z}[i]$	$(1+i) = \{(1+i)z \mid z \in \mathbb{Z}[i]\}$ $(2) = \{2z \mid z \in \mathbb{Z}[i]\}$	$(1+i) = \{(1+i)z \mid z \in \mathbb{Z}[i]\}$	$(1+i) = \{(1+i)z \mid z \in \mathbb{Z}[i]\}$
$\mathbb{Z}/p\mathbb{Z}$ where $p$ is prime	$(0) = \{0\}$ $(1) = \mathbb{Z}/p\mathbb{Z}$	$(0) = \{0\}$	$(0) = \{0\}$
$\mathbb{R}$	$(0) = \{0\}$	None	$(0) = \{0\}$
$\mathbb{C}$	$(0) = \{0\}$	None	$(0) = \{0\}$
$\mathbb{Z}[x]$	$(2) = \{2f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(3) = \{3f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(x) = \{xf(x) \mid f(x) \in \mathbb{Z}[x]\}$	$(2) = \{2f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(x) = \{xf(x) \mid f(x) \in \mathbb{Z}[x]\}$	None
$M_n(\mathbb{R})$	$(E_{11}) = \{AE_{11}B \mid A, B \in M_n(\mathbb{R})\}$	None	None
$\mathbb{H}$ (Quaternions)	$(1+i) = \{(1+i)q \mid q \in \mathbb{H}\}$	None	None

Ring (Set)	Principal Ideals	Prime Ideals	Maximal Ideals
$\mathbb{Z}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$
$\mathbb{R}[x]$	$(x) = \{xf(x) \mid f(x) \in \mathbb{R}[x]\}$ $(x-1) = \{(x-1)f(x) \mid f(x) \in \mathbb{R}[x]\}$ $(x^2+1) = \{(x^2+1)f(x) \mid f(x) \in \mathbb{R}[x]\}$	$(x)$	$(x-1)$
$\mathbb{C}[x]$	$(x) = \{xf(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x-i) = \{(x-i)f(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x+i) = \{(x+i)f(x) \mid f(x) \in \mathbb{C}[x]\}$	$(x-i)$  $(x+i)$	$(x-i)$  $(x+i)$
$\mathbb{Z}/6\mathbb{Z}$	$(2) = \{0, 2, 4\}$ $(3) = \{0, 3\}$	None	$(2) = \{0, 2, 4\}$ $(3) = \{0, 3\}$
$\mathbb{Z}[i]$	$(1+i) = \{(1+i)z \mid z \in \mathbb{Z}[i]\}$ $(2) = \{2z \mid z \in \mathbb{Z}[i]\}$	$(1+i)$	$(1+i)$
$\mathbb{Z}/p\mathbb{Z}$ where $p$ is prime	$(0) = \{0\}$ $(1) = \mathbb{Z}/p\mathbb{Z}$	$(0)$	$(0)$
$\mathbb{R}$	$(0) = \{0\}$	None	$(0)$
$\mathbb{C}$	$(0) = \{0\}$	None	$(0)$
$\mathbb{Z}[x]$	$(2) = \{2f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(3) = \{3f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(x) = \{xf(x) \mid f(x) \in \mathbb{Z}[x]\}$	$(2)$  $(x)$	None
$M_n(\mathbb{R})$	$(E_{11}) = \{AE_{11}B \mid A, B \in M_n(\mathbb{R})\}$	None	None
$\mathbb{H}$ (Quaternions)	$(1+i) = \{(1+i)q \mid q \in \mathbb{H}\}$	None	None

Ring (Set)	Examples of Ideals	Examples of Principal Ideals	Examples of Prime Ideals	Examples of Maximal Ideals
$\mathbb{Z}$	$(0), (2), (3)$ $(4), (6)$ $(0)$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$	$(2) = \{2k \mid k \in \mathbb{Z}\}$ $(3) = \{3k \mid k \in \mathbb{Z}\}$ $(5) = \{5k \mid k \in \mathbb{Z}\}$
$\mathbb{R}[x]$	$(0), (x), (x-1)$ $(x^2+1)$	$(x) = \{xf(x) \mid f(x) \in \mathbb{R}[x]\}$ $(x-1) = \{(x-1)f(x) \mid f(x) \in \mathbb{R}[x]\}$ $(x^2+1) = \{(x^2+1)f(x) \mid f(x) \in \mathbb{R}[x]\}$	$(x)$	$(x-1)$
$\mathbb{C}[x]$	$(0), (x), (x-i)$ $(x+i)$	$(x) = \{xf(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x-i) = \{(x-i)f(x) \mid f(x) \in \mathbb{C}[x]\}$ $(x+i) = \{(x+i)f(x) \mid f(x) \in \mathbb{C}[x]\}$	$(x-i)$ $(x+i)$	$(x-i)$ $(x+i)$
$\mathbb{Z}/6\mathbb{Z}$	$(0), (2), (3)$	$(2) = \{0, 2, 4\}$ $(3) = \{0, 3\}$	None	$(2) = \{0, 2, 4\}$ $(3) = \{0, 3\}$
$\mathbb{Z}[i]$	$(0), (1+i), (2)$	$(1+i) = \{(1+i)z \mid z \in \mathbb{Z}[i]\}$ $(2) = \{2z \mid z \in \mathbb{Z}[i]\}$	$(1+i)$	$(1+i)$
$\mathbb{Z}/p\mathbb{Z}$ where $p$ is prime	$(0), (1)$	$(0) = \{0\}$ $(1) = \mathbb{Z}/p\mathbb{Z}$	$(0)$	$(0)$
$\mathbb{R}$	$(0)$	$(0) = \{0\}$	None	$(0)$
$\mathbb{C}$	$(0)$	$(0) = \{0\}$	None	$(0)$
$\mathbb{Z}[x]$	$(0), (2), (3)$	$(2) = \{2f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(3) = \{3f(x) \mid f(x) \in \mathbb{Z}[x]\}$ $(x) = \{xf(x) \mid f(x) \in \mathbb{Z}[x]\}$	$(2)$  $(x)$	None
$M_n(\mathbb{R})$	$(0), (E_{11})$	$(E_{11}) = \{AE_{11}B \mid A, B \in M_n(\mathbb{R})\}$	None	None
$\mathbb{H}$ (Quaternions)	$(0), (1+i)$	$(1+i) = \{(1+i)q \mid q \in \mathbb{H}\}$	None	None