


# Riemann Surfaces and Algebraic Curves

A framework for understanding Elliptic Curves

Ji, Yonghyeon

PART I — MULTIVARIABLE CALCULUS




# Riemann Surfaces and Algebraic Curves

*A Framework for Understanding Elliptic Curves*

## Part I — Multivariable Calculus

Ji, Yonghyeon

December 29, 2025



WINTER 2026

## The FTC hierarchy

Name	Formula
FTC I (Accumulation)	$\frac{d}{dx} \left( \int_a^x f(t) dt \right) = f(x).$
FTC II (Evaluation)	$\int_a^b f'(x) dx = f(b) - f(a).$
Fundamental Theorem of Line Integrals	$\int_C \nabla \phi \cdot d\mathbf{r} = \phi(B) - \phi(A).$
Green's Theorem	$\oint_{\partial R} P dx + Q dy = \iint_R \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$
Stokes' Theorem (3D)	$\oint_{\partial S} \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} dS.$
Divergence Theorem	$\iint_{\partial V} \mathbf{F} \cdot \mathbf{n} dS = \iiint_V (\nabla \cdot \mathbf{F}) dV.$
Generalized Stokes	$\int_{\partial \Omega} \omega = \int_{\Omega} d\omega.$

## Copyright

Copyright © 2025 by Ji, Yonghyeon All rights reserved.

No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the publisher, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law.

## Changelog

v1.0 2025-12-29 Initial release.

# Contents

<b>1</b>	<b>Fundamental Theorem of Calculus</b>	<b>3</b>
1.1	Gradient Vector Fields . . . . .	4
1.2	Green's Theorem . . . . .	4
1.3	Divergence Theorem . . . . .	6
1.4	Stokes' Theorem . . . . .	6
<b>2</b>	<b>Differential Forms</b>	<b>7</b>
<b>3</b>	<b>Winding Numbers and Complexification</b>	<b>8</b>

# 1 Fundamental Theorem of Calculus

## Fundamental Theorem for Gradient Fields

If  $\mathbf{F} = \nabla f$  is a conservative vector field and  $C$  is a smooth curve from  $A$  to  $B$ , then

$$\int_C \mathbf{F} \cdot d\mathbf{r} = f(B) - f(A).$$

## Green's Theorem

For a positively oriented, simple closed curve  $C$  bounding a region  $R$  in the plane,

$$\oint_C P dx + Q dy = \iint_R \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

## Divergence Theorem

Let  $\mathbf{F}$  be a vector field defined on a region  $E$  with closed boundary surface  $S$  (outward-oriented). Then

$$\iiint_E \nabla \cdot \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} dS.$$

## Stokes' Theorem

Let  $S$  be an oriented surface with boundary curve  $C$ , and let  $\mathbf{F}$  be a vector field. Then

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}.$$

## Triple Integral

To integrate a scalar function  $f(x, y, z)$  over a region  $E$  in  $\mathbb{R}^3$ ,

$$\iiint_E f(x, y, z) dV.$$

## 1.1 Gradient Vector Fields

1. Let  $\mathbf{F} = \langle 2x, 2y \rangle$ . Show that  $\mathbf{F}$  is conservative and compute

$$\int_C \mathbf{F} \cdot d\mathbf{r}$$

where  $C$  is any path from  $(0, 0)$  to  $(1, 1)$ .

2. Determine whether the vector field  $\mathbf{F} = \langle y, x \rangle$  is conservative. If so, find a potential function.
3. Let  $f(x, y, z) = xyz$ . Compute  $\nabla f$  and evaluate the line integral of  $\nabla f$  over the path from  $(1, 0, 0)$  to  $(1, 2, 3)$ .
4. Let  $\mathbf{F} = \nabla f$  for  $f(x, y) = x^2 + y^2$ . Compute the line integral over a circular path from  $(1, 0)$  to  $(0, 1)$  and explain the result.

## 1.2 Green's Theorem

1. Use Green's Theorem to evaluate

$$\oint_C x \, dy - y \, dx$$

where  $C$  is the unit circle oriented counterclockwise.

2. Let  $\mathbf{F} = \langle y^2, 2xy \rangle$ . Use Green's Theorem to evaluate the line integral around the boundary of the square  $[0, 1] \times [0, 1]$ .
3. Evaluate

$$\oint_C (x + y)dx + (x - y)dy$$

where  $C$  is the triangle with vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(1, 1)$  oriented counterclockwise.

4. Determine if

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

for  $\mathbf{F} = \langle y, -x \rangle$  around a circle of radius  $r$  centered at the origin.

### 1.3 Divergence Theorem

1. Let  $\mathbf{F} = \langle x, y, z \rangle$ . Use the Divergence Theorem to compute the flux across the surface of the unit sphere.
2. Let  $\mathbf{F} = \langle x^2, y^2, z^2 \rangle$ . Compute both the divergence and the surface integral over the unit cube  $[0, 1]^3$ .
3. Use the Divergence Theorem to find the outward flux of  $\mathbf{F} = \langle yz, xz, xy \rangle$  through the unit cube.
4. Let  $\mathbf{F} = \langle x, -y, z \rangle$ . Verify the Divergence Theorem on the upper hemisphere of radius 1 centered at the origin.

### 1.4 Stokes' Theorem

1. Let  $\mathbf{F} = \langle -y, x, 0 \rangle$ . Use Stokes' Theorem to compute the circulation around the boundary of the disk  $x^2 + y^2 \leq 1$  in the  $xy$ -plane.
2. Let  $\mathbf{F} = \langle z, 0, x \rangle$ . Use Stokes' Theorem on the triangular surface with vertices at  $(0, 0, 0)$ ,  $(1, 0, 0)$ ,  $(0, 1, 0)$ .
3. Compute both sides of Stokes' Theorem for  $\mathbf{F} = \langle y, z, x \rangle$  on the surface  $z = 0$  bounded by the unit circle.
4. Use Stokes' Theorem to show that

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

if  $\mathbf{F}$  is the gradient of some scalar field  $f$ .

## 2 Differential Forms

TBA



### 3 Winding Numbers and Complexification

TBA