

# Abstract Algebra II

Ji, Yong-hyeon

May 4, 2025

We cover the following topics in this note.

- Group Action
  - Cayley Theorem
  - Normal Subgroups
  - TBA
- 

## Contents

### Group Action

**Definition.** Let  $(G, *)$  be a group and let  $X \neq \emptyset$ . A **(left) group action** of  $G$  on  $X$  is a function

$$\cdot : G \times X \rightarrow X, \quad (g, x) \mapsto g \cdot x$$

satisfying the followings: for all  $g, h \in G$  and all  $x \in X$ ,

- (i) (Identity)  $e \cdot x = x$ , where  $e \in G$  is the identity element of  $G$ ;
- (ii) (Compatibility)  $(g * h) \cdot x = g \cdot (h \cdot x)$ .

The pair  $(X, \cdot)$  (or simply  $X$ ) is then called a  $G$ -set.

**Note (Notation).** If a group  $G$  acts on a set  $X$ , one commonly writes:  $G \curvearrowright X$ .

**Remark.** A right group action of  $G$  on  $X$  is a function  $\cdot : X \times G \rightarrow X, \quad (x, g) \mapsto x \cdot g$  satisfying:

- (i)  $x \cdot e = x$  for all  $x \in X$ ;
- (ii)  $(x \cdot g) \cdot h = x \cdot (gh)$  for all  $g, h \in G, x \in X$ .

**Example** (Scalar Multiplication on a Vector Space). Let  $\mathbb{F}$  be a field, and let  $X = \mathbb{F}^n$  be the  $n$ -dimensional vector space over  $\mathbb{F}$ . Consider the multiplicative group of nonzero scalars in  $\mathbb{F}$ :


$$G = (\mathbb{F}^\times, \times), \quad \text{where } \mathbb{F}^\times = \mathbb{F} \setminus \{0\}.$$

We define an action  $G \curvearrowright X$  by scalar multiplication:

$$\begin{aligned} \cdot : \mathbb{F}^\times \times \mathbb{F}^n &\longrightarrow \mathbb{F}^n \\ (\lambda, \mathbf{v}) &\longmapsto \lambda \cdot \mathbf{v} \end{aligned}$$

where the product  $\lambda \cdot \mathbf{v}$  is defined componentwise. Then

- (i)  $1 \cdot \mathbf{v} = \mathbf{v}$  for all  $\mathbf{v} \in \mathbb{F}^n$ .
- (ii)  $(\lambda\mu) \cdot \mathbf{v} = \lambda \cdot (\mu \cdot \mathbf{v})$  for all  $\lambda, \mu \in \mathbb{F}^\times, \mathbf{v} \in \mathbb{F}^n$ .



**Example** (Conjugation Action on the Group Itself). Let  $G$  be any group, and consider  $X = G$ . Define an action of  $G$  on itself by conjugation:

$$G \curvearrowright G, \quad (g, x) \mapsto g \cdot x := g * x * g^{-1}.$$

Then

(i)  $e \cdot x = e * x * e^{-1} = x$  for all  $x \in G$ .

(ii) Note that

$$\begin{aligned} (g * h) \cdot x &= (g * h) * x * (g * h)^{-1} \\ &= (g * h) * x * (h^{-1} * g^{-1}) \\ &= g * (h * x * h^{-1}) * g^{-1} \\ &= g * (h \cdot x) * g^{-1} \\ &= g \cdot (h \cdot x). \end{aligned}$$

Thus, this is a left group action.

**Example** (Trivial  $G$ -Set). Let  $G$  be any group and define the set  $X = \{x\}$ , a singleton. Define the action

$$G \curvearrowright X, \quad (g, x) \mapsto g \cdot x := x \quad \text{for all } g \in G.$$

This is the **trivial action**, where every group element acts as the identity on  $X$ :

(i)  $e \cdot x = x$ .

(ii)  $(g * h) \cdot x = x = g \cdot (h \cdot x)$ .

**Example** (Action on Coset Space  $G/H$ ). Let  $(G, *)$  be a group, and let  $H \leq G$ . Let  $X = G/H$  be the set of left cosets of  $H$  in  $G$ , i.e.,

$$X = G/H = \{gH \mid g \in G\}.$$

Define an action

$$G \curvearrowright G/H, \quad (g, aH) \mapsto (ga)H.$$

This is well-defined because if  $a_1H = a_2H$ , then  $a_1^{-1}a_2 \in H$ , so:  $ga_1H = ga_2H$ . Since

(i)  $e \cdot aH = aH$ ;

(ii)  $(gh) \cdot aH = g \cdot (h \cdot aH)$ ,

this is a **transitive action**.

### Group Elements Act as Permutations

**Proposition.** Let  $G$  be a group action on a set  $X$  via a left action  $G \curvearrowright X$ , given by  $(g, x) \mapsto g \cdot x$ . Then for each  $g \in G$ , the map

$$\sigma_g : X \rightarrow X, \quad x \mapsto g \cdot x$$

is one-to-one and onto. That is,  $\sigma_g \in \text{Sym}(X)$ , the group of all permutations of  $X$ .

Proof. TBA

□

### Group Actions Induce Permutation Representations

**Theorem.** Let  $G$  be a group action on a set  $X$  via a left group action  $G \curvearrowright X$ ,  $(g, x) \mapsto g \cdot x$ . For each  $g \in G$ , define the bijection  $\sigma_g : X \rightarrow X$  by  $\sigma_g(x) := g \cdot x$ . Then the map

$$\phi : G \rightarrow \text{Sym}(X), \quad g \mapsto \sigma_g,$$

is a **group homomorphism** from  $G$  to the symmetric group  $\text{Sym}(X)$ . In other words, for all  $g, h \in G$ ,

$$\phi(g * h) = \sigma_{g * h} = \sigma_g \circ \sigma_h = \phi(g) \circ \phi(h).$$

**Remark.** A group action  $G \curvearrowright X$  is equivalent to a group homomorphism  $G \rightarrow \text{Sym}(X)$ , i.e., a **permutation representation** of  $G$ .

Proof. TBA

□

### Cayley Theorem

**Theorem.** Let  $G$  be a group. Consider the action of  $G$  on itself by left multiplication. For each  $g \in G$ , define

$$\sigma_g : G \longrightarrow G, \quad x \mapsto g \cdot x.$$

Then the map

$$\phi : G \longrightarrow \text{Sym}(G), \quad g \mapsto \sigma_g$$

is an **injective group homomorphism** (group monomorphism). In particular,

$$\phi(G) \simeq G \quad \text{and} \quad \phi(G) \leq \text{Sym}(G).$$

Proof.

□

## Normal Subgroups

### Existence of the Quotient Group

**Proposition.** Let  $(G, *)$  be a group and let  $H \leq G$  be a subgroup. Define a binary operation  $\boxtimes$  on the set of left cosets  $G/H$  by

$$(g * H) \boxtimes (g' * H) = (g * g') * H$$

where  $g, g' \in G$ . Then this operation is well-defined (and makes  $G/H$  into a group) if and only if

$$g * h * g^{-1} \in H.$$

for all  $g \in G, h \in H$ .

*Proof.*  $(\Rightarrow)$  Let  $g \in G$  and  $h \in H$ . Then

$$h * g^{-1} \in H \implies g * H = g(h)$$

$(\Leftarrow)$

□

### Normal Subgroup

**Definition.**

## References

- [1] 수학의 즐거움, Enjoying Math. “수학 공부, 기초부터 대학원 수학까지, 23. 추상대수학 (d) 군 작용과 케일리-정리 group action and Cayley theorem” YouTube Video, 29:20. Published October 23, 2019. URL: <https://www.youtube.com/watch?v=5SQfrH83HfA&t=1040s>.
- [2] 수학의 즐거움, Enjoying Math. “수학 공부, 기초부터 대학원 수학까지, 24. 추상대수학 (e) 정규부분군의 정의 def of normal subgroups” YouTube Video, 23:00. Published October 25, 2019. URL: <https://www.youtube.com/watch?v=3UJILZr4CNo>.

## Appendices

### Orbit and Stabilizer

**Definition.** Let  $G$  be a group acting on a set  $X$  via a (left) group action:

$$G \curvearrowright X, \quad (g, x) \mapsto g \cdot x.$$

Let  $x \in X$ .

(1) The **orbit** of  $x$  under the action of  $G$  is defined by

$$\text{Orb}_G(x) := G \cdot x = \{g \cdot x \mid g \in G\} \subseteq X.$$

This is the set of all elements of  $X$  to which  $x$  can be moved by the action of  $g \in G$ .

(2) The **stabilizer subgroup** of an  $x \in X$ , also called the **isotropy subgroup** or **fixer**, is defined by

$$\text{Stab}_G(x) := \{g \in G \mid g \cdot x = x\}.$$

This is a subgroup of  $G$  consisting of all elements that fix  $x$  under the action.

**Remark.**

- The orbits form a *partition* of  $X$ ; that is,  $X$  is the disjoint union of its orbits under the action of  $G$ .